



**HAL**  
open science

## Understanding groundwater systems and their functioning through the study of stable water isotopes in a hard-rock aquifer (Maheshwaram watershed, India)

Philippe Négrel, Hélène Pauwels, Benoît Dewandel, Jean-Marie Gandolfi, Cédric Mascré, Mohamed-Salem Ahmed

### ► To cite this version:

Philippe Négrel, Hélène Pauwels, Benoît Dewandel, Jean-Marie Gandolfi, Cédric Mascré, et al.. Understanding groundwater systems and their functioning through the study of stable water isotopes in a hard-rock aquifer (Maheshwaram watershed, India). *Journal of Hydrology*, 2011, 397 (1-2), pp.55-70. 10.1016/j.jhydrol.2010.11.033 . hal-00552062

**HAL Id: hal-00552062**

**<https://brgm.hal.science/hal-00552062>**

Submitted on 5 Jan 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **UNDERSTANDING GROUNDWATER SYSTEMS AND THEIR**  
2 **FUNCTIONING THROUGH THE STUDY OF STABLE**  
3 **WATER ISOTOPES IN A HARD-ROCK AQUIFER**  
4 **(MAHESHWARAM WATERSHED, INDIA)**  
5

6 **Ph. Negrel<sup>α, 1</sup>, H. Pauwels<sup>1</sup>, B. Dewandel<sup>1, 2</sup>, J.M. Gandolfi<sup>1, 2</sup>, C. Mascré<sup>1, 2</sup>, S. Ahmed<sup>2, 3</sup>**  
7

8 <sup>1</sup> BRGM, Avenue C. Guillemin, BP 36009, 45060 Orléans Cedex 02, FRANCE.

9 [p.negrel@brgm.fr](mailto:p.negrel@brgm.fr), [h.pauwels@brgm.fr](mailto:h.pauwels@brgm.fr), [b.dewandel@brgm.fr](mailto:b.dewandel@brgm.fr), [jm.gandolfi@brgm.fr](mailto:jm.gandolfi@brgm.fr),

10 <sup>2</sup> CEFIRES, Indo-French Center for Ground water Research, NGRI, Uppal Road, 500 007 Hyderabad, India

11 <sup>3</sup> NGRI, Uppal Road, 500 007 Hyderabad, India

12 [shakeelahmed@ngri.res.in](mailto:shakeelahmed@ngri.res.in)

---

13 **Abstract**  
14

15 Groundwater degradation through abstraction, contamination, etc., shows a world-wide  
16 increase and has been of growing concern for the past decades. In this light, the stable  
17 isotopes of the water molecule ( $\delta^{18}\text{O}$  and  $\delta^2\text{H}$ ) from a hard-rock aquifer in the Maheshwaram  
18 watershed (Andhra Pradesh, India) were studied. This small watershed (53 km<sup>2</sup>) underlain by  
19 granite, is endorheic and representative of agricultural land use in India, with more than 700  
20 bore wells in use. In such a watershed, the effect of overpumping can be severe and the  
21 environmental effects of water abstraction and contamination are of vital importance. A  
22 detailed and dynamic understanding of groundwater sources and flow paths in this watershed  
23 thus is a major issue for both researchers and water managers, especially with regards to water  
24 quality as well as the delimitation of resources and long-term sustainability.

25 To this end, the input from monsoon-precipitation was monitored over two cycles, as well as  
26 measuring spatial and temporal variations in  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  in the groundwater and in  
27 precipitation. Individual recharge from the two monsoon periods was identified, leading to  
28 identification of periods during which evaporation affects groundwater quality through a  
29 higher concentration of salts and stable isotopes in the return flow. Such evaporation is further  
30 affected by land use, rice paddies having the strongest evapotranspiration.

---

<sup>α</sup> Corresponding author: [p.negrel@brgm.fr](mailto:p.negrel@brgm.fr); Tel.: +33 2 38 64 39 69, Fax: +33 2 38 64 34 46

31  
32

33 **Keywords:** Evaporation, Groundwater, India, Monsoon, Recharge, Stable Isotopes

34

## 35 **1. Introduction**

36 Groundwater flow and storage in hard-rock areas are of great interest and importance to  
37 researchers and water managers (e.g., De Silva and Weatherhead, 1997; Ballukraya and  
38 Sakthivadivel, 2002), from a viewpoint of both water quantity (Gupta and Singh, 1988) and  
39 quality (Robins and Smedley, 1994), as well as for delimiting resources and aquifers (e.g.,  
40 Singhal et al., 1988). In terms of hydrogeology, hard rocks are heterogeneous and anisotropic,  
41 groundwater flow generally being controlled by fissure networks (Maréchal et al., 2004).

42 Typically, hard-rock aquifers occupy the first tens of metres below ground (Detay et  
43 al., 1989). Their hydrodynamic properties derive primarily from weathering processes (Wyns  
44 et al., 1999; Taylor and Howard, 2000; Dewandel et al., 2006), where hostrock minerals are  
45 transformed into mainly clay-rich materials at surface equilibrium (Tardy, 1971; Nahon,  
46 1991). From the top downward, hard-rock aquifers consist of two main layers: saprolite or  
47 regolith—a clay-rich material derived from prolonged in-situ weathering of the hostrock—  
48 and a fissured layer characterized by dense sub-horizontal and -vertical fracturing in the first  
49 metres, below which fissure density decreases with depth (e.g. Dewandel et al., 2006). Such  
50 fractures either pre-existed or were caused by the swelling of certain minerals (e.g. biotite in  
51 granite), resulting in a local volume increase that generates cracks and fracturing. The fresh  
52 basement below is only locally permeable, where tectonic joints and fractures are present.

53 In India, as in other rapidly developing countries, groundwater use for domestic,  
54 industrial and agricultural activities is growing fast. Groundwater now irrigates 27 million  
55 hectares of farmland in India, a larger area than that irrigated by surface water (21 million ha).  
56 This change in water use has been extremely rapid since the start of the ‘Green Revolution’ in  
57 the 1970s; the number of bore wells has also increased enormously in the past 40 years, from  
58 less than one million in 1960 to over 26 million in 2007. For that reason, groundwater

59 resources are under great stress, especially in hard-rock and (semi)arid areas, due to the  
60 abstraction of large quantities of water through pumping for irrigation that undermines the  
61 sustainability of water availability and agricultural development. Over the last decade, much  
62 of India, and particularly Andhra Pradesh, Karnataka, Maharashtra, Madhya Pradesh and  
63 Rajasthan, has suffered from drought, severe drops in groundwater level, and an alarming  
64 deterioration of water quality. It is thus imperative that groundwater resources be carefully  
65 managed and that institutions in charge of water management be equipped with suitable tools  
66 (Dewandel et al., 2007b).

67 Sustainable use of groundwater is quantitatively related to such factors as the volume  
68 of the existing resource, recharge, and associated environmental factors. From a groundwater-  
69 quality viewpoint, sustainability implies the prevention of deterioration of groundwater  
70 quality beyond acceptable and well-defined limits.

71 Stable isotopes of the water molecule have been investigated throughout India, as a  
72 support of fluoride-release investigations (Tirumalesh et al., 2007), as a tool for recharge and  
73 water-pathway characterization (Gupta et al., 2005; Sukhija et al., 2005; Mukherjee et al.,  
74 2007), for geothermal-water investigations (Majumdar et al., 2005), and for isotopic  
75 fingerprinting of paleo-climates in groundwater (Kulkarni et al., 1995; Sukhija et al., 1998).  
76 However, the trend of increasing groundwater exploitation has induced a major use of isotope  
77 tracing over the past three decades (Kumar et al., 1982; Gupta and Deshpande, 2005;  
78 Mukherjee et al., 2007; Négrel et al., 2007; Murad and Krishnamurthy, 2008). The main  
79 contribution of stable isotopes in fractured-rock hydrogeology concerns the identification of  
80 variably mixed groundwater reservoirs, of groundwater flow, and of resource renewal  
81 (Fontes, 1980, Praamsma et al., 2009).

82 The Maheshwaram watershed is representative of watersheds in southern India in  
83 terms of geology, overpumping of its hard-rock aquifer (more than 700 classical open-end

84 wells in use), its cropping pattern (rice dominating), and its rural socio-economy mainly based  
85 on traditional agriculture (Dewandel et al., 2007a). Groundwater resources face chronic  
86 depletion, reflected in a strongly declining water table induced by a negative groundwater  
87 balance (Maréchal et al., 2006).

88 The aim of this work, based on a detailed study of the isotopic composition of  
89 groundwater and rainfall in the Maheshwaram watershed (Négrel et al., 2007), was to  
90 investigate the use of stable isotopes of the water molecule for tracing and fingerprinting the  
91 processes of groundwater-recharge (e.g. the monsoon input) and water-use in this typical rural  
92 Indian watershed experiencing agricultural water-resource overexploitation. As return flow  
93 was suspected to be an important process, a major goal of this investigation was to determine  
94 its potential impact on groundwater.

## 95 **2. General features of the catchment**

96 The 53 km<sup>2</sup> Maheshwaram watershed (Fig. 1) is located 35 km south of Hyderabad (Ranga  
97 Reddy District, State of Andhra Pradesh). The area has a relatively flat topography with  
98 elevations between 590 and 670 m above sea level and no perennial streams (Kumar and  
99 Ahmed, 2003).

100 The semi-arid climate is controlled by the periodicity of the monsoon (rainy season:  
101 June–October). Mean annual precipitation is around 750 mm, more than 90% of which falls  
102 during the monsoon season. The mean annual temperature is about 26 °C, but during summer,  
103 from March to May, maximum temperatures can reach 45 °C. The ratio of potential  
104 evaporation from soil plus transpiration by plants (1800 mm/year) against the total quantity of  
105 rainwater, yields an aridity index of 0.42, typical of semi-arid areas (Maréchal et al., 2006).  
106 Surface streams are dry most of the year, except for a few days after very heavy rainfall  
107 during the monsoon.

108 The area is underlain by Archean granite (Fig. 1; Dewandel et al., 2006). The rock is  
109 mainly orthogneissic granite with porphyritic K-feldspar. Intrusive leucocratic granite with a  
110 lower biotite content forms small hills and boulder-strewn outcrops. The weathering profile of  
111 these granites generally shows a thin layer of red soil (10–40 cm), a 1 to 3 m thick layer of  
112 sandy regolith, a 10 to 15 m thick layer of laminated saprolite, fractured granite that occupies  
113 the next 15 to 20 m and, farther down, fresh granite.

114 The groundwater budget of the depleted unconfined aquifer in the watershed can be  
115 summarized as follows. Horizontal groundwater inflow and outflow of the aquifer are very  
116 low compared to other terms (about 1 mm/year) and their difference is close to nil (Maréchal  
117 et al., 2006), baseflow representing groundwater discharge to streams or springs being nil  
118 (Kumar and Ahmed, 2003; Maréchal et al., 2006). Groundwater recharge is mainly direct as  
119 no permanent or temporary streams exist and only two tanks can serve as a source of indirect  
120 recharge (Dewandel et al., 2007b). The irrigation-return flow, according to Dewandel et al.  
121 (2007a), is variable in the area as a function of land-use (higher in rice paddies than in flower  
122 or vegetable plots) and evaporation from the water table is quasi nil (Maréchal et al., 2006).  
123 The water table for June 2002 (Maréchal et al., 2006) roughly follows the topographic slope,  
124 as is common in flat hard-rock areas. Water-table levels fluctuate between 610 and 619 masl,  
125 compared to a flat topography with elevations of 670 to 690 m. The water table is thus  
126 everywhere in the fissured aquifer layer, but local cones of depression occur in areas where  
127 groundwater abstraction is high due to strong pumping for irrigation.

### 128 **3. Sampling procedures and analytical methods**

129 About 700 open-end wells fitted with submersible pumps, with depths ranging from 30 to  
130 60 m, are used for irrigation in the Maheshwaram watershed (Maréchal et al., 2006). The  
131 water samples were collected from these operational agricultural wells. Water temperature,

132 electrical conductivity (EC) and pH of each sample were measured *in situ*, EC with a  
133 conductivity meter standardized to 20 °C and pH using a pH electrode previously calibrated  
134 with standard buffers. Samples for stable-isotope determinations were stored in 50 ml high-  
135 density polyethylene bottles with small necks and caps sealed with paraffin film. The stable  
136 isotopes <sup>2</sup>H and <sup>18</sup>O were measured using a Finnigan MAT 252 mass spectrometer with a  
137 precision of 0.1‰ vs. SMOW (Standard Mean Ocean Water) for δ<sup>18</sup>O and of 0.8‰ for δ<sup>2</sup>H.  
138 Isotopic compositions are plotted in the usual δ-scale in ‰ with reference to V-SMOW  
139 (Vienna Standard Mean Ocean Water) according to  $\delta_{\text{sample}} (\text{‰}) = \{(R_{\text{sample}} / R_{\text{standard}}) - 1\} * 1000$ ,  
140 where R is the <sup>2</sup>H/<sup>1</sup>H or <sup>18</sup>O/<sup>16</sup>O atomic ratio. Chloride analyses were performed by ion  
141 chromatography (uncertainty of 10% with a coverage factor k = 2), after filtration through  
142 0.45 mm acetate filters (Pauwels et al., 2007).

143 In January 2006, after the rainy period, we selected a network of 24 open-end wells  
144 from the over 700 existing ones in the watershed for the groundwater collection to be  
145 representative of watershed recharge. In March 2006, the same network of open-end wells  
146 was sampled plus two others. One of the rare surface waters, a tank fed by the monsoon rains  
147 and afterwards only affected by evaporation, was also sampled several times to be  
148 representative of evaporation. Further sampling concerned 23 of the open-end wells from the  
149 network in July 2006, 11 open-end wells in November 2006, and 9 open-end wells in June  
150 2008 and February 2009. During the different collection surveys, groundwater temperatures  
151 were 25.9 to 30.1 °C.

152 In addition to the groundwater surveys, monsoon rainfall was collected for isotope  
153 characterization of the rainwater input into the watershed. To this end, rainfall gauge RF  
154 (Fig. 1) was designed and installed in the meteorological station of Maheshwaram village. As  
155 the samples were collected on a monthly basis, the isotopic composition of the sample was  
156 likely to be modified by evaporation. This was minimized by a special construction of the rain



157 gauge, which included a 1.2-m-deep concrete excavation with a water depth of around 30 cm,  
158 in which a cool box was placed. A 15-cm-diameter tube connected the 5-litre collection bottle  
159 to the rain-gauge funnel. A lid for lowering the evaporation and preserving the collecting  
160 system from direct input of the monsoon sealed the excavation.

161 During this study, we tried to reproduce the effect of evaporation on the basis of pan  
162 evaporation (Jhajharia et al., 2009). Evaporated water samples were taken from a galvanized  
163 evaporation pan installed far from vegetation and about 10 cm above the ground. The  
164 operating water level in the pan at the start of the experiment was 270 mm and the volume  
165 was 109.9 L from the rim. The experiment covered 55 days between November 2 and  
166 December 27, 2007, and concerned the analysis of Cl<sup>-</sup> contents and the stable isotopes δ<sup>2</sup>H  
167 and δ<sup>18</sup>O, as described by Kattan (2008) from Syria.

## 168 **4. Results**

### 169 4.1. δ<sup>2</sup>H AND δ<sup>18</sup>O ISOTOPIC SIGNATURES OF THE MONSOON

170 The collection station in Maheshwaram village was operational from June to November 2006  
171 and six monthly samples were collected. The results are given in Table 1. The amount of  
172 water ranged from 0.25 L in November to 4 L in September 2006. In all, 11 L of rainwater  
173 were collected for 10.5 L recorded by the meteorological station. Moreover, the linear  
174 relationship (R squared = 0.996) observed for the individual points (Iso\_Rainfall gauge = 1.05  
175 x Meteo\_Rainfall gauge + 0.02; n = 7) confirms that all the monsoon rainfall was collected  
176 for the isotope survey (Table 1). The 2006 monsoon gave 517 mm of precipitation input on  
177 the watershed, which is considered as a low monsoon. The isotopes in the precipitation varied  
178 between -1.2‰ in June and -6.9‰ in October for δ<sup>18</sup>O, and between 0.2‰ and -44.4‰ for  
179 δ<sup>2</sup>H. The mean weighted isotope compositions for the precipitation are -3.83‰ and -21.69‰  
180 for δ<sup>18</sup>O and δ<sup>2</sup>H, respectively (Table 1). These values are less depleted than those recorded at

181 the nearby Hyderabad station with a mean weighted  $\delta^{18}\text{O}$  between April and November 1998  
182 of -6.5‰ (Sukhija et al., 2006), or than those from the Lower Maner Basin (Andhra Pradesh,  
183 north of Maheshwaram) where monthly composite precipitation samples in 1977 gave ranges  
184 of -4.1 up to -5.1‰ for  $\delta^{18}\text{O}$  and of -25.7 up to -33.4 for  $\delta^2\text{H}$  (Kumar et al., 1982).

185 During the main rainy season (June to September), the west coast of India receives  
186 influx from the Arabian Sea as the monsoon blows in from the southwest and is uplifted by  
187 the Western Ghats. This section of the southwest monsoon is called the “Arabian Sea branch”  
188 (hereafter referred to as SW monsoon), generally with a low depletion of heavy isotopes  
189 (Deshpande et al., 2003; Gupta et al., 2005). In the Bay of Bengal, the monsoon current then  
190 takes a south-easterly turn, entering the Indo-Gangetic plains after being replenished with  
191 moisture from the Bay of Bengal. This section of the Indian monsoon is called the “Bay of  
192 Bengal branch” (hereafter referred to as NE monsoon) generally yielding a much larger  
193 depletion of heavy isotopes (Deshpande et al., 2003; Gupta et al., 2005).

194 In Maheshwaram, almost 70% of the rainfall is contributed by the SW monsoon from  
195 June to September and the rest by the NE monsoon from October to November (Kumar and  
196 Ahmed, 2003). During the 2006 monsoon this was quite different, with 54% of the rain falling  
197 from June to August and 46% from September to November. This indicates a change in the  
198 distribution between the two monsoons, as has been observed regularly since the 1990s with  
199 mean values of 59% and 41%, respectively, for the SW and NE monsoons (data are from the  
200 meteorological station in Maheshwaram). During the collection period, the  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  fell  
201 in two groups as illustrated on Figure 2. The first group yielded slightly depleted values of -  
202 1.2 to -2.6‰ ( $\delta^{18}\text{O}$ ) and 0.2 to -11.4‰ ( $\delta^2\text{H}$ ) that represent the SW-monsoon input between  
203 June and August, while the second one, between September and November, yielded a much  
204 larger depletion of heavy isotopes with values of -5.5 to -6.9‰ ( $\delta^{18}\text{O}$ ) and -34.4 to -44.4‰  
205 ( $\delta^2\text{H}$ ), a typical range for the NE monsoon. Considering the distribution of the two monsoons,

206 the SW monsoon became less and less dominant and thus the mean value of the rainfall was  
207 less and less depleted for  $\delta^{18}\text{O}$  (around -3.83‰ for this study), compared to the previously  
208 reported values of less than -5‰ for  $\delta^{18}\text{O}$  (Kumar et al., 1982; Sukhija et al., 2006).

209 Overall, when compared to the values from the SW and NE monsoons, the mean  
210 weighted values are depleted for the NE monsoon (-43.1 and -6.43‰ for  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$ ,  
211 respectively) while they are enriched for the SW one (-3.5 and -1.6‰ for  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$ ,  
212 respectively), which agrees with previous studies (Deshpande et al., 2003; Gupta et al., 2005).  
213 The NE monsoon was found to have values of around -60.7 and -8.8‰ for  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$ ,  
214 while the SW monsoon gave -17.1 and -3.0‰ for  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$ .

215 We reproduced the monsoon collection in 2008 from June to November with a further  
216 six monthly samples. Similar to the 2006 monsoon, 12 L of rainwater was collected giving  
217 620 mm of precipitation input on the Maheshwaram watershed. The isotopes in the 2008  
218 precipitation varied in the same range as those of the 2006 monsoon (Table 1), with  $\delta^{18}\text{O}$   
219 ranging from 0.1‰ to -7.4‰ and  $\delta^2\text{H}$  from 5.6‰ to -49.7‰. The mean weighted isotope  
220 compositions for the precipitation are -3.39‰ and -17.05‰ for  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$ , respectively  
221 (Table 1). As illustrated on Figure 2, the two mean values are very close. As for the 2006  
222 monsoon (Fig. 1), the two monsoon branches can be seen on Figure 2.

223 The rainwater-isotope data collected at the Maheshwaram station define a linear  
224 relationship  $\delta^2\text{H} = 7.64 \pm 0.26 \times \delta^{18}\text{O} + 7.80 \pm 1.18$  ( $R^2 = 0.989$ ,  $n = 12$ ). We used FREML  
225 (Functional Relationship Estimation by Maximum Likelihood, AMC, 2002) to give the best  
226 estimate of the slope and intercept of the co-variation. This local meteoric water line falls  
227 between the global meteoric water line  $\delta^2\text{H} = 8\delta^{18}\text{O} + 10$  of Craig (1961) and the linear  
228 relationship  $\delta^2\text{H} = 7.6\delta^{18}\text{O} + 6.3$  given by Kumar et al. (1982). The 1997 and 1998 monsoon  
229 data from the *Global Network of Isotopes in Precipitation* (IAEA/GNIP database on the web:

230 <http://isohis.iaea.org>), the data of which define a linear relationship given as  $\delta^2\text{H} = 6.91 \times$   
231  $\delta^{18}\text{O} + 2.50$  ( $R^2 = 0.989$ ,  $n = 12$ ), differ from those obtained during this study.

232

#### 233 4.2. $\delta^2\text{H}$ AND $\delta^{18}\text{O}$ ISOTOPIC SIGNATURES OF GROUNDWATER

234 Results of isotopic analyses for the groundwater collected at different periods are given in  
235 Table 2. The isotope composition in the open-end well network shows a wide range in isotope  
236 compositions from -0.6 to -4.1‰ for  $\delta^{18}\text{O}$  and from -6.6 to -33.1‰ for  $\delta^2\text{H}$ .

237 The main causes of variations in the stable-isotope signature of groundwater are  
238 natural variations in the isotopic composition of rainfall, mixing with pre-existing waters, and  
239 evaporation during percolation through soil and/or the unsaturated zone (Kendall and  
240 McDonnell, 1998). In view of the temperatures generally encountered in the subsurface, the  
241 stable isotopes of water can be considered as conservative and not affected by exchanges with  
242 soil or rock (Barth, 2000).

243 The  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  values of the groundwater samples collected in January are plotted  
244 on Figure 2 with the local meteoric water line and its 95% uncertainty envelopes (LMWL),  
245 and the global meteoric water line (GMWL, Craig, 1961); those earlier defined by Kumar et  
246 al. (1982) plot in the 95% LMWL uncertainty envelopes and are not shown on the figure.  
247 Figure 2 shows only the data obtained in January, which represent recharge that mixed with  
248 pre-existing water. Figure 2 also shows all sampling periods (January plus March, July and  
249 November 2006, June 2008 and February 2009). Groundwater in the watershed collected in  
250 January 2006 defines a co-variation with a statistically significant correlation when tested  
251 with Pearson's parametric or Spearman's nonparametric tests for correlation with values of  
252 0.985 (at  $\alpha = 0.05$ ). Using FREML, the equation is  $\delta^2\text{H} = 6.17 \pm 0.25 \times \delta^{18}\text{O} - 0.54 \pm 0.67$  ( $n =$   
253 24). Groundwater also defines a co-variation in March, June, November 2006, June 2008 and  
254 February 2009 with Pearson and Spearman's R coefficients higher than 0.98. The equations

255 are given in Table 3 for all periods and few differences can be observed considering the 95%  
256 uncertainty envelopes.

257

#### 258 4.3. IMPACT OF EVAPORATION ON $\delta^2\text{H}$ AND $\delta^{18}\text{O}$ ISOTOPIC SIGNATURES

259 The  $\delta^2\text{H}$ – $\delta^{18}\text{O}$  relationship due to evaporation of water can be predicted on the basis of a  
260 steady-state isotope-balance model using temperature, humidity, isotopes in precipitation and  
261 a liquid-vapour equilibrium model. This is based on the Craig-Gordon model (see review in  
262 Gat, 2008). A slope of around 4 in a  $\delta^2\text{H}$ – $\delta^{18}\text{O}$  relationship is generally observed either in  
263 surface water during dry-season flow (Négrel and Lachassagne, 2000; Gibson et al., 2008), or  
264 in the prediction of the Craig-Gordon model for an open-water dominated evaporative system.  
265 Allison et al. (1984) showed a slope of between 4 and 6 in a  $\delta^2\text{H}$ – $\delta^{18}\text{O}$  relationship for the  
266 residual liquid when water evaporates from lakes and rivers or saturated soil, and of between  
267 2 and 5 when evaporation affects an unsaturated soil profile.

268 A pan evaporation experiment was carried out for 55 days between 2 November and  
269 27 December 2007. Starting with a Cl<sup>-</sup> of 4.2 mg L<sup>-1</sup>, it reached 33 mg L<sup>-1</sup> at the end of the  
270 experiment, leading to a concentration factor of 7.9. At the same time, the water volume  
271 decreased from 109.9 L at the start to reach 20.6 L at the end, i.e. a factor of 5.3. The  $\delta^2\text{H}$  and  
272  $\delta^{18}\text{O}$  started with respective values of -15.11 and -1.58‰ and reached respectively +65.06 and  
273 +17.73‰ 55 days later; this corresponds to the equation

$$274 \quad \delta^2\text{H} = 4.19 \pm 0.09 \times \delta^{18}\text{O} - 9.17 \pm 0.98 \text{ (using FREML, } R^2 = 0.998, n = 5)$$

275 reflecting an evaporation relationship with a slope of 4.19±0.09 (Fig. 3a). This value fully  
276 agrees with the evaporation slopes of about 4.7 given by Navada et al. (1999) for Rajasthan  
277 and of around 4.9 found by Kulkarni et al. (1995) in the central Indo-Gangetic plain.

278 One of the rare surface waters, a tank in the centre of the watershed, was sampled in  
279 March and September 2006 and in February 2009 to be representative of evaporation, as the

280 tank was fed by monsoon rain and afterwards was only affected by evaporation. The  $\delta^2\text{H}$   
281 and  $\delta^{18}\text{O}$  values of the tank water plot along the line defined by the pan-evaporation  
282 experiment, corroborating the effect of evaporation on this surface water (Fig. 3a).

283 Deuterium excess ('D-excess') in groundwater samples is expressed as  $\delta\text{D} - 8\delta^{18}\text{O}$ .  
284 The 'D-excess' values are  $<10$ , ranging from 1.1 to 7.5 in January, from -0.6 to 7.6 in March,  
285 from -9 to 7.5 in June, from -5.7 to 7.3 in November 2006, and from -4.5 to 7.2 in June 2008.  
286 The tank sample shows a 'D-excess' of -7.9 in March, -2.2 in November and -28.2 in  
287 February 2009. The lowest 'D-excess' values in the open-end wells are always observed for  
288 the M35 (around -9). The 'D-excess' values are plotted versus  $\delta^{18}\text{O}$  on Figure 4.

289 The water loss through evaporation in semi-arid zones is illustrated on Figure 3b  
290 where  $\delta^{18}\text{O}$  values for the pan experiment and the tank are plotted versus Cl contents. Starting  
291 from the mean monsoon (2006 and 2008), dissolved salts in the evaporation experiment  
292 concentrate by evaporation steps and the  $\delta^{18}\text{O}$ - $\delta^2\text{H}$  values have a slope reflecting kinetic  
293 fractionation; the Cl- $\delta^{18}\text{O}$  relationship should thus be positively correlated as any increase in  
294 evaporation would result in isotopic enrichment as well as in a higher Cl content. The Cl-  
295  $\delta^{18}\text{O}$  relationship defines a steep slope as less isotope variations would occur with increasing  
296 salinity (Fig. 3b). Compared to this natural evolution, the tank also integrates the input of Cl  
297 through human and agricultural activities, causing an increase in Cl content that can be quite  
298 large, as shown by the February 2006 sample.

299

## 300 **5. Discussion**

### 301 5.1. DISTRIBUTION OF RECHARGE IN SPACE

302 The  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  values of the groundwater samples collected in January 2006 lie closely  
303 around the local meteoric water line (LMWL, Fig. 2), confirming that the groundwater most

304 probably comes from present-day precipitation, though with a significant shift to the right of  
305 the LMWL. This reflects only evaporation during the percolation through soil and saprolite,  
306 as evaporation from the water table is nil (Maréchal et al., 2006) although an enrichment in  
307 heavy isotopes by irrigation-return flow is possible as well. Percolation through soil and  
308 saprolite may generate the shift to the right of the LMWL; Lee et al. (2007) showed that soil  
309 waters in a  $\delta^{18}\text{O}$ – $\delta^2\text{H}$  space plot close to the local meteoric water lines, indicating recharge  
310 from year-round precipitation and negligible evaporation, even during the hot summer season.  
311 Mathieu and Bariac (1996), studying groundwater recharge, found fast and direct infiltration  
312 through conducting fissured zones, generating  $\delta^{18}\text{O}$  values at a depth of a few metres that  
313 were close to the mean rainfall input, while slow infiltration through soil and the upper 0.5 m  
314 of weathered basement rock generated enriched  $\delta^{18}\text{O}$  values. Sami (1992), in South Africa,  
315 also found some evaporative enrichment in groundwater during recharge, as did Navada et al.  
316 (1999) and Nair et al. (2001) in Rajasthan who showed a similar  $\delta^2\text{H}$ – $\delta^{18}\text{O}$  relationship in  
317 groundwater as we did, and also found similar  $\delta^{18}\text{O}$  values at depth (>10m) as well as  
318 enriched values in the first metre of soil and saprolite.

319 The  $\delta^2\text{H}$ – $\delta^{18}\text{O}$  relationship (Fig. 2) intercepts the LMWL around  $\delta^{18}\text{O} = -5.5\text{‰}$ ,  
320 suggesting this value to be the approximate  $\delta^{18}\text{O}$  composition of rainwater. However, this  
321 value of  $-5.5\text{‰}$  disagrees with the weighted mean monsoon value of  $-3.39$  to  $-3.83\text{‰}$   
322 determined in this study, and also disagrees with that of Kumar et al. (1982) who gave a  $\delta^{18}\text{O}$   
323 ranging from  $-4.1$  to  $-5.1\text{‰}$ . Moreover, the distribution of groundwater values on Figure 2  
324 shows some points that plot around the LMWL close to the mean monsoon value.  
325 Evaporation of representative water of the mean monsoon value alone cannot be the process  
326 that causes a shift from the LMWL, considering the mean  $\delta^{18}\text{O}$  composition of the monsoon  
327 ( $-3.39$  to  $-3.83\text{‰}$ ) and the pan-evaporation experiment that led to a slope of  $4.19 \pm 0.09$ , quite  
328 different from the groundwater one ( $6.17 \pm 0.25$ ). This suggests different stages of groundwater

329 recharge with isotope signatures in close connection with those of the dual monsoon as shown  
330 also by Deshpande et al. (2003) for shallow groundwater in southern India. Thus, the first step  
331 is related to input from the SW monsoon, resulting in slightly depleted  $\delta^{18}\text{O}$  values, while the  
332 second one is related to input from the NE monsoon yielding much more depleted  $\delta^{18}\text{O}$   
333 values. The linear relationship defined in the groundwater thus reflects the mixing between  
334 the two rainfall inputs, explaining the data distribution along the  $\delta^{18}\text{O}$  line ranging from -5‰  
335 to -0.5‰. Some groundwater clearly reflects a recharge due to the SW monsoon without any  
336 significant influence of the NE one, which may reflect a saturated soil that limited rainwater  
337 percolation during the second monsoon stage. On the contrary, the few points that are more  
338 depleted when compared to mean monsoon values are more influenced by the NE monsoon.

339         The  $6.17 \pm 0.25$  slope of the linear relationship between all points in the  $\delta^2\text{H}-\delta^{18}\text{O}$  plot,  
340 differs from that of LMWL ( $7.64 \pm 0.26$ ) and is caused by a larger influence of evaporation  
341 affecting groundwater recharge during the SW monsoon. The water loss through evaporation  
342 in semi-arid areas assumes importance as the heavy-isotope content of groundwater becomes  
343 enriched relative to that of precipitation (Fontes, 1980; Navada et al., 1999; Nair et al., 2001;  
344 Kattan, 2008). This suggests that the transfer velocity from rainfall towards groundwater  
345 storage through soil and the unsaturated zone is sufficiently low, and that post-precipitation  
346 evaporation during the passage in soil and the unsaturated zone is an active process, having a  
347 greater impact for the second part of the monsoon period than for the first part, as also  
348 suggested by Navada et al. (1999) in Rajasthan, Majumdar et al. (2005) in eastern India, and  
349 Gupta et al. (2005) in central India. However, this is contrary to the observations by Stüber et  
350 al. (2003) in West Bengal and by Négrel et al. (2007) on the Subarnarekha River system  
351 (Jharkhand State).

352         This is further shown by the plot of ‘D-excess’ vs.  $\delta^{18}\text{O}$  (Fig. 4). The ‘D-excess’  
353 values in groundwater samples are less than 10 (range -0.6 to 7.6) and fluctuate together with



354 the  $\delta^{18}\text{O}$  (Pearson and Spearman's R coefficients of around -0.85). Therefore, 'D-excess'  
355 values of around 8 in groundwater may be inherited from precipitation during the monsoon as  
356 their similarity with the  $\delta^{18}\text{O}$  monsoon values confirms minor evaporation (Deshpande et al.,  
357 2003; Gupta et al., 2005). 'D-excess' values of less than 5, however, suggest significant  
358 evaporation of rainwater, leaving the residual groundwater with lower 'D-excess' values and  
359 depleted  $\delta^{18}\text{O}$ . The water in the tank plots along the same relationship, with 'D-excess' values  
360 close to the lowermost ones observed in groundwater (Fig. 4). This confirms that evaporation  
361 has a similar impact on ground- and surface waters.

362

## 363 5.2. EVOLUTION OVER TIME OF STABLE ISOTOPES IN GROUNDWATER

### 364 5.2.1. *Water dynamics at the watershed scale*

365 Groundwater in the watershed is defined by co-variations in March, June, November 2006,  
366 June 2008, and February 2009 (Pearson and Spearman's R coefficients  $>0.98$ , Fig. 2). The co-  
367 variation slope in March ( $6.44\pm 0.18$ , Table 3) fully agrees with that observed in January  
368 ( $6.17\pm 0.25$ ), and a set of slopes lower than 6 (Table 3) is observed for June 2006 ( $5.30\pm 0.19$ ),  
369 November 2006 ( $4.80\pm 0.43$ ), June 2008 ( $5.11\pm 0.30$ ) and February 2009 ( $5.16\pm 0.32$ ), though  
370 generally higher than that caused by evaporation ( $4.19\pm 0.09$ , Fig. 2). Groundwater in the  
371 watershed is extensively used for irrigation, which implies irrigation-return flow (Dewandel et  
372 al., 2007a) that may be affected by evaporation. Intensive pumping may result in mixing  
373 between enriched irrigation-return-flow water and more depleted groundwater (Basu et al.,  
374 2002; Zheng et al., 2005).

375 The rural socio-economy in the Maheshwaram watershed is based on traditional  
376 agriculture, the main crop being rice that requires much irrigation water. In semi-arid climates  
377 (Simpson et al., 1992), a major part of the water used for rice production returns to the  
378 atmosphere through the combined effects of evaporation and transpiration. Both processes  
379 cause enrichment of the residual water in heavier isotopes, evaporation leading to stronger

380 enrichment than transpiration (Clark and Fritz, 1997). Simpson et al. (1992), studying the  
381 flooding of rice paddies in Australia, found a slope of the  $\delta^2\text{H}-\delta^{18}\text{O}$  relationship for the  
382 return-flow water similar to that observed in evaporation pans in the same region. Therefore,  
383 evaporation would cause a shift for the slope of the  $\delta^2\text{H}-\delta^{18}\text{O}$  co-variation in groundwater  
384 towards the lower value by the end of the irrigation period in June and May, which would  
385 remain low up to the end of the recharge period (after the end of the monsoon in November,  
386 plus the transfer through the unsaturated zone). Maréchal et al. (2006) calculated that around  
387 40% of the pumped water returns to the aquifer, and Dewandel et al. (2007a) showed the  
388 irrigation-return-flow coefficients (ratio between pumping abstraction versus return flow) to  
389 be 10% for flowers, 25% for vegetables and around 50% for rice. This means that the effect of  
390 the return flow on the  $\delta^2\text{H}-\delta^{18}\text{O}$  signature of groundwater is variable as a function of land  
391 use. Taken as a whole over the watershed, the observed small effect on  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  in  
392 groundwater, which show about the same values as those observed in January during the  
393 collection period (Table 3), suggests that the return-flow process has a relatively minor impact  
394 on the  $\delta^2\text{H}-\delta^{18}\text{O}$  signature of groundwater even if it is important (around 80% of the aquifer  
395 volume on a yearly basis, Dewandel et al., 2006). This will be further discussed hereafter  
396 when comparing the data at the open-end well scale, as a working hypothesis could be the  
397 scale effect because land-use patterns differ in the watershed (de Condappa, 2005), with  
398 paddy fields dominating in some parts compared to grape and fruit growing in others.

399

#### 400 *5.2.2 Downscaling water dynamics: land-use effect at the bore-well scale*

401 There is some variation in the stable-isotope composition between sites, suggesting a  
402 heterogeneous origin of the water in the aquifer; this first heterogeneity is related to the  
403 variation in recharge as demonstrated before. Investigation of the water dynamics should  
404 consider specific use of a well, for which several features can be stated. First, there is a

405 variation or lack of variation in site(s) that may reflect different processes. Hereafter, we  
406 consider the case of open-end wells located in parts of the watershed with different water use.  
407 First we will investigate the effect of moderate use of groundwater for cultivation from three  
408 open-end wells. The first is the little-used open-end well M57 that supplies a dress factory,  
409 the two others are open-end wells M12 and M14, respectively used for drip irrigation of grape  
410 vines and mango trees. After that, we consider the larger use of groundwater from open-end  
411 wells M30, 35, 50 and 53, used for rice growing and vegetable- and flower cultivation.

412 Records for January, March, June and November 2006 were collected for open-end  
413 wells M12, 14 and 57 (Fig. 5) plus June 2008 and February 2009 for M12 and M57. For  
414 open-end well M14 there is no variation between the periods, and the data plot close to the  
415 meteoric water lines and the mean weighted monsoon value of 2006. This suggests that  
416 recharge is rapid and concerns the two monsoon periods, as the January values for  $\delta^{18}\text{O}$  and  
417  $\delta^2\text{H}$  reflecting recharge (Fig. 2) fall on the meteoric water lines. The groundwater is not  
418 affected by evaporation and the  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  values of the groundwater mimic those of the  
419 mean weighted monsoon, suggesting that recharge occurs during the monsoon period from  
420 June to December, with physical parameters of soil and saprolite that allow fast and direct  
421 recharge (Mathieu and Bariac, 1996). It is obvious that evaporation, either from percolation  
422 through soil and saprolite, or directly from the water table, is nil (Maréchal et al., 2006) as  
423 the  $\delta^2\text{H}-\delta^{18}\text{O}$  values in the aquifer agree with the meteoric water input. Following the  
424 recharge period, the other surveys show similar  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  values, suggesting that the  
425 water body supplying the water in open-end well M14 is the same throughout the period,  
426 leading to the conclusion that no return flow occurs during drip irrigation of mango trees.

427 Open-end well M57 functioned similarly to open-end well M12 for the first five  
428 periods, without variation in the  $\delta^2\text{H}-\delta^{18}\text{O}$  values between the first four periods whose data  
429 plot close to the meteoric water lines and the mean weighted monsoon value of 2006. This

430 agrees with recharge during the June to December monsoon period, with soil and saprolite  
431 characteristics that allow fast and direct recharge to be dominant. The groundwater around  
432 open-end well M57 supplies a dress factory and a recent increase in the pumping frequency  
433 has shown a shift of the last collected sample (February 2009) along the evaporation line. The  
434 response of the aquifer to this change in water abstraction is relatively rapid as the time  
435 interval is around 8 months (between June 2008 and February 2009).

436 Open-end well M12 functions differently. Even though the three periods of January,  
437 March and June 2006 had similar  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  values, they all plot along the evaporation  
438 line. For November 2006, after the recharge period, the groundwater values are close to those  
439 of the mean monsoon, again suggesting rapid recharge (Mathieu and Bariac, 1996) without  
440 any impact from evaporation. In June 2008, the value was close to that of November 2006,  
441 but plotted differently to those observed in June 2006 and February 2009, plotting close to  
442 November 2006 and June 2008 values but different from the March 2006 one. This suggests  
443 that recharge was sufficiently minor to balance the enriched  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  values observed  
444 between January and June 2006, but was strong enough to affect the groundwater  $\delta^{18}\text{O}$  and  
445  $\delta^2\text{H}$  values between November and February 2009. But, as the  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  values in  
446 groundwater differ from the meteoric water lines through evaporation during the period  
447 between January and June 2006, this may indicate that the water body from which the water is  
448 pumped was affected by returned irrigation flow. On the other hand, the water body seems to  
449 be less affected by evaporation during the period from November 2008 to February 2009,  
450 suggesting less water use in this area where grape vines are drip irrigated. A rice paddy is  
451 located close to the open-end well, affecting the groundwater body during the rice-growing  
452 season (not all the time) through a partly identified return irrigation flow.

453 We now consider open-end wells M30, 35, 50 and 53, used for rice growing and  
454 vegetable- and flower cultivation. Data for all five periods were collected for open-end wells

455 M50, 53 and 30, and for four periods for open-end well M35 (Fig. 5, cont'd). All wells are  
456 located near paddy fields and used throughout the irrigation period for flooding the paddies  
457 twice a day, all points evolving along the evaporation line. This agrees with groundwater  
458 pumping during the irrigation period from the same water body, any compositional change  
459 being only controlled by returned irrigation flow. The January values for the open-end wells  
460 reveal that all were affected by evaporation as the values depart from the meteoric water lines,  
461 but all are also different from the mean weighted 2006 monsoon value, suggesting once again  
462 that fast and direct recharge is dominant. Moreover, they evolve similarly during the irrigation  
463 period. The groundwater from well M53 falls on the evaporation line, with the March and  
464 June 2006 points being close to that of the January recharge, indicating abstraction from the  
465 same water body during the irrigation period. The points collected in November 2006 and  
466 February 2009 show a slightly more evaporated signature, probably reflecting a returned  
467 irrigation flow rather than a change in the water origin, which was also observed in June  
468 2008. The groundwater from well M50 evolved similarly from recharge in January plotting  
469 along the evaporation line up to March, and June samples that migrated along the evaporation  
470 line. One key observation concerns the location along the evaporation line of the February  
471 2009 samples that plot with the most strongly shifted values, indicating that either the return  
472 flow was large at that time, or that the monsoon recharge was not yet visible in the  $\delta^{18}\text{O}$  and  
473  $\delta^2\text{H}$  signatures of the groundwater. For well M35 the January and March samples do not  
474 differ in signature, but evaporation strongly affected the June ones resulting in a heavy-  
475 isotope enrichment of the water. Such evidence of evaporation is not linked to processes  
476 affecting the water table and the only way to explain such evaporation is by large returned  
477 irrigation flow after enrichment of the water through evaporation in the paddy fields and  
478 transpiration from the rice (Simpson et al., 1992). The June samples for both 2006 and 2008  
479 thus represent a clear mixture of the groundwater that was used for irrigation between January

480 and March with the strongly evaporated returned irrigation flow. The November sample lies  
481 on the same evaporation line as the other samples of the same well, but with  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$   
482 values between those of June and March, showing a shift back to the monsoon value along the  
483 LMWL corresponding to the end of the 2007 monsoon. This implies that the groundwater  
484 collected in November was partly controlled by the signature of the (mostly evaporated)  
485 return flow and by the recharge signature of the SW Monsoon. For February 2009, similar to  
486 well M50,  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  plot with the most shifted values, again indicating that the  
487 groundwater is largely controlled by the signature of the evaporated return flow. Considering  
488 that the February sample still includes part of the 2008 monsoon, the  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  values  
489 indicate a shift back to the monsoon value and thus that the maximum of the  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$   
490 signatures of the groundwater affected by return flow may be higher.

491 In case of such binary mixing, the following equation applies:

$$492 \quad M = a \text{ EM}_1 + (1 - a) \text{ EM}_2 \quad (\text{eq. 1})$$

493 where M corresponds to the mixture,  $\text{EM}_1$  and  $\text{EM}_2$  to the two end-members, and  $a$  being the  
494 mixing parameter, i.e. the proportion of  $\text{EM}_1$  and  $\text{EM}_2$  in the mixing. This gives for  $a$

$$495 \quad a = (\text{M} - \text{EM}_2) / (\text{EM}_1 - \text{EM}_2) \quad (\text{eq. 2})$$

496 Considering the November sample as the mixture in M50, the June samples as  $\text{EM}_1$   
497 and the January one as  $\text{EM}_2$ , mixing parameter  $a$  can be calculated by using equation 2. The  
498 result shows that around  $70\% \pm 0.5$  (propagated error according to classical formulae of  
499 equations 2 derivatives) of the water in the November sample comes from  $\text{EM}_1$  (June  
500 samples) and around  $30\% \pm 0.1$  (propagated error) from  $\text{EM}_2$  (January recharge). Parameter  
501  $a$  suggests that most of the recharge through soil and the unsaturated zone occurs between  
502 November and January. For the M35 case, we can also consider the November sample as the  
503 mixture and the January sample as the recharge ( $\text{EM}_2$ ), defining the returned flow as the most  
504 shifted value with  $\delta^{18}\text{O}$  around  $0.5\text{‰}$ . Using equation 2, mixing parameter  $a$  gives around

505 50% for each end-member, showing that groundwater like that of November can be affected  
506 for around  $65\% \pm 3$  (propagated error) by return flow, agreeing with the estimate given by  
507 Dewandel et al. (2007a) for the irrigation return-flow coefficients of around 50% for rice.

508 Open-end well M30 is characterized by a more marked NE-monsoon recharge with  
509 values close to the mean weighted monsoon value. It experienced evaporation as the recharge  
510 values (January sample) depart from the meteoric water lines and the March and June 2006  
511 samples plot along an evaporation line, both values being similar but with a more marked  
512 evaporation signal compared to the January sample. The June and November 2008 and  
513 February 2009  $\delta^2\text{H}$  and  $\delta^{18}\text{O}$  values are increasingly marked by the evaporation signal, though  
514 with a less evaporated signal than that found in well M35. The influence of returned flow is  
515 strongly suggested. Considering the November sample as the mixture, the February samples  
516 as  $\text{EM}_1$  and the January one as  $\text{EM}_2$ , mixing parameter  $a$  can be calculated with equation 2  
517 showing that around  $80\% \pm 12$  (propagated error) of the water in the November sample comes  
518 from  $\text{EM}_1$  (February sample) and around  $20\% \pm 4$  (propagated error) from  $\text{EM}_2$  (January  
519 recharge). Here, parameter  $a$  suggests that groundwater like the November one can be  
520 affected around  $80 \pm 12\%$  by return flow, a higher value than the 50% estimate of Dewandel  
521 et al. (2007a) on the irrigation return-flow coefficients for rice.

522 The M35 and M30 calculations led to the conclusion that isotopes of the water  
523 molecule, at the bore-well scale, may help in quantifying the land-use effect, i.e. return flow.

524

### 525 *5.2.3 Effect of land use: impact on salinity and stable-isotope signature of the groundwater*

526

527 In order to evaluate the source of salt in groundwater at the watershed scale, chloride  
528 concentrations were plotted against the  $\delta^{18}\text{O}$  signature of the groundwater (Fig. 6). As the  
529 bedrock contains no evaporites, chloride concentrations in groundwater do not derive from  
530 weathering and must originate from both evaporation of rainwater and human activity

531 (Négrel, 1999; 2006). Data were plotted with those from Sukhija et al. (2006), obtained in  
532 weathered/fractured granite near Hyderabad (India), 30 km north of Maheshwaram. The  
533 highest  $\text{Cl}^-$  concentration in rainwater,  $(\text{Cl})_{\text{ref}}$ , was determined according to the method  
534 defined by Négrel et al. (1993); it was calculated with the chloride content in rainwater, a  
535 mean weighted value of close to  $1.9 \text{ mg L}^{-1}$  during the monsoon, multiplied by the  
536 concentration factor  $F$  (Négrel, 1999). This factor  $F$  represents the concentration effect of  
537 evapotranspiration and is related to the total quantity of rainwater  $P$  (in mm) and the  
538 evapotranspiration process  $E$  (in mm) by the equation  $F = P/(P-E)$ . According to the data  
539 given by Maréchal et al. (2006) for the watershed,  $F$  ranges between 4.5 and 11, leading to a  
540 maximum of rain-derived  $\text{Cl}$  of 8 to  $20 \text{ mg L}^{-1}$ . The lowest chloride concentrations observed  
541 in Maheshwaram groundwater are  $<5 \text{ mg L}^{-1}$  (Pauwels et al., 2007) and thus only some  
542 samples can be considered as representative of the natural background by rain input without  
543 human influence. However, the samples with the highest chloride concentrations are also  
544 those with the highest nitrate concentrations, suggesting a common origin for these two  
545 elements (domestic sewage, *in-situ* sanitation, livestock farming, fertilizer spreading). Some  
546 groundwater with a  $\delta^{18}\text{O}$  signature similar to that of the mean monsoon rain input, i.e. around  
547  $-3.5$  to  $-4\text{‰}$ , shows a large variation in  $\text{Cl}$  content (Fig. 6), from less than  $\text{Cl}_{\text{ref}}$  with a  
548 minimum of  $2.6 \text{ mg L}^{-1}$  up to  $250 \text{ mg L}^{-1}$ , whereas with a more enriched  $\delta^{18}\text{O}$  signature of  
549 around  $-3\text{‰}$  the  $\text{Cl}$  content is  $>400 \text{ mg L}^{-1}$ . This could be due to several steps of evaporation  
550 year after year. However, if mixing occurs, due to groundwater exploitation for example, a  
551 poor correlation might be observed (Kattan, 2008).

552 On Figure 6, according to the pan-evaporation experiment, the line where groundwater  
553 variations are due only to recharge (i.e. monsoon input, Fig. 3a, b) and evaporation  
554 (percolation through soil and saprolite), should show a weak impact on the  $\delta^{18}\text{O}$  and  $\text{Cl}^-$   
555 contents, as it starts from the mean rain input but never reaches the maximum of rain-derived



556  $\text{Cl}^-$  of up to  $20 \text{ mg L}^{-1}$ . Thus, considering one-step evaporation processes, the  $\text{Cl}^-$  content  
557 should only reach a low value, around  $5 \text{ mg. L}^{-1}$ .

558 In order better to understand the joint evolution of  $\text{Cl}^-$  and  $\delta^{18}\text{O}$ , we created a daily  
559 mass-balance model covering the 20-year period of overpumping of the Maheshwaram  
560 watershed, which started close to the beginning of the Green Revolution. Overexploitation  
561 can be seen in the strongly declining water table caused by a negative groundwater balance  
562 (Maréchal et al., 2006). Starting with about 10 open-end wells in 1985, the watershed  
563 contained more than 150 open-end wells in 1995 and up to 700 in 2005. For  $\text{Cl}$ , rainfall is the  
564 only source to be considered in the model as man-induced  $\text{Cl}$  inputs do not influence the O-  
565 isotopes. Furthermore, no  $\text{Cl}$  sink is considered as overpumping of the groundwater leads to  
566 endorheic functioning of the watershed (Maréchal et al., 2006) and groundwater becomes  
567 increasingly salty though without reaching oversaturation with respect to  $\text{Cl}$ -solid phases  
568 (Pauwels et al., 2007). Soils in the watershed cannot act as  $\text{Cl}$  sinks as they are clayey/sandy  
569 without an organic layer (de Condappa, 2005).

570 The model considers a constant mean value for rainfall and an identical recharge every  
571 year. For rainfall values, we used the mean weighted value of  $\text{Cl}$  during the monsoon period  
572 ( $1.9 \text{ mg L}^{-1}$ ), and a mean weighted  $\delta^{18}\text{O}$  value was used for the June-August period and  
573 another for September-November, representing the dual monsoon (Deshpande et al., 2003).

574 The volumes of recharge, pumping and return flow for the simulation are from  
575 Dewandel et al. (2007). Even if the volume of water in the aquifer varies within a year, it  
576 returns to the same initial value by the end of every year. The initial condition was estimated  
577 and concerns the mean piezometric elevation of the water table (617 m on average according  
578 our field data over 2002, 2003, 2004; see also Maréchal et al., 2006), the mean bedrock  
579 elevation (604 m), and a specific yield of 0.014 (Dewandel et al., 2006, 2007a). The aquifer  
580 volume can be calculated and thus for one aquifer volume the rainfall volume (recharge) is

581 0.5, the volume of the aquifer pumped is 0.96 and the return flow is 0.48 of the pumped  
582 volume. Recharge occurs in this model over 120 days. One should also consider the twice  
583 yearly rice harvest (Dewandel et al., 2007a). The first rainy season between June and October,  
584 Kharif, has an average duration of 137 days when watering is from rainfall and irrigation, the  
585 second period between November and April, Rabi, covering 166 days when watering is from  
586 irrigation only (Perrin et al., 2006). We also considered a watering period of one month before  
587 planting when the pumped volume is 0.1 the normal aquifer volume (Dewandel et al., 2007a).

588 The isotopic signature of the aquifer  $\delta^{18}\text{O}_T$  can be calculated as follows:

$$589 \quad \delta^{18}\text{O}_T = \delta^{18}\text{O}_{j-1} \times V_{j-1}/V_T + \delta^{18}\text{O}_{\text{RF}} \times V_{\text{RF}}/V_T - \delta^{18}\text{O}_p \times V_p/V_T + \delta^{18}\text{O}_{\text{Rd}} \times V_{\text{Rd}}/V_T \quad (\text{Eq. 3})$$

590 where:

- 591 •  $\delta^{18}\text{O}_{\text{RF}}$  is the isotopic signature of the return flow corresponding to  $\delta^{18}\text{O}_p + a$  ( $\delta^{18}\text{O}_p$  being  
592 the isotopic signature of the pumped water, and  $a$  the daily enrichment in heavy isotopes,  
593 assumed from an evaporation pan with a value of 0.35);
- 594 •  $\delta^{18}\text{O}_{\text{Rd}}$  is the isotopic signature of rainfall, for which two  $\delta^{18}\text{O}$  values were used (-1.61‰  
595 for June-August and -6.43 ‰ for September-November);
- 596 •  $V_{\text{RF}}$  is the volume of return flow, daily calculated with the equation  $V_{\text{RF}} = 0.48 \times V_p$ ;
- 597 •  $V_p$  is the pumped volume, daily calculated according to  $V_p = 0.96 \times V_{T-1}/\text{days of pumping}$   
598 (166 for the Kharif period and 137 for the Rabi period);
- 599 •  $V_{\text{Rd}}$  is the volume of direct recharge, daily calculated according to  $V_{\text{Rd}} = 0.5 \times V_p/120$ ;
- 600 •  $V_T$  is the total volume according to  $V_T = V_{T-1} + V_{\text{RF}70} + V_{\text{Rd}70} - V_p$  (the volume of return  
601 flow and direct recharge are considered over 70 days after the beginning of pumping and  
602 rainfall, and  $V_{T-1}$  is the aquifer volume the day before).

603 A similar equation was used for simulating the Cl-content variations:

$$604 \quad \text{Cl}_T = \text{Cl}_{j-1} \times V_{j-1}/V_T + \text{Cl}_{\text{RF}} \times V_{\text{RF}}/V_T - \text{Cl}_p \times V_p/V_T + \text{Cl}_{\text{Rd}} \times V_{\text{Rd}}/V_T \quad (\text{Eq. 4})$$

605

606 The results of the simulation are presented in Figure 7, the annual changes covering 10, 15  
607 and 20 years. This allows evaluating both intra-annual and inter-annual variations. From these  
608 changes, we can state, first, that the chloride content in the groundwater increases as a time  
609 function over 10 to 20 years of aquifer exploitation, while, second, the  $\delta^{18}\text{O}$  signature does  
610 not follow the same evolution and stays within the same range of -3 to -3.5‰ over 10 to 20  
611 years of aquifer exploitation. Propagated errors on the  $\delta^{18}\text{O}$ , according to classical formulae of  
612 equations 3 derivative agree with the stability of the signal as the values of the error never  
613 exceed 0.4‰. After 15 years of aquifer overpumping, the simulation results match some of  
614 the observed chloride contents (e.g. around 26 mg L<sup>-1</sup>) for the groundwater as do the 20-year  
615 simulation results with a chloride content of around 34 mg L<sup>-1</sup>. It is worth noting that the  
616 propagated errors on the Cl content, calculated similarly to that of eq.3, never exceed 1 mg L<sup>-1</sup>  
617 and thus the increase of the Cl content as given by the model is not impacted by the  
618 uncertainties on the calculation. The simulation after the maximum of groundwater  
619 exploitation of 20 years thus yields a Cl content of up to 34 mg L<sup>-1</sup>, including recharge,  
620 evaporation, pumping and return flow, but the maximum chloride content does not correspond  
621 to the highest  $\delta^{18}\text{O}$  signature as illustrated on Figure 8 for the 10<sup>th</sup> year of aquifer exploitation.  
622 The chloride content mainly follows the aquifer-volume fluctuation, whereas the  $\delta^{18}\text{O}$   
623 variations are due to the effects of three processes, i.e. pumping, evaporation and return flow,  
624 during the Rabi season. However, the increase of the  $\delta^{18}\text{O}$  value in the simulation is larger  
625 with the direct recharge from the first monsoon. The second monsoon has an opposite effect,  
626 the isotopic groundwater signature being more negative. Nevertheless, even if the  $\delta^{18}\text{O}$   
627 signature is marked by the monsoon, the simulation again leads to the conclusion that  
628 evaporation causes enrichment in heavy isotopes. The mean  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  signatures are  
629 higher for the rice-paddy areas than for the other parts of the watershed, clearly showing that  
630 land-use (paddy fields, grapes, fruit, etc.), and management of the groundwater resource

631 directly affect the hydrochemistry and isotope signatures ( $\delta^{18}\text{O}$ ,  $\delta^2\text{H}$ ) of the groundwater,  
632 notwithstanding the effect of additional anthropogenic inputs, like fertilizer applications, on  
633 the hydrochemistry.

634 The high  $\text{Cl}^-$  contents of  $<430 \text{ mg. L}^{-1}$  (sample M20, Jan. 2006) and  $370 \text{ mg. L}^{-1}$   
635 (sample M15, Nov. 2006) show a slight deviation from the mean weighted monsoon value in  
636 terms of  $\delta^{18}\text{O}$ , suggesting a very small impact of the evaporation process and a large  
637 anthropogenic influence that may arise from different sources but cannot be simulated by the  
638 model (Figs. 6 and 7). In terms of input,  $\text{Cl}^-$  and  $\text{NO}_3^-$  reach very high contents of around  
639  $940 \text{ mg. L}^{-1}$  for Cl (Pauwels et al., 2007) as shown on Figure 6. The data from Sukhija et al.  
640 (2006) show a similar relationship, with most “modern” water being  $\text{Cl}^-$  enriched, owing to  
641 the fact that one sample of “old” (a few hundreds to thousands of years old) water had a high  
642  $\text{Cl}^-$  content without any divergence of the  $\delta^{18}\text{O}$  value. When nitrates and chlorides are of  
643 organic origin, the chemical composition is marked by a lower  $\text{NO}_3^-/\text{Cl}^-$  ratio than when they  
644 have a mineral source (fertilizer). This variability of the  $\text{NO}_3^-/\text{Cl}^-$  ratio suggests a variety of  
645 inputs, which agrees with land use in the basin. Poultry farming and a population of 15,000  
646 inhabitants provide organic compounds, whereas mineral fertilizer is used in paddies and  
647 vegetable gardens.

## 648 **6. Conclusions**

649 Stable-isotope analyses of precipitation and groundwater from a tropical hard-rock aquifer  
650 (Maheshwaram watershed, Andhra Pradesh, India) defined a detailed and dynamic picture of  
651 groundwater sources and flow paths. This endorheic watershed is representative of South  
652 Indian conditions in terms of geology (hard rock), land-use, overpumping of the hard-rock  
653 aquifer (over 700 bored open-end wells in use), and its socio-economic context. The stable  
654 isotopes revealed the following features.

655           Precipitation samples from two entire monsoon cycles allowed defining the NE and  
656 SW monsoon signals, the latter being enriched in stable isotopes compared to the first. The  
657 monsoon samples define a Local Meteoric Water Line (LMWL) that is close to the Global  
658 Meteoric Water Line (GMWL), and allow calculating the mean weighted monsoon signal.

659           Groundwater data collected in January allowed defining the recharge isotopic  
660 signature in the watershed that fluctuates between the two monsoons, with a shift from the  
661 LMWL due to evaporation during percolation through soil and saprolite. Groundwater from  
662 the irrigation period (March and June) plots in the same range as that defined in January,  
663 reflecting few changes at the watershed scale.

664           Using individual open-end wells for investigating the downscaling of water dynamics,  
665 shows that, for a moderate groundwater use for fruit- and grape growing and light industry,  
666 the data plot either close to the meteoric water lines and the mean weighted monsoon value,  
667 or on an evaporation line. The first observation suggests that recharge is rapid as the monsoon  
668 signal is not transformed during percolation through soil and saprolite, while the second  
669 observation suggests a lower recharge rate.

670           Open-end wells with a heavier use of groundwater for rice paddies and vegetable- and  
671 flower growing show a different functioning. Groundwater generally follows the evaporation  
672 line, with or without deviation from the mean weighted monsoon data, and a returned  
673 irrigation flow with a more evaporated signature can be identified while a mixing model  
674 allows calculating the proportion of returned water in one case.

675           Future work should consider variations in the hydrogeological system (layered,  
676 fractured, etc.) and in groundwater collected from different boreholes at various depths, for  
677 designing a 3D-isotopic framework. This would be necessary for identifying possible  
678 flowpaths in conjunction with a larger exploitation of the groundwater resource. Such work  
679 would also aid in generalizing the use of other isotope tools such as Nd and Sr or Pb, and of

680 newly developed isotope systematics like Ca or Si, in other catchments that may face similar  
681 structural problems of groundwater overexploitation. This is of primary importance as  
682 groundwater flow and -storage in hard-rock areas are major issues for researchers and water  
683 managers, especially with regard to water quality as well as the definition of resources and  
684 aquifers and of long-term sustainability.

685

686 **Acknowledgments.** This study was carried out at the Indo-French Centre for Groundwater Research (BRGM-  
687 NGRI). The authors thank the French Ministry of Foreign Affairs and the French Embassy in India for their  
688 support. This paper has benefited from the research and technical assistance provided by Mohamed Wajiduddin  
689 (“fabuleux Wajid”), S. Roy and F. Jouin, and from the collaboration of C. Flehoc (BRGM Mass-Spectrometer  
690 team) who made the stable-isotope analyses. Tom Bullen is warmly thanked for his comments on the first  
691 version of this paper. We are grateful to Dr. H.M. Kluijver for proofreading and editing the English text. We  
692 thank the anonymous reviewers for providing critical comments that improved this manuscript.

## 693 **References**

694

695 Allison, G.B., Barner, C.J., Hughes, M.W., Leaney, F.W.J. 1984. Effect of climate and vegetation on  $\delta^{18}\text{O}$  and  
696  $\delta\text{D}$  profiles in soils. *In*: Proceedings of IAEA Symposium, Vienna 1984, 105-123.

697

698 AMC, 2002. Analytical Methods Committee. Technical brief No. 10. "Fitting a linear functional relationship to  
699 data with error on both variables" (Royal Society of Chemistry).

700

701 Barth, S.R. 2000. Stable isotope geochemistry of sediments-hosted groundwater from a Late-Proterozoic-Early  
702 Mesozoic section in Central Europe. *Journal of Hydrology* 235, 72-87.

703

704 Ballukraya, P.N., Sakthivadivel, R. 2002. Over exploitation and artificial recharge of hard rock aquifers of South  
705 India: issues and options. IWMI-Tata Water Policy Research program, International Water Management  
706 Institute, 1-14, available at <http://www.iwmi.org/iwmi-tata>.

707

708 Basu, A.R., Jacobsen, S.B., Poreda, R.J., Dowling, C.B., Aggarwal, P.K. 2002. Response to Harvey, 2002:  
709 groundwater flow in the Ganges delta. *Science* 296, 1563.

710

711 Clark, I., Fritz, P. 1997. *Environmental Isotopes in Hydrogeology*. Lewis Publishers, Boca Raton, New York,  
712 328p.

713

714 Craig, H. 1961, Isotopic variations in meteoric waters. *Science*, 133, 1702.

715

716 de Condappa, D. 2005. Étude de l'écoulement d'eau à travers la Zone Non-Saturée des aquifères de socle à  
717 l'échelle spatiale du bassin versant. Application à l'évaluation de la recharge au sein du bassin versant de  
718 Maheshwaram, Andhra Pradesh, Inde. PhD thesis, University of Grenoble, 361p.

719

720 Deshpande, R.D., Bhattacharya, S.K., Jani, R.A., Gupta, S.K. 2003. Distribution of oxygen and hydrogen  
721 isotopes in shallow ground waters from Southern India: influence of a dual monsoon system. *Journal of*  
722 *Hydrology* 271, 226-239.

723

724 De Silva, C.S., Weatherhead, E.K. 1997. Optimising the dimensions of agrowells in hard-rock aquifers in Sri  
725 Lanka. *Agricultural Water Management* 33, 117-126.  
726  
727 Detay, M., Poyet, P., Emsellem, Y. Bernardi, A. Aubrac, G. 1989. Development of the saprolite reservoir and its  
728 state of saturation: influence on the hydrodynamic characteristics of drillings in crystalline basement (in French).  
729 *Compt. Rend. Acad. Sci. Paris II* 309, 429-436.  
730  
731 Dewandel, B., Lachassagne, P., Wyns, R., Maréchal, J.C., Krishnamurthy, N.S. 2006. A generalized 3-D  
732 geological and hydrogeological conceptual model of granite aquifers controlled by single or multiphase  
733 weathering. *Journal of Hydrology* 330, 260-284.  
734  
735 Dewandel, B., Gandolfi, J.M., de Condappa, D., Ahmed, S. 2007a. An efficient methodology for estimating  
736 irrigation return flow coefficients of irrigated crops at watershed and seasonal scale. *Hydrological Processes*,  
737 DOI: 10.1002/hyp.6738.  
738  
739 Dewandel, B., Gandolfi, J.M., Ahmed, S. Subrahmanyam, K. 2007b. A Decision Support Tool for sustainable  
740 Groundwater Management in semi-arid hard-rock areas with variable agro-climatic scenarios. *Current Sciences*  
741 92, 1093-1102.  
742  
743 Fontes, J.Ch. 1980. Environmental isotopes in groundwater hydrology. *Handbook of environmental isotopes*  
744 *geochemistry*, Fritz P. and Fontes, J.Ch. (eds.), Elsevier, Amsterdam, Oxford, New York, 75-140.  
745  
746 Gat J.R. 2008. The isotopic composition of evaporating waters - review of the historical evolution leading up to  
747 the Craig-Gordon model. *Isotopes in Environmental and Health Studies* 44, 5-9.  
748  
749 Gibson, J.J., Sadek, M.A., Stone, D.J.M., Hughes, C.E., Hankin, S., Cendon, D.I., Hollins, S.E. 2008.  
750 Evaporative isotope enrichment as a constraint on reach water balance along a dryland river. *Isotopes in*  
751 *Environmental and Health Studies* 44, 83-98.  
752  
753 Gupta, C.P., Singh, V.S. 1988. Flow regime associated with partially penetrating large-diameter wells in hard  
754 rocks. *Journal of Hydrology* 103, 209-217.  
755  
756 Gupta, S.K., Deshpande, R.D. 2005. Groundwater isotopic investigations in India: What has been learned?  
757 *Current Science* 89, 825-835.  
758  
759 Gupta, S.K., Deshpande, R.D., Bhattacharya, S.K., Jani, R.A. 2005. Groundwater  $\delta^{18}\text{O}$  and  $\delta\text{D}$  from central  
760 Indian Peninsula: influence of the Arabian Sea and the Bay of Bengal branches of the summer monsoon. *Journal*  
761 *of Hydrology* 303, 38-55.  
762  
763 Jhajharia, D., Shrivastava, S.K., Sarkar, D., Sarkar, S. 2009. Temporal characteristics of pan evaporation trends  
764 under the humid conditions of northeast India. *Agricultural and Forest Meteorology* 149, 763-770.  
765  
766 Kattan, Z. 2008. Estimation of evaporation and irrigation return flow in arid zones using stable isotope ratios and  
767 chloride mass-balance analysis: Case of the Euphrates River, Syria. *Journal of Arid Environments* 72, 730-747.  
768  
769 Kendall, C., McDonnell, J.J. 1998. *Isotope Tracers in Watershed Hydrology*. Elsevier, 839p.  
770  
771 Kulkarni, K.M., Navada, S.V., Rao, S.M., Nair, A.R., Kulkarni, U.P., Sharma, S. 1995. Effect of the Holocene  
772 climate on composition of groundwater in parts of Haryana, India. IAEA-SM-336/36. In *Proceedings of the*  
773 *Symposium on isotopes in water resources management*, Vienna, March 1995. 439-454.  
774  
775 Kumar, B., Athavale, R.N., Sahay, K.S.N. 1982. Stable isotope geohydrology of the Lower Maner Basin,  
776 Andhra Pradesh, India, *Journal of Hydrology* 59, 315-330.  
777  
778 Kumar, D., Ahmed, S. 2003. Seasonal behaviour of spatial variability of groundwater level in a granitic aquifer  
779 in monsoon climate. *Current Science* 84, 188-196.  
780  
781 Lee, K.S., Kim, J.M., Lee, D.R., Kim, Y., Lee, D. 2007. Analysis of water movement through an unsaturated soil  
782 zone in Jeju Island, Korea using stable oxygen and hydrogen isotopes. *Journal of Hydrology* 345, 199-211.

783 Majumdar, N., Majumdar, R.K., Mukherjee, A.L., Bhattacharya, S.K., Jani, R.A. 2005. Seasonal variations in  
784 the isotopes of oxygen and hydrogen in geothermal waters from Bakreswar and Tantloi, Eastern India:  
785 implications for groundwater characterization. *J. Asian Earth Sci.* 25, 269-278.

786  
787 Mathieu, R., Bariac, T. 1996. An isotopic study ( $^2\text{H}$  and  $^{18}\text{O}$ ) of water movements in clayey soils under a  
788 semiarid climate. *Water Resources Research* 32, 779-789.

789  
790 Maréchal J.C., Dewandel, B. Subrahmanyam. K. 2004. Contribution of hydraulic tests at different scales to  
791 characterize fracture network properties in the weathered-fissured layer of a hard rock aquifers. *Water Resources*  
792 *Res.*, 40, W11508.

793  
794 Maréchal, J.C., Dewandel, B., Ahmed, S., Galeazzi, L., Zaidi, F.K. 2006. Combined estimation of specific yield  
795 and natural recharge in a semi-arid groundwater basin with irrigated agriculture. *Journal of Hydrology* 329, 281-  
796 293.

797  
798 Mukherjee, A., Fryar, A.E., Rowe, H.D. 2007. Regional-scale stable isotopic signatures of recharge and deep  
799 groundwater in the arsenic affected areas of West Bengal, India. *Journal of Hydrology* 334, 151-161.

800  
801 Murad A.A., Krishnamurthy R.V. 2008. Factors controlling stable oxygen, hydrogen and carbon isotope ratios in  
802 regional groundwater of the United Arab Emirates (UAE). *Hydrological Processes* 22, 1922-1931.

803  
804 Nahon, D.B. 1991. Introduction to the petrology of soils and chemical weathering. J. Wiley & Sons Ed., 313p.

805  
806 Nair, A.R., Navada, S.V., Kulkarni, K.M. Kulkarni, U.P., U.P., Joseph, T.B. 1999. Environmental isotopes  
807 studies in the arid regions of western Rajasthan, India. *TecDoc IAEA 1207*, 41-57.

808  
809 Navada, S.V., Nair, A.R., Sinha, U.K., Kulkarni, U.P., Joseph, T.B. 1999. Application of isotopes and chemistry  
810 in unsaturated zone in arid areas of Rajasthan, India. *TecDoc IAEA 1246*, 119-130.

811  
812 Négrel, Ph., 1999. Geochemical study of a granitic area - The Margeride Mountains, France: chemical element  
813 behavior and  $^{87}\text{Sr}/^{86}\text{Sr}$  constraints. *Aquatic Geochemistry* 5, 125-165.

814  
815 Négrel, Ph., Lachassagne, P. 2000. Geochemistry of the Maroni River (French Guyana) during low water stage:  
816 Implications for water rock interaction and groundwater characteristics. *Journal of Hydrology* 237, 212-233.

817  
818 Négrel, Ph. 2006. Water-granite interaction: clues from strontium, neodymium and rare earth elements in  
819 saprolite, sediments, soils, surface and mineralized waters. *Applied Geochemistry* 21, 1432-1454.

820  
821 Négrel, Ph., Dewandel, B., Gandolfi, J.M., Dayal, A.M., Pauwels, H., Roy, S., Flehoc, C. 2007. Stable isotope  
822 hydrogeology of the Maheshwaram watershed (Andhra Pradesh, India). 3<sup>rd</sup> International Groundwater Conference  
823 (IGC-2007) on Water, Environment and Agriculture, Feb. 7-10, 2007, Combinatore, India. In *Groundwater*  
824 *Resources Assessment, Recharge and Management*, Macmillan Advanced Research Series, M.V. Ranghaswami,  
825 K. Palanisami, C. Mayilswami eds., 214-221.

826  
827 Négrel, Ph., Machard de Gramont, H., Lemièrre, B., Billaud, P., Sengupta, B. 2007. Hydrogeochemical  
828 processes, mixing and isotope tracing in hard rock aquifers and surface waters from the Subarnarekha river  
829 basin, (east Singhbhum district, Jharkhand state, India). *Hydrogeology Journal* 15, 1535-1552.

830  
831 Pauwels, H., Roy, S., Négrel, Ph., Ahmed, S., Dewandel, B., Gandolfi, J.M., Atal, S. 2007. Genesis and control  
832 of fluoride concentrations in groundwater of a small agricultural watershed (Maheshwaram, Andhra Pradesh,  
833 India). 3<sup>rd</sup> International Groundwater Conference (IGC-2007) on Water, Environment and Agriculture, Feb. 7-  
834 10, 2007, Combinatore, India. In *Groundwater Quality and Environment*, Macmillan Advanced Research Series,  
835 M.V Ranghaswami, K. Palanisami, C. Mayilswami eds., 158-164.

836  
837 Perrin, J., Dewandel, B., Aulong, S., Ahmed, S., Hrkal, Z., Krazny, J., Mascré, C., Massuel, S., Mukherji, A.,  
838 Samad, M. 2006. SUSTWATER Project Final Scientific Report. BRGM report RP-56913-FR, 137 p.

839  
840 Praamsma, T., Novakowski, K., Kyser, K., Hall, K. 2009. Using stable isotopes and hydraulic head data to  
841 investigate groundwater recharge and discharge in a fractured rock aquifer. *Journal of Hydrology* 366, 35-45.

842



843 Robins, N.S., Smedley, P.L. 1994. Hydrogeology and hydrogeochemistry of a small hard-rock island — the  
844 heavily stressed aquifer of Jersey. *Journal of Hydrology* 163, 249-269.  
845

846 Sami, K. 1992. Recharge mechanisms and geochemical processes in a semi-arid sedimentary basin, Eastern  
847 Cape, South Africa. *Journal of Hydrology* 139, 27-48.  
848

849 Simpson, H.J., Herczeg, A.L., Meyer, W.S. 1992. Stable isotope ratios in irrigation water can estimate rice crop  
850 evaporation. *Geophysical Research Letters* 19, 377-380.  
851

852 Singhal, D.C., Niwas, S., Singhal, B.B.S. 1988. Integrated approach to aquifer delineation in hard rock terrains-  
853 A case study from the Banda District, India. *Journal of Hydrology* 98, 165-183.  
854

855 Sukhija, B.S., Reddy, D.V., Nagabhushanam, P. 1998. Isotopic fingerprints of paleoclimates during the last  
856 30,000 years in deep confined groundwaters of Southern India. *Quaternary Research* 50, 252-260.  
857

858 Sukhija, B.S., Reddy, D.V., Nagabhushanam, P., Bhattacharya, S.K., Jani, R.A., Kumar, D. 2006.  
859 Characterisation of recharge processes and groundwater flow mechanisms in weathered-fractured granites of  
860 Hyderabad (India) using isotopes. *Hydrogeology Journal* 14, 663-674.  
861


862 Tardy, Y. 1971. Characterization of the principal weathering types by the geochemistry of waters from some  
863 European and African crystalline massifs. *Chemical Geology* 7, 253-271.  
864

865 Taylor, R., Howard, K. 2000. A tectono-geomorphic model of the hydrogeology of deeply weathered crystalline  
866 rock: evidence from Uganda. *Hydrogeology Journal*. 8, 279-294.  
867


868 Tirumalesh, K., Shivanna, K., Jalihal, A.A. 2006. Isotope hydrochemical approach to understand fluoride release  
869 into groundwaters of Ilkal area, Bagalkot District, Karnataka, India. *Hydrogeology Journal* 15, 589-598.  
870

871 Wyns, R., Gourry, J.C., Baltassat, J.M., Lebert, F. 1999. Caractérisation multiparamètres des horizons de  
872 subsurface (0-100 m) en contexte de socle altéré, in 2<sup>ème</sup> Colloque GEOFCAN, edited by I. BRGM, IRD,  
873 UPMC, pp. 105-110, Orléans, France.  
874


875 Zheng, Y., Datta, S., Stute, M., Dhar, R., Hoque, M.A., Rahman, M.W., Ahmed, K.M., Schlosser, P., van Geen,  
876 A. 2005. Stable isotopes (<sup>18</sup>O, <sup>2</sup>H) and arsenic distribution in the shallow aquifers in Araihasar, Bangladesh. EOS  
877 Transactions American Geophysical Union, 86 (52), Fall Meeting Supplement, Abstract H31B-1305.  
878  
879  
880  
881  
882  
883  
884 **Figure and table captions**  
885



886  
 887 Figure 1. Location of the Maheshwaram watershed (located 35 km south of Hyderabad) in Andhra Pradesh  
 888 (India), grey arrows indicate the main wind direction (and monsoon origin) for the periods June to September  
 889 and October to December (Deshpande et al., 2003). Simplified geological map of the Maheshwaram watershed  
 890 with location of sampled boreholes (Mxx). Black triangle: rain gauge in Maheshwaram village.  
 891




892  
 893 Figure 2.  $\delta^{18}\text{O}$ - $\delta^2\text{H}$  plot for groundwater from the Maheshwaram watershed in January (Fig. 2), March, June and  
 894 November 2006, June 2008 and February 2009. GMWL is the global meteoric water line ( $\delta^2\text{H} = 8 \delta^{18}\text{O} + 10$ ;  
 895 Craig, 1961). LMWL is the local meteoric water line defined in this study shown with its 95% uncertainty  
 896 envelopes.  $\delta^{18}\text{O}$ - $\delta^2\text{H}$  monsoon samples and weighted means are shown for 2006 and 2008 monsoons. The  
 897 monsoon data (Kumar et al., 1982) are from the Lower Maner Basin (Andhra Pradesh, north of Maheshwaram).  
 898 Groundwater in the watershed collected in January 2006 defines a co-variation (black line), the equation is  
 899  $\delta^2\text{H} = 6.17 \pm 0.25 \times \delta^{18}\text{O} - 0.54 \pm 0.67$  ( $n = 24$ ).






900


901 Figure 3.  $\delta^{18}\text{O}$  -  $\delta^2\text{H}$  plot for groundwater from the Maheshwaram watershed in January (Fig. 4a) and pan-  
 902 evaporation experiment (Fig. 4b). Tank-sample data are from the only surface water in the watershed. GMWL,  
 903 LMWL, and monsoon, mean-weighted-monsoon as on Figure 2.  $\delta^{18}\text{O}$  - Cl content plot of pan-evaporation  
 904 experiment; mean-weighted-monsoon and tank surface water.





905

906 Figure 4, 'D-excess' vs.  $\delta^{18}\text{O}$  plot for groundwater from the Maheshwaram watershed and for the monsoons (see  
 907 text).






909  
 910 Figure 5a, b. Seasonal  $\delta^{18}\text{O}$  vs.  $\delta^2\text{H}$  variations for selected groundwater collected in the Maheshwaram watershed  
 911 (moderate use of groundwater for cultivation through open-end wells M12, 14 and 57 and greater use of  
 912 groundwater through open-end wells M30, 35, 50 and 53; the monsoon, mean-weighted-monsoon, and the slope  
 913 of the evaporation line are shown as well. As on Figures 2 and 3, GMWL is the global meteoric water line and  
 914 LMWL is the local meteoric water line.




915  
 916 Figure 6. Cl<sup>-</sup> concentration vs. δ<sup>18</sup>O plot for groundwater from the Maheshwaram watershed, including data for  
 917 the monsoon, and from the study by Sukhija et al. (2006) in the catchment that comprises Hyderabad and its  
 918 suburbs. Dashed line corresponds to the one-step evaporation line following the evaporation experiment.





919

920 Figure 7. Results of the simulation for Cl<sup>-</sup> contents and δ<sup>18</sup>O signatures over a period of 20 years within the  
 921 Maheshwaram catchment, considering yearly evolution at 10, 15 and 20 years of overpumping of the aquifer.



922  
 923  
 924  
 925  
 926  
 927  
 928  
 929

Figure 8. Evolution along the 10th year of overpumping of the Maheshwaram aquifer, for aquifer volume, Cl content, and  $\delta^{18}\text{O}$ .\*\*

Period	$\delta^2\text{H} \text{‰}$	$\delta^{18}\text{O} \text{‰}$	Collected volume Rainfall gauge	Collected volume Meteo gauge	$\text{Cl}^-$
	V-SMOW ( $\pm 0.8\text{‰}$ )	V-SMOW ( $\pm 0.1\text{‰}$ )	L	L	mg. L <sup>-1</sup>
<b>Monsoon 2006</b>					
June 2006	-2.3	-1.2	0.70	0.60	4.75
July 2006	-11.4	-2.6	1.75	1.50	2.10
August 2006	0.2	-1.2	3.50	2.94	1.53
September 2006	-43.4	-6.4	4.00	4.18	1.30
October 2006	-44.4	-6.9	0.80	0.77	2.55
November 2006	-34.4	-5.5	0.25	0.40	3.30
Mean weighted monsoon 2006	-21.69	-3.83	-	-	1.86
<b>Monsoon 2008</b>					
June 2008	5.6	0.1	0.75	-	nd
July 2008	-6.4	-2.1	2.84	-	nd
August 2008	-12.6	-3	5.30	-	nd
September 2008	-49.7	-7.4	1.45	-	nd
October 2008	-30.2	-4.6	1.25	-	nd
November 2008	-34.1	-5.9	0.40	-	nd
Mean weighted monsoon 2008	-17.05	-3.39	-	-	-

930

931 Table 1. Results of stable isotopes of the water molecule ( $\delta^{18}\text{O}$  and  $\delta^2\text{H}$ ), collected volumes, and  $\text{Cl}^-$  contents for  
932 monsoon precipitation collected in Maheshwaram village in 2006.

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

Sample	T °C	EC μS cm <sup>-1</sup>	Cl <sup>-</sup> mg. L <sup>-1</sup>	δ <sup>2</sup> H ‰ V-SMOW (±0.8‰)	δ <sup>18</sup> O ‰ V-SMOW (±0.1‰)	D-excess
<b>January 2006</b>						
A11	27.5	850	34.41	-14.3	-2.3	4.1
M02	28.4	1440	166.22	-13.3	-2.0	2.7
M03	27.2	770	40.64	-11.5	-2.1	5.3
M06	29.8	685	36.98	-19.1	-2.7	2.5
M08	26.6	700	37.58	-9.6	-1.6	3.2
M12	26.0	750	17.34	-18.2	-2.8	4.2
M14	27.6	900	29.75	-22.9	-3.8	7.5
M15	27.9	3310	263.21	-17.0	-2.9	6.2
M16	27.3	1800	238.00	-23.4	-3.8	7.0
M19N	27.4	1470	167.04	-14.1	-2.1	2.7
M20	27.5	3570	429.15	-17.4	-2.7	4.2
M25	28.1	700	31.45	-16.6	-2.5	3.4
M28	26.6	670	15.65	-13.8	-2.0	2.2
M29	25.9	1390	126.84	-16.5	-2.5	3.5
M30	26.3	670	11.70	-26.9	-4.1	5.9
M33	27.7	600	11.78	-23.3	-3.8	7.1
M35N	27.1	1060	77.91	-13.0	-1.8	1.4
M36N	28.8	750	31.86	-14.8	-2.4	4.4
M44	26.7	940	55.77	-3.7	-0.6	1.1
M46	29.3	800	68.91	-16.7	-2.6	4.1
M50	26.9	770	36.82	-17.2	-2.5	2.8
M51	27.3	1830	232.62	-8.6	-1.3	1.8
M53	28.4	810	11.35	-17.3	-2.8	5.1
M57	28.6	1100	51.66	-22.4	-3.6	6.4
<b>March 2006</b>						
A11	27.4	820	29.62	-15.5	-2.3	2.9
M02	28.5	970	97.91	-33.1	-4.7	4.5
M03	26.8	800	41.12	-11.0	-2.0	4.6
M06	28.6	800	45.50	-10.5	-1.5	1.5
M08	27.0	660	38.97	-6.8	-1.1	2.0
M12	26.5	797	24.30	-18.7	-2.9	4.5
M14	27.6	940	42.89	-22.0	-3.7	7.6
M15	28.8	3270	290.07	-17.5	-3.1	7.3
M16	nd	1780	235.28	-24.6	-3.8	5.8
M19N	27.1	180	243.59	-13.1	-1.9	2.1
M20	29.1	1077	57.17	-26.6	-3.9	4.6
M25	29.1	590	17.43	-17.8	-2.7	3.7
M28	27.8	690	13.84	-13.6	-2.2	4.0
M29	26.8	930	66.32	-15.9	-2.2	1.7
M30	27.5	1460	10.81	-24.7	-3.6	4.1
M33	28.2	610	12.18	-24.5	-3.9	6.7
M35N	27.4	1140	96.80	-11.7	-1.5	0.3
M36N	28.6	760	31.96	-15.8	-2.8	6.6
M37	28.6	1490	237.33	-4.8	-0.6	0.0
M44	26.0	940	60.41	-4.6	-0.5	-0.6
M46	28.6	790	65.34	-14.7	-2.3	3.7
M48	27.5	650	24.46	-31.2	-4.6	5.6
M50	27.4	650	40.37	-14.7	-2.1	2.1
M51	27.5	1770	184.74	-9.7	-1.4	1.5
M53	28.0	790	10.72	-17.0	-2.7	4.6
M57	30.1	1080	44.79	-22.1	-3.7	7.5
Tank	nd	nd	nd	-10.3	-0.3	-7.9
<b>June 2006</b>						

A11	28.8	790	31.73	-13.9	-2.3	4.5
M2	nd	nd	173.78	-16.3	-2.4	2.9
M3	28.2	840	48.85	-10.8	-2.0	5.2
M6	27.2	690	38.51	-16.6	-2.3	1.8
M8	nd	nd	45.31	-9.1	-1.4	2.1
M12	nd	nd	24.62	-19.3	-3.1	5.5
M14	28.6	890	34.30	-23.7	-3.9	7.5
M15	28.9	1480	208.46	-22.2	-3.6	6.6
M16	27.8	1740	238.38	-23.2	-3.8	7.2
M25	28.3	650	25.28	-15.8	-2.5	4.2
M29	nd	nd	52.53	-20.9	-2.9	2.3
M30	27.5	720	12.56	-24.3	-3.6	4.5
M33	28.2	640	15.82	-22.9	-3.6	5.9
M35N	nd	nd	60.69	-6.6	0.3	-9.0
M36N	29.0	780	32.59	-15.9	-2.5	4.1
M37	28.7	1160	199.16	-7.8	-0.9	-0.6
M44	nd	nd	70.63	-11.7	-1.8	2.7
M46	28.7	756	57.27	-17.7	-2.7	3.9
M48	nd	nd	6.14	-26.0	-3.9	5.2
M50	nd	nd	43.66	-13.7	-2.0	2.3
M51	27.4	1610	197.50	-12.0	-1.8	2.4
M53	28.9	810	13.90	-17.2	-2.8	5.2
M57	28.8	1060	49.27	-23.1	-3.7	6.5
<hr/> <b>November 2006</b>						
M2	28.3	1940	170.28	-17.0	-2.5	3.0
M8	nd	Nd	45.74	-8.8	-1.4	2.7
M12	nd	Nd	13.72	-22.3	-3.7	7.5
M14	nd	Nd	34.95	-23.4	-3.9	7.8
M15	28.2	4930	369.61	-16.7	-3.0	7.3
M29	27.5	1350	66.52	-21.7	-3.3	4.7
M30	nd	Nd	9.48	-20.0	-3.0	3.8
M35	nd	1280	56.70	-8.1	-0.3	-5.7
M50	27.4	1300	46.58	-11.9	-1.7	1.7
M53	28.9	1120	12.29	-15.6	-2.4	3.6
M57	nd	nd	49.47	-22.8	-3.7	6.9
<hr/> <b>June 2008</b>						
M6	27.9	nd	34.10	-11.0	-1.7	2.6
M8	28.7	780	51.15	-11.0	-1.7	2.6
M12	29.2	700	25.73	-21.7	-3.6	6.7
M30	28.3	700	9.66	-21.3	-3.2	4.6
M35	27.7	1200	86.44	-5.2	-0.1	-4.5
M48	28.4	nd	50.41	-18.7	-2.9	4.6
M50	27.7	920	49.10	-13.2	-1.9	2.3
M53	30.1	870	16.55	-14.2	-2.2	3.7
M57	30.6	1510	194.06	-22.0	-3.7	7.2
<hr/> <b>February 2009</b>						
M2	28.8	1303	154.71	-15.2	-2.4	3.8
M6	27.6	752	31.73	-16.3	-2.2	1.4
M8	23	596	25.88	-18.6	-3.1	6.5
M12	28.6	763	37.30	-21.3	-3.5	6.5
M30	27.7	683	8.40	-19.1	-2.7	2.7
M35	27.5	1085	61.65	-3.0	0.0	-3.3
M50	26.9	982	46.83	-12.2	-1.7	1.3
M53	29.6	857	14.00	-15.4	-2.4	3.8
M57	32	2000	359.01	-19.3	-3.1	5.4
<hr/> <b>Tank</b>						

Tank March 2006	nd	nd	13.6	-10.3	-0.3	-7.9
Tank November 2006	nd	nd	18.7	-5.4	-0.4	-2.2
Tank February 2009	nd	nd	50.42	38.4	8.4	-28.2

nd = non-determined values

959

960 Table 2. Results of field variables (EC in  $\mu\text{S}/\text{cm}$ ; temperature in  $^{\circ}\text{C}$ ; pH) and stable isotopes of the water  
 961 molecule ( $\delta^{18}\text{O}$  and  $\delta^2\text{H}$ ), in ‰ with reference to V-SMOW (Vienna Standard Mean Ocean Water) for  
 962 groundwater collected in the Maheshwaram watershed.

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

<b>Period</b>	<b>Equation</b>	<b>n</b>	<b>R<sup>2</sup></b>
January 2006	$\delta^2\text{H} = 6.17 \pm 0.25 \times \square \delta^{18}\text{O} - 0.54 \pm 0.67$	24	> 0.98
March 2006	$\delta^2\text{H} = 6.44 \pm 0.18 \times \delta^{18}\text{O} - 0.24 \pm 0.50$	26	> 0.98
June 2006	$\delta^2\text{H} = 5.30 \pm 0.19 \times \square \delta^{18}\text{O} - 3.33 \pm 0.54$	23	> 0.98
November 2006	$\delta^2\text{H} = 4.80 \pm 0.43 \times \square \delta^{18}\text{O} - 4.49 \pm 1.22$	11	> 0.98
June 2008	$\delta^2\text{H} = 5.11 \pm 0.30 \times \square \delta^{18}\text{O} - 3.44 \pm 0.77$	9	> 0.98
February 2009	$\delta^2\text{H} = 5.16 \pm 0.32 \times \square \delta^{18}\text{O} - 3.51 \pm 0.81$	9	> 0.98

996

997 Table 3. Results of  $\delta^{18}\text{O}$  and  $\delta^2\text{H}$  relationships for the January, March, June and November 2006, June 2008,  
998 February 2009 collection periods, n is the number of open end wells collected and R<sup>2</sup> is the correlation  
999 coefficient.

1000

1001

1002

1003