

Behaviour of lithium and its isotopes during weathering in the Mackenzie Basin, Canada

Romain Millot, Nathalie Vigier, Jérôme Gaillardet

► To cite this version:

Romain Millot, Nathalie Vigier, Jérôme Gaillardet. Behaviour of lithium and its isotopes during weathering in the Mackenzie Basin, Canada. Geochimica et Cosmochimica Acta, 2010, 74 (14), p. 3897-3912. 10.1016/j.gca.2010.04.025 . hal-00553199

HAL Id: hal-00553199 https://brgm.hal.science/hal-00553199

Submitted on 6 Jan 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

3

8 9

10

11 12 13

Behaviour of lithium and its isotopes during weathering in the Mackenzie Basin, Canada 2

Romain Millot (1)^{*}, Nathalie Vigier (2) and Jérôme Gaillardet (3) (1) BRGM, Metrology Monitoring Analysis Department, Orléans, France Corresponding author, e-mail: r.millot@brgm.fr (2) CRPG, CNRS, Nancy-Université, Vandœuvre-lès-Nancy, France (3) IPGP, Equipe de Géochimie et Cosmochimie, Université Paris 7, Paris, France

14 Abstract

15 We report Li isotopic compositions, for river waters and suspended sediments, of about 40 rivers sampled within 16 the Mackenzie River Basin in northwestern Canada. The aim of this study is to characterize the behaviour of Li 17 and its isotopes during weathering at the scale of a large mixed lithology basin. The Mackenzie river waters 18 display systematically heavier Li isotopic compositions relative to source rocks and suspended sediments. The 19 range in δ^7 Li is larger in dissolved load (from +9.3 to +29.0%) compared to suspended sediments (from -1.7 to 20 +3.2‰), which are not significantly different from δ^7 Li values in bedrocks. Our study shows that dissolved Li is 21 essentially derived from the weathering of silicates and that its isotopic composition in the dissolved load is 22 inversely correlated with its relative mobility when compared to Na. The highest enrichment of ⁷Li in the dissolved 23 load is reported when Li is not or poorly incorporated in secondary phases after its release into solution by mineral 24 dissolution. This counterintuitive observation is interpreted by the mixing of water types derived from two different 25 weathering regimes producing different Li isotopic compositions within the Mackenzie River Basin. The incipient 26 weathering regime characterizing the Rocky Mountains and the Shield areas produces ⁷Li enrichment in the fluid 27 phase that is most simply explained by the precipitation of oxy-hydroxide phases fractionating Li isotopes. The 28 second weathering regime is found in the lowland area and produces the lower δ^7 Li waters (but still enriched in ⁷Li 29 compared to bedrocks) and the most Li-depleted waters (compared to Na). Fractionation factors suggest that the 30 incorporation of Li in clay minerals is the mechanism that explains the isotopic composition of the lowland rivers. 31 The correlation of boron and lithium concentrations found in the dissolved load of the Mackenzie rivers suggests 32 that precipitation of clay minerals is favoured by the relatively high residence time of water in groundwater. In the 33 Shield and Rocky mountains, Li isotopes suggest that clay minerals are not forming and that secondary minerals 34 with stronger affinity for ⁷Li appear. 35 Although the weathering mechanisms operating in the Mackenzie Basin need to be characterized more precisely, 36 the Li isotope data reported here clearly show the control of Li isotopes by the weathering intensity. The spatial 37 diversity of weathering regimes, resulting from a complex combination of factors such as topography, geology, 38 climate and hydrology explains, in fine, the spatial distribution of Li isotopic ratios in the large drainage basin of 39 the Mackenzie River. There is no simple relationship between Li isotopic composition and chemical denudation 40 fluxes in the Mackenzie River Basin.

42	Keywords: lithium isotopes, river waters, silicate weathering, sediments, erosion, Mackenzie Basin
43	Submitted on September 22 nd 2009 – Revised on April 22 nd 2010
44	7 198 words (without the references and captions)

1. INTRODUCTION

46

Assessing the behaviour of lithium and the distribution of Li isotopes during weathering is of major importance for studying water/rock interactions at the surface of the Earth. This is because lithium (⁶Li ~7.5% and ⁷Li ~92.5%) is a fluid-mobile element and, due to the large relative mass difference between its two stable isotopes, it is subject to significant low temperature mass fractionation which provides key information on the nature of weathering processes.

Recent studies have shown that the range of δ^7 Li values spans more than 40‰ at the Earth 53 surface ($\delta^7 \text{Li} = ({}^7 \text{Li}/{}^6 \text{Li}_{\text{sample}} / {}^7 \text{Li}/{}^6 \text{Li}_{\text{L-SVEC}}$ -1) x10³), (e.g. Pistiner and Henderson 2003, 54 55 Rudnick et al. 2004, Kisakürek et al. 2004, Huh et al. 2004, Kisakürek et al. 2005, Hathorne and James 2006, Pogge von Strandmann et al. 2006, Vigier et al. 2009). Furthermore, the 56 world-wide range of δ^7 Li in river waters is between +6 and +33‰ (Huh et al. 1998, 2001). 57 There are as yet few studies concerning mixed lithology basins, but these suggest that river 58 water δ^7 Li is not affected by differences in catchment lithology (Kisakürek et al. 2005, Huh et 59 al. 1998). Rather, lithium isotopes are thought to be strongly affected by isotope fractionation 60 during secondary mineral formation and the degree and type of silicate weathering 61 (Kisakürek et al. 2004, Huh et al. 2004, Kisakürek et al. 2005, Hathorne and James 2006, 62 Pogge von Strandmann et al. 2006, Vigier et al. 2009). 63

To date, both the magnitude of the Li isotopic fractionation associated with water-rock interaction processes, and the factors controlling these fractionations, are poorly understood. However, both field and experimental studies have shown that ⁶Li is preferentially retained by secondary minerals during silicate weathering (Pistiner and Henderson 2003, Kisakürek et al. 2004, Pogge von Strandmann et al. 2006, Vigier et al. 2009). Accordingly, the fractionation of Li isotopes is dependent upon the extent of chemical weathering. Large fractionation seems to occur during superficial weathering while little fractionation is observed during more

intense or prolonged weathering in stable environments (Huh et al. 1998, 2001, Pogge von
Strandmann et al. 2006).

73 Lithium isotopic fractionation has been documented in numerous natural environments, with 74 experimental and natural data (Huh et al. 2001, Pistiner and Henderson 2003, Tomascak 2004). It has been shown that partial dissolution of basalts does not result in fractionation of 75 lithium isotopes, but that dissolution of granitic rock can cause fractionation (Pistiner and 76 77 Henderson 2003). In addition, adsorption onto mineral surfaces can also be a mechanism of 78 Li isotopic fractionation in the hydrosphere. Sorption of Li from aqueous solutions by mineral phases at the temperature of the Earth's surface has been highlighted by Taylor and Urey 79 (1938), as well as by Anderson et al. (1989). Li sorption experiments utilizing several 80 minerals (Pistiner and Henderson 2003) have revealed that Li isotopic fractionation does not 81 82 occur when this element is not incorporated into the structure of the solid (physical sorption). When Li is incorporated by stronger bonds (chemical sorption), an isotopic fractionation is 83 observed that is dependent on the chemical structure of the minerals (Anghel et al. 2002). 84 The most significant Li isotopic fractionation factor (A solution - solid) ~ +14‰ has been measured 85 86 for gibbsite. For Li sorption processes between solution and different solids (clay minerals 87 such as kaolinite and vermiculite, and freshwater sediments), Zhang et al. (1998) have observed even higher Li isotopic fractionation factors (Δ solution - solid ~ +22‰). 88

89 In the present paper, we have undertaken a systematic study of the weathering products (both dissolved load and suspended sediments) of the Mackenzie River Basin, one of the 90 major high latitude river basins of North America. This area is of particular interest because it 91 92 has previously been extensively studied and is well characterized from a lithology and 93 weathering mass budget point of view (Vigier et al. 2001, Millot et al. 2002, Millot et al. 2003, 94 Gaillardet et al. 2003). Also, there is a clear contrast between mountains and plains, with a fourfold increase of silicate chemical weathering rates in the plains (Millot et al. 2003). A 95 direct comparison of silicate weathering rates and river Li isotope compositions can be 96 97 performed. In this paper, we evaluate the potential of Li isotopes as a tracer of silicate 98 weathering processes in large mixed lithology basins at the scale of a large river basin.

100

101

2. GEOLOGICAL SETTING AND SAMPLE LOCATIONS

This study focuses on the Mackenzie River basin and adjacent river basins (Stikine, Nass, 102 Skeena, Fraser, and Yukon rivers). Most of the river waters and sediments are from the 103 Mackenzie River Basin itself (Figure 1), which is located in northwestern Canada under arctic 104 105 and subarctic climate. The mainstream of the Mackenzie River was sampled at two locations 106 (samples CAN96-6 at the river mouth and CAN96-25 upstream). The main tributaries of the Mackenzie River are the Red Arctic River (CAN96-7), the Liard River (CAN96-26), the Peel 107 River (CAN96-5), the Peace River (CAN96-15 and CAN96-37), the Slave River (CAN96-38) 108 109 and the Athabasca River (CAN96-42).

110 The Mackenzie River basin is composed of three main structural units, corresponding to 111 different geomorphological zones: namely, the Rocky and the Mackenzie Mountains in the western part, the Interior Platform (lowlands), and the Canadian Shield (in the eastern part), 112 113 (Figure 1). Rivers belonging to a transition-zone correspond to basins draining the foothills located between the Rockies and the low-lying plains. From a mass budget point of view, the 114 mainstream of the Mackenzie is mostly fed by tributaries draining the Interior Plain. River 115 samples from the adjacent basins of the Yukon and Fraser Rivers are also considered here 116 117 because their geological contexts are similar to those of the Upper Liard and Athabasca sub-118 basins, both located in the Rocky Mountains. Finally, the main rivers draining the Western Canadian orogenic belt were also analysed (Nass, Skeena and Stikine River). 119

Sampled rivers therefore drain a large range of lithologies: volcanics and immature volcanoclastic sediments in the Western Canadian orogenic belt (Stikine terrane), carbonates and slates in the Mackenzie Mountains, marine and non-marine sedimentary rocks (Cambrian to Cretaceous limestones, shales and sandstones) in the Interior plains of the basin, and old silicate rocks (Archean granites and gneisses) in the Slave Province of the Canadian Shield. The chemical weathering conditions and physical erosion rates differ greatly depending on

the runoff and basin lithology. Chemical denudation rates in this arctic zone are rather low

127 compared to other major river basins, as determined by Millot et al. (2003) and Gaillardet et
128 al. (2003). Full details concerning river samples (geological and hydrological contexts) have
129 been published in Millot et al. (2003) for the Mackenzie River basin, in Millot et al. (2002) for
130 the rivers of the Canadian Shield and in Gaillardet et al. (2003) for rivers of the Western
131 Cordillera.

- 132
- 133

3. ANALYTICAL METHODS

134

3.1. Sampling methodology, major and trace element concentration measurements 136

137 River waters were sampled twice during the high flow stage of the summer season (August 1996 and June 1999). In the field, 10-15 L of river water was collected in acid-washed 138 containers for major and trace elements, and isotopic measurements (Millot et al. 2003). 139 140 Samples were filtered a few hours later, using a Sartorius frontal filtration unit (0.2 µm cellulose acetate filter, 142 mm diameter). After filtration, samples were stored in acid-141 washed polypropylene bottles. The river sediments were collected after drying and 142 centrifugation of the retentate of river water filtration. The samples for cations and trace 143 144 element concentrations and Li isotope analyses were acidified to pH = 2 with ultrapure 145 HNO₃. Bottles for anion analyses were not acidified.

Lithium concentrations (for river waters and suspended sediment samples) were determined by ICP-MS (VG PlasmaQuad II+) with indium as an internal standard, with a precision of \pm 5%.

Major and trace elements are reported in Millot et al. (2003) for the Mackenzie River basin, in Millot et al. (2002) for the rivers of the Canadian Shield, and in Gaillardet et al. (2003) for rivers of the Western Cordillera respectively.

152

153 **3.2. Lithium isotope measurements**

154

Lithium isotopic compositions were measured using a Neptune Multi-Collector ICP-MS (Millot et al. 2004). ⁷Li/⁶Li ratios were normalized to the L-SVEC standard solution (NIST SRM 8545, Flesch et al. 1973) following the standard-sample bracketing method. Typical in-run precision on the determination of δ^7 Li is about 0.1-0.2‰ ($2\sigma_m$, standard error of the mean).

159 Chemical separation of Li from the matrix was achieved before isotope analyses, following a 160 procedure modified from the technique of James and Palmer (2000) that uses a cationic 161 exchange resin (a single column filled with 3 mL of BioRad AG® 50W-X12 resin, 200-400 162 mesh) and HCI acid media (0.2N) for 30 ng of Li. Blanks for the total chemical extraction 163 were less than 30 pg of Li, which is negligible since this represents a blank/sample ratio of 164 $<10^{-3}$.

Successful quantitative measurement of Li isotopic compositions requires 100% Li recovery during laboratory processing. Therefore, frequent column calibrations were performed and repeated analysis of L-SVEC standard processed through the columns shows that no isotope fractionation occurred as a result of the purification process.

Accuracy and reproducibility of the isotopic measurements were tested by repeated analyses of three Li standard solutions (namely ⁶Li-N, LiCl-N and ⁷Li-N, Carignan et al. 2007). Mean values of δ^7 Li = -8.0‰ ± 0.3 (2 σ , n = 38), δ^7 Li = +10.1‰ ± 0.2 (2 σ , n = 46) and δ^7 Li = +30.2‰ ± 0.3 (2 σ , n = 89), were obtained for ⁶Li-N, LiCl-N and ⁷Li-N respectively over a period of 10 months. Thus, long-term reproducibility of the Li mass analysis is better than 0.3‰ at the two sigma level for standard solutions (σ : standard deviation).

The accuracy and reproducibility of the entire method (purification procedure + mass analysis) were tested by repeated measurement of a seawater standard solution (IRMM BCR-403) after separation of Li from the matrix, for which we obtained a mean value of $\delta^7 \text{Li} =$ +30.8‰ ± 0.4 (2 σ , n = 15) over the period of the duration of the analyzes. This mean value is in a good agreement with our long-term measurements ($\delta^7 \text{Li} =$ +31.0‰ ± 0.5, 2 σ , n = 30, Millot et al. 2004) and with other values reported in the literature (see for example Carignan et al. 2004 and Tomascak 2004 for a compilation).

For suspended sediments, as well as for sands and rocks, a total digestion of 50 mg of 182 crushed sample was performed over 4 days at 100°C in a closed beaker with a mixture of 183 184 three ultrapure acids: 4 mL of HF (23N), 1 mL of HNO₃ (14N) and 0.1 mL of HClO₄ (12N). The solution was subsequently evaporated to dryness and 4 mL of HCl acid (6N) was added 185 and left for a further 4 days at 100°C. Sample aliquots (30 ng of Li) of the residue of the acid 186 dissolution were then dissolved in 0.5 mL of HCI (0.2N), before being placed on cation 187 exchange columns for Li separation. Accuracy and reproducibility of the procedure for solid 188 samples (dissolution + purification procedure + mass analysis) were tested by repeated 189 measurement of the JB-2 basalt standard (Geological Survey of Japan) which gave a mean 190 value of $\delta^7 Li = +4.9\% \pm 0.6$ (2 σ , n = 17), in good agreement with published values (see 191 Jeffcoate et al. 2004, Tomascak 2004 and Carignan et al. 2007 for data compilation). 192

- 193
- 194

4. RESULTS

195

196 4.1. Dissolved phase

197

Lithium concentrations (Table 1) in the Mackenzie basin river waters display a wide range of 198 values, from 0.05 µmol/L (for the Skeena River in the W. Cordillera) to 1.29 µmol/L (for the 199 Hay River located in the lowlands). The Mackenzie River at the river mouth (CAN96-6) has a 200 201 Li concentration of 0.57 µmol/L. The main tributaries, which are mainly located in the lowlying plains, show a mean value of Li of 0.63 μ mol/L, (n = 11). The group of rivers draining 202 the Interior Platform display the highest Li contents (mean value of 0.85 µmol/L, n = 9). In 203 contrast, rivers of the Shield and the W. Cordillera areas display the lowest Li levels (~0.16 204 μ mol/L, n = 4 and ~0.08 μ mol/L, n = 3, respectively). Also, rivers draining the Rocky 205 Mountains have low Li contents (mean value of 0.30 µmol/L, n = 11). Finally, Li contents for 206 rivers located in the transition zone are intermediate between those from the Rockies and 207 208 those from the low-lying plains (mean Li = $0.54 \mu mol/L$, n = 4).

Compared to the concentrations of dissolved Li reported in the pioneering work of Reeder et 209 al. (1972) for the Mackenzie river system (Li ranging from <0.7 to 4.5 µmol/L), our values are 210 211 clearly lower in the present case (from 0.05 to 1.29 µmol/L). This difference is likely to be 212 related to the method used for the Li determination in waters. Indeed, Reeder et al. (1972) reported a detection limit of 0.70 µmol/L (~5 µg/L) by atomic adsorption in the early 1970's, 213 whereas it is now ~0.01 µmol/L using modern ICP-MS techniques. Consequently, it is 214 obvious that Reeder et al. (1972) were not able to measure low lithium concentrations in 215 216 some of the rivers of the Mackenzie Basin (indeed Li was not detected in 49 samples of their 217 suite of 101 river samples).

Lithium isotopic compositions in the Mackenzie river waters (Table 1) range from $\delta^7 Li = +9.3$ 218 (for the Smoky River located in the transition zone) to +29.0‰ (for the Yellowknife River 219 located in the Slave Province). Rivers from the Lowland regions vary from +10.1‰ to 220 +16.8‰ with an average of +13.2‰. Rivers from the Rockies display higher δ^7 Li values (from 221 +12.2‰ to +20.5‰, with an average of +15.7‰), when compared to rivers draining the 222 plains. Rivers draining the W. Cordillera and the Canadian Shield show the heaviest isotope 223 compositions, with δ^7 Li ranging between +25.2 to +29.0% for the Shield rivers and between 224 +21.6 and +24.9% for the western orogenic belt (Stikine terrane). 225

An inverse relationship of Li isotopic composition with Li concentrations is observed for the Mackenzie river system (Figure 2). It is clear from this figure that the different physiographic provinces in the basin are characterized by different domains. The large tributaries display δ^7 Li values that essentially reflect the mixture of waters from the mountains and waters from the lowlands, in the range +12.7 to +16.5‰ (with an average of +14.6‰). The Mackenzie River at river mouth (CAN96-6) shows a δ^7 Li value of +15.0‰, close to both values reported by Huh et al. (1998) for the Mackenzie river mainstream (+17.9 and +15.7‰, respectively).

233

4.2. River sediments and bedrocks

Lithium concentrations in suspended sediments of the Mackenzie basin rivers (Table 2) range between 17.8 μ g/g (for the Mackenzie River at river mouth) and 57.8 μ g/g (for the Trout River in the Rockies). Lithium contents of the Mackenzie suspended sediments show almost the same range as that determined for the suspended sediments of the Orinoco basin (from 5.2 to 71.3 μ g/g, see Huh et al. 2001) and are close to the mean value of 35 ± 11 μ g/g reported for the Upper Continental Crust (UCC) by Teng et al. (2004).

The isotopic composition of the suspended sediments range from -1.7% to +3.2%, with a 242 243 mean value of +0.4‰ (n = 20, Table 2), identical, within uncertainties, to the estimate of the UCC (δ^7 Li from -2 to +2 ‰, Teng et al. 2004). Suspended sediments from the main 244 tributaries of the Mackenzie Basin display slightly negative δ^7 Li values (from -1.6 to -0.2‰) 245 whereas δ^7 Li of suspended sediments of rivers located in the Rocky Mountains are slightly 246 positive (from +0.3 to +3.2‰). The suspended sediments of the Nass and Stikine Rivers (W. 247 Cordillera) exhibit positive and similar isotope signatures ($\delta^7 Li = +1.0$ and +1.2%248 respectively). As for Li content, the range of δ^7 Li for suspended sediments of the Mackenzie 249 Basin is comparable with the values obtained by Huh et al. (2001) for silicate regions of the 250 Orinoco Basin. δ^7 Li values measured in suspended sediments are also in good agreement 251 with the range reported by Kisakürek et al. (2005) for the Himalayan rivers (-3.9 to +3.0%). 252

The sand sample collected at the mouth of the Mackenzie River (CAN96-6) has a Li concentration of 47.4 μ g/g, which is lower than the Li content of the corresponding suspended material (57.8 μ g/g) and a Li isotopic composition of -0.5‰, not significantly different from that of the corresponding suspended sediments (-0.9‰).

Bedrocks (carbonates and black shales) sampled in the basin, and assumed to be representative of the main type of rocks exposures, display Li contents ranging between 1.5 and 64.4 μ g/g (Table 3). The carbonate samples from the Rocky Mountains (CAN99-65) display the lowest concentration (1.5 μ g/g), in agreement with literature data reported for other carbonates (e.g. Huh et al. 2001, Hathorne and James 2006). In contrast, Li contents in blackshales from the Interior Platform are high (29.1 and 64.4 μ g/g). Two glacial tills sampled

within the Slave Province (Yellowknife and Prosperous Lake) display different Li
 concentrations of 16.2 and 54.6 μg/g respectively.

The carbonate sample (CAN99-65, Rocky Mountains) has the highest δ^7 Li value (+7.2%), in 265 agreement with other values reported for continental carbonates by Huh et al. (2001). This 266 value is likely to reflect a mixing between pure marine carbonate ($\delta^7 Li > +25\%$). Tomascak 267 2004, Hall et al. 2005, Hathorne and James 2006, Vigier et al. 2007) and detrital phases with 268 much lower δ^7 Li values and higher Li contents. δ^7 Li for the blackshales are negative (-1.0 and 269 270 -1.1‰ for the Interior plains and for the Transition zone samples respectively), but remain close to the estimate of the UCC. Finally, the two glacial till samples display relatively similar 271 δ^7 Li values of +4.8 and +5.2‰ respectively. 272

- 273
- 274

5. DISCUSSION

- 275
- 276 **5.1. Sources of dissolved lithium**
- 277

The sources of dissolved lithium in the river waters of the Mackenzie Basin can be assumed 278 279 to be either released by rock weathering (including solubilisation of the bedrock, and destabilisation or desorption from secondary minerals), derived from the atmosphere (via 280 precipitation or wet deposition), or derived from groundwater input. First of all, atmospheric 281 input can be evaluated using chloride concentrations if no evaporite is present within the 282 283 drainage basin. This is the case for most of the rivers of the Mackenzie Basin, as shown by Millot et al. (2002), Millot et al. (2003) and Gaillardet et al. (2003). In particular, rivers draining 284 the Shield area, characterized by low chemical weathering rates, are good candidates for 285 estimating the atmospheric input of Li. Rivers draining the Canadian Shield have low 286 287 dissolved Li contents (from 0.07 to 0.2 μ mol/L) and display the highest δ^7 Li values, close to that of the ocean (δ^7 Li = +31.0‰), a priori suggesting a marine origin for Li via seasalt input 288 289 through precipitation. Then, if we assume that all chloride in the Shield rivers (with Li/Cl =

5x10⁻³ and δ^7 Li = +25.2‰ on average) is of atmospheric origin and that marine salts have seawater Li/Cl molar ratios of 5x10⁻⁵ and δ^7 Li = +31.0‰, it can be calculated that only a small fraction of riverine Li (< 1%) in these rivers originates from seasalt input through precipitation. The same calculation for rivers draining the Western Cordillera (i.e. the rainy region: 1500-1750 mm/year compared to 350 mm/yr in the Shield area) indicates that a maximum of only ~0.4% riverine Li could have been supplied by precipitation.

We provided evidence in a previous paper (Millot et al. 2002), that the rivers of the Shield 296 297 area (Slave Province) are characterized by a chloride excess compared to a pure marine input, and we attributed their relatively high CI concentrations to the dissolution of evaporite-298 type aerosols derived from the dry area located south of the Great Slave Lake. Cambrian 299 300 evaporites occur at the base of the sedimentary basin of the lowlands, and these have 301 contributed to redeposited salt in modern lakes. The Salt River is one of the rivers that is highly influenced by the dissolution of these formations. Although this river was not analysed 302 in the present work, Li and Na concentrations of the Salt River (Li = 4.5 µmol/L and Na = 303 304 276.45 mmol/L) have been reported by Reeder et al. (1972). The typical Li/Na ratio of this river is 1.6x10⁻⁵ mol/mol. Considering this ratio along with the highest CI concentration found 305 306 in rivers draining the Slave Province (80 µmol/L), we estimate there is a negligible Li input from evaporites in the Shield rivers (less than 0.4% of lithium derived from evaporite). 307

308 In summary, our calculations indicate that the contribution of Li coming from precipitation and wet deposition is negligible, and that dissolved Li in all sampled rivers is mainly derived from 309 weathering of silicate and carbonate rocks. Although carbonates have relatively low Li 310 concentrations (< 1 μ g/g) compared to silicates (higher than 50 μ g/g), their higher dissolution 311 312 rates may make them a significant source of Li in carbonate rich-regions. Millot et al. (2002) 313 showed that the areas of the Mackenzie Basin which are most influenced by carbonate 314 dissolution are the Rocky and Mackenzie Mountains. In contrast, the lowlands are less 315 influenced by this lithology and the Shield area displays no input from carbonate weathering. 316 The proportions of lithium derived from silicate and carbonate weathering can be calculated

317

11

from the results of the inversion model (Millot et al. 2003) which allowed us to determine, for

each major element present in the dissolved phase, the proportion derived from the weathering of silicate and carbonate end-members within the Mackenzie River Basin. We use a mean Li/Ca molar ratio of 15x10⁻⁶ for carbonates, in agreement with Li/Ca molar ratios ranging from 10x10⁻⁶ to about 20x10⁻⁶ reported in carbonates elsewhere (Hall and Chan 2004, Lear and Rosenthal 2006, Hathorne et al. 2009), and determine that the carbonate end-member contribution is very low for lithium. Between 90.5% and 99.4% of dissolved lithium is thus derived from silicate weathering in the Mackenzie River Basin (Figure 3).

325 These estimates agree well with previous studies that have inferred that most riverine Li derives from silicate lithologies (Huh et al. 1998, Kisakürek et al. 2005). In addition, the 326 absence of a correlation between Li and Sr isotopic ratios (from Millot et al. 2003) argues 327 against a carbonate weathering control on riverine Li isotopic composition. We therefore 328 conclude that Li concentrations in the dissolved load of rivers in the Mackenzie Basin are 329 essentially controlled by silicate weathering. Dissolved Li isotopic compositions are however 330 highly variable within the Mackenzie river basin (from +9.3 to +29.0%), suggesting variable 331 332 fractionation of Li isotopes during silicate weathering.

333

334 **5.2. Li isotopes and silicate weathering regimes**

335

336

5.2.1. Evidence for isotope fractionation during chemical weathering

337

River dissolved loads are systematically enriched in ⁷Li relative to bedrocks and suspended 338 339 particles (Figure 4). In this regard, lithium is similar to boron (Lemarchand and Gaillardet 2006). However, the dissolved loads exhibit a large range of δ^7 Li values across the sampled 340 rivers. A part of this variability may derive from the δ^7 Li of bedrocks, which might differ 341 between the main provinces of the Mackenzie Basin. Literature data indicate that the Li 342 isotopic composition of continental granites (-2 to +3‰, Teng et al. 2004, 2006, 2008, 2009) 343 is slightly different from the shales and loesses from which they were derived (-3 to +5‰). 344 345 The Li isotopic composition of the Canadian Shield bedrock (mainly consisting of granites

and associated crystalline rocks, Millot et al. 2002) is averaged in the glacial till samples 346 reported in Table 3. In concordance with literature data, the δ^7 Li values measured in till 347 samples are higher than those measured in the bedrocks from the lowlands and the Rocky 348 349 Mountains (which are mainly recycled materials such as shales, blackshales, slates and a few low grade metamorphic rocks). In Figure 4, we report δ^7 Li values in silicate bedrocks 350 compiled from the literature together with the results from this study. Suspended sediment 351 352 and sand samples from the Mackenzie rivers fall within the range of bedrocks defined by our bedrock samples and by literature data (Figure 4). However, a positive trend between $\delta^7 Li$ 353 and Na/AI is observed (Figure 5). Due to the high solubility of Na during chemical weathering 354 compared to AI which is immobile, the Na/AI ratio can be regarded as "a weathering index". 355 The low Na/AI ratios associated with low δ^7 Li values suggests that lithium isotopes are 356 fractionated by chemical weathering processes. The direction of the fractionation in the 357 358 suspended sediments is consistent with the enrichment of ⁷Li in the dissolved load since the lowest Na/AI are associated with the lowest δ^7 Li. Since the suspended sediments of the 359 Mackenzie rivers are essentially derived from sedimentary rocks (the Shield area does not 360 361 contribute significant sediment), it is difficult to conclude whether the Na loss recorded in the Na/AI ratio is due to present day weathering conditions, or whether the low Na/AI ratios are 362 inherited from previous cycles of erosion and sedimentation. Examples of shale composites 363 reported in the literature, such as the PAAS or NASC (Post Archean Australian Shale and 364 365 North American Shale Composite: Holland 1978, Taylor and MacLennan 1985) do indeed show Na depletion compared to granodioritic Upper Continental Crust (Taylor and 366 367 MacLennan 1985, Gaillardet et al. 1999). Given the low grade silicate weathering in the Mackenzie River Basin (Millot et al. 2003), and the overlap of bedrock and suspended 368 369 sediment Li isotopes compositions (Figure 4), we favour the hypothesis that Na/AI in the 370 suspended sediments essentially records bedrock geochemistry rather than present day 371 weathering products. Therefore the trend of increasing Li isotopic composition with Na/AI ratio (Figure 5) is likely inherited from the bedrock history. From a mass budget perspective, 372

it is not easy to estimate the partitioning of Li transported in the dissolved and particulate 373 load because chemical weathering does not proceed at steady state (Vigier et al. 2001). 374 375 However, the above considerations suggest that the dissolved load clearly reflects the fractionation of Li isotopes during chemical weathering (with the enrichment of ⁷Li in the 376 fluid), while the suspended phase is not significantly different from the bedrock. In other 377 words, the dissolved load is sensitive to water rock interactions, but solids are not, because 378 379 chemical weathering is too weak. Investigating in more detail the isotopic composition of Li in 380 river suspended sediments as a function of weathering rate is beyond the scope of the present paper, but would be interesting to do at large scale. The next section focuses on the 381 dissolved load. 382

383

384

5.2.2. Influence of the weathering regime

385

We show in the following that the range of Li isotopic composition measured in the rivers of the Mackenzie Basin is related to the weathering regime.

The Li/Na^{*} ratio in the dissolved load, when compared to that in the bedrock can be used as an index of Li mobility compared to Na. Both elements are alkali metals, thus mobile during water-rock interactions. As shown by a number of previous studies, in contrast to Na, Li is likely to be incorporated into secondary minerals, such as clays. By analogy with the study of Georg et al. (2007) concerning the mobility of Si isotopes during chemical weathering in lceland , we can define f_{Li} , the fraction of Li remaining in solution relative to Na, as:

394

395

$$f_{Li} = \frac{(Li / Na^*)_{dissolved}}{(Li / Na)_{bedrock}}$$
(1)

396

where Na^{*} is the Na concentration corrected for atmospheric input and for evaporite dissolution using the results of Millot et al. (2003). Correction for evaporite dissolution only significantly affects a few samples. A value of 1 for f_{Li} means that chemical weathering is 400 congruent. While dissolution is usually thought to be incongruent, precipitation of secondary
401 phases is the most probable process that will fractionate Li and Na, Li being known to have
402 affinity for clay minerals, while Na is much more mobile and hardly reincorporated into clay
403 minerals.

404 We used the suspended sediment compositions, where available, to estimate the bedrock Li/Na ratio. In the Canadian Shield province, because rivers at the time of sampling 405 contained very low levels of suspended sediment, we used two glacial till samples (Table 3) 406 407 which display contrasting Li/Na ratios. The calculated fLi in the rivers of the Mackenzie basin range between 0.1 (in the lowlands and main tributaries) to 0.8 in the rivers of the Shield 408 area. Intermediate values are found in the Rocky Mountains and in the Western Cordillera. 409 This indicates that the reincorporation of Li into secondary phases is more important in the 410 411 lowlands and main tributaries than in the Cordillera and Rocky Mountain samples. In Figure 6, the dissolved Li/Na in the Shield rivers were normalized to the Prosperous Lake till 412 sample. If the till sample CAN96-31 (Yellowknife) is used as a reference for the parent 413 414 bedrock in the Slave Province, then f_{Li} values higher than 1 are found. Values of f_{Li} higher 415 than 1 would mean a preferential leaching of Li compared to Na. Although Prosperous Lake till sample is our preferred bedrock estimate here, we nevertheless consider fLi higher than 1 416 417 (and CAN96-31 as a source rock) as a possible scenario. Despite these uncertainties on the bedrock in the Shield province, f_{Li} provides a relative indication of Li mobility during 418 419 weathering.

When the isotopic composition of dissolved lithium is plotted as a function of f_{Li} , a correlation appears (Figure 6). This correlation can be explained by a binary mixing between waters draining areas characterized by two markedly different weathering regimes.

The first end-member needed to account for the correlation between f_{Li} and $\delta^7 Li$ of the dissolved load corresponds to a weathering regime where Li is significantly precipitated in the solids compared to Na, most probably incorporated into Li-rich secondary minerals (clays). The lowlands and the transition zone areas best characterize this weathering regime since they display the lowest f_{Li} values. Numerous studies have shown that Li isotopes are

fractionated during the formation of secondary minerals. The ~13‰ isotopic shift between the 428 composition of the dissolved load and the bedrock in the lowlands is, to a first order, 429 430 compatible with the estimated clay-solution isotopic fractionation at equilibrium (Taylor and Urey 1938, Chan et al. 1992, Zhang et al. 1998, Pistiner and Henderson 2003, Vigier et al. 431 2008). The formation of secondary solids, being kinetically limited, requires a favorable 432 environment. Using boron isotopes, Lemarchand and Gaillardet (2006) have suggested that 433 434 most of the dissolved boron in rivers is derived from the shallow groundwater system (drift 435 aquifers). As shown by Lemay (2002), high lithium concentrations (3-15 µmol/L) are also measured in these groundwaters. The correlation between boron and lithium concentrations 436 that links both rivers and groundwaters (Figure 7) strongly argues for a significant input of Li 437 to the lowland rivers from groundwaters. Because groundwaters have long residence times 438 relative to surface waters, chemical weathering reactions can proceed further than in the 439 Shield or Rocky Mountains areas where the residence time of water is not sufficiently long 440 (Lemarchand and Gaillardet 2006). We therefore suggest that the residence time of water in 441 442 groundwater counteracts the effect of low temperature and favors the formation of secondary 443 minerals that fractionate Li isotopes (Figure 6) at equilibrium.

The second endmember in Figure 6 corresponds to the Shield and Rocky Mountains areas, 444 and displays the highest δ^7 Li and f_{Li} values. This situation may appear as a paradox because 445 fLi indicates no or low Li incorporation in secondary products of weathering, while the isotopic 446 shift between Li in the bedrock and dissolved Li is at its maximum (from +16 to +22‰). The 447 448 Rocky Mountains and the Shield province are areas characterized by particularly low chemical weathering rates (Millot et al. 2003), in contrast to the lowlands. These regions 449 have been deglaciated relatively recently and are characterized by abundant bare rocks and 450 451 poorly developed soils. The Rocky and Mackenzie Mountains areas have relatively high river 452 sediment yields with sediments mostly derived from glacial erosion (Millot et al. 2003) and not from soils. The Shield area is characterized by important glacial till deposits. Several 453 hypotheses can be invoked to explain the enrichment of ⁷Li in the waters of the Rocky 454 Mountains and of the Shield area. The most evident possibility is that the formation of some 455

secondary mineral fractionating Li isotopes. As indicated by the fLi values close to 1, the 456 incorporation of Li into secondary phases should not be quantitatively important in these 457 458 regions but should however be visible in terms of isotope fractionation. A good candidate is Mn and Fe-oxyhydroxide minerals. Adsorption of Li onto Fe and Mn oxyhydroxide surfaces 459 have been reported by Chan and Hein (2007) with preferential enrichment of ⁶Li in the 460 deposit, thus consistent with the observed heavy isotope enrichment in the fluids. A recent 461 462 study reports the enrichment of the heavy isotope of Li in the dissolved load of glacial and non-glacial rivers from Greenland and attributes it to the preferential uptake of ⁶Li during 463 inner sphere sorption of Li⁺ on the Fe-oxyhydroxide surfaces (Wimpenny et al. 2010). The 464 same mechanism can be invoked here. The difference between the study of Wimpenny et al. 465 (2010) and our study is that the Shield rivers, that have ⁷Li-enriched waters, are not sulfate-466 rich and therefore that sulfide oxidation, a mechanism proposed by the authors to explain the 467 abundance of Fe-oxides in Greenland, is not applicable to the Shield rivers. The presence of 468 Fe-oxyhydroxides minerals in the Shield rivers could be tested by analysing Mn and Fe river 469 470 deposits often found in river bedloads, at the surface of blocks and boulders for example. In 471 addition, saturation state calculations (using PHREEQC modelling) indicate that the waters are supersaturated with respect to goethite in this area. 472

A second possibility to explain the enrichment of ⁷Li in the dissolved load of the Shield and 473 474 Mackenzie Mountain rivers could be a fractionation of Li isotopes during mineral dissolution. 475 Experimental work on basalt dissolution does not show evidence of Li isotope fractionation during dissolution (Pistiner and Henderson 2003), but we can imagine a number of natural 476 processes potentially able to fractionate Li isotopes during mineral dissolution. The 477 preferential release of ⁷Li could be possible either by the preferential dissolution of ⁷Li-478 479 enriched phases, or as a transient effect of the leaching process with formation of unstable secondary phases such as Si-gel layers around minerals. So far, Li isotope heterogeneity at 480 the mineral scale has not been observed within a single bedrock, but we cannot exclude the 481 existence of low temperature minerals that would show ⁷Li enrichments. What is interesting 482 here, is that the ⁷Li enrichment in waters is observed in the geological context of glacial 483

484 erosion (still active in the Rockies and in the form of glacial deposits in the Shield area). The 485 abundance of fine glacial flour allows specific weathering reactions to occur, that have been 486 observed experimentally (in columns of grind granitic material, White et al. 1998) or in nature 487 (Anderson et al. 1989). In particular fast dissolving accessory phases like carbonates and 488 sulfides have been shown to contribute significantly to solute export in the early stages of 489 weathering.

490 Whatever the reason for Li isotope fractionation, our data do show that incipient weathering of rocks can produce ⁷Li enriched fluids. The similar enrichments found in Greenland 491 (Wimpenny et al. 2010) or in very recently deglaciated moraine in the granitic Swiss Alps 492 (Lemarchand et al. 2009) shows that Li isotopes are very sensitive to the early stages of 493 chemical weathering. Future studies will be necessary to investigate Li isotope fractionation 494 in incipient weathering conditions and the isotopic heterogeneity of bedrocks to precisely 495 determine the mechanisms that result in the preferential release of heavy Li isotopes in 496 497 solution.

According to the above discussion, all sampled rivers in the Mackenzie Basin have an 498 499 isotopic composition reflecting a mixture of water masses that have interacted with bedrock minerals according to two contrasting weathering regimes. On one hand, oxyhydroxide 500 formation and/or leaching of comminuted bedrock produces ⁷Li-enriched waters, similarly to 501 502 that occurring in glacial and non glacial rivers from Greenland. On the other hand, the more 503 optimum weathering conditions of the low-lying plain aquifers will produce waters with Li isotopic compositions lower by about 20%. As indicated by the previous studies of our group, 504 the weathering rates of silicates in these two contrasted environments differ by a factor of 3-505 4, probably accelerated in the lowlands by the contribution of groundwaters (Lemarchand 506 507 and Gaillardet 2006) and the chelating effects of organic matter (Millot et al. 2003). The Li isotopic data presented here show that the Li isotopic signature of waters is more influenced 508 by the precipitation of specific secondary phases than weathering rates. The relationship 509 between chemical weathering rates of silicates (calculated from the inversion of major 510 element concentrations and Sr isotopic ratios by Millot et al. 2003), and δ^7 Li measured in the 511

dissolved loads is shown on Figure 8a. There is no clear relationship at the scale of the 512 Mackenzie River Basin between the Li isotopic composition and the specific flux of chemical 513 514 denudation, showing that secondary mineral formation is not necessarily related to denudation fluxes. In particular, the high fluxes observed for the Western Cordillera and 515 visible in Figure 8 are due to the combined effects of runoff and the volcanic nature of 516 bedrocks (Gaillardet et al. 2003). The absence of a relationship between chemical 517 denudation rate and water lithium isotopic composition is in contrast with the relationship 518 observed on small monolithologic rivers in Iceland (Vigier et al. 2009). 519

Two trends of δ^7 Li with total dissolved load derived from silicate weathering in mg/L are 520 observed (Figure 8b), corresponding to the two weathering regimes described above. The 521 first trend of increasing ⁷Li-enrichment with dissolved solutes only contains samples from the 522 Shield and Rocky-Mackenzie Mountains from the first weathering regime described above. 523 524 By contrast, the waters most influenced by the second weathering regime show constant Li isotopic composition with total dissolved load concentration derived from silicate weathering. 525 Figure 8b confirms the broad idea that Li isotopes are indicative of the regime weathering. 526 Regimes characterized by secondary clay formation have Li isotopic composition between 10 527 528 and 15‰ and show no relationship with silicate weathering rates, while weathering regimes characterized by incipient weathering, or weathering of comminuted bedrocks, are 529 characterized by higher Li isotopic composition. 530

531

532

533

6. CONCLUSIONS

The Li isotopic composition measured in the rivers of the Mackenzie Basin shows that ⁷Li is enriched in the dissolved load and that δ^7 Li can vary by 20‰ within a large river basin. The δ^7 Li of the particulate load reflects that of the bedrock and ranges between -2 and +3‰. Our study shows that dissolved Li in river waters is essentially derived from the weathering of silicates and that Li isotopic ratios of the dissolved load depends on the weathering regime of silicates.

If Na is considered as a soluble element not incorporated in secondary phases, river waters 540 showing Li/Na ratios much lower than the bedrock ratio have the lowest isotopic 541 542 compositions. Although the fractionation coefficients between solution and newly formed solids remain poorly known and need to be better constrained by experimental work, our 543 isotopic data are consistent with the incorporation of Li into clay minerals at equilibrium. 544 Groundwaters, in particular shallow aguifers in the low-lying plains of the basin, seem to play 545 546 an important role in the acquisition of the dissolved Li isotope composition, probably because high residence times favor the formation of clay mineral. The most ⁷Li-enriched waters are 547 observed in the Rocky Mountains and in the Shield areas of the basin. In these areas, by 548 analogy to previous work in Greenland (Wimpenny et al. 2010), we suggest that high $\delta^7 Li$ 549 values in rivers result from the fractionation associated with Li sorption onto Mn and Fe-550 oxyhydroxide surfaces. However, another mechanism could be that leaching is likely to be a 551 552 non negligible process of water-rock interaction due to surficial or incipient weathering, being able to produce ⁷Li-enriched waters. Experimental work will be necessary for better 553 understanding the behaviour of Li isotopes in incipient weathering regimes. 554

To summarize, this large scale study shows that dissolved Li is essentially derived from 555 556 silicate weathering but shows that silicate weathering does not have a unique isotopic signature. A very large variability of Li isotopic ratios exists within the Mackenzie Basin and is 557 strongly spatialized. This is an important result if Li isotopes in the ocean are to be used as a 558 proxy for the secular evolution of the Earth's surface. This study suggests that, rather than 559 560 weathering fluxes, the weathering regime controls the Li isotopic signature of silicate weathering. Future work will have to address the role of the different parameters that define 561 the weathering regime: time, climate, hydrological setting, residence time of water, biological 562 563 parameters.

564

565 Acknowledgements:

566

This work was supported by the French program funded by the INSU-CNRS (PNSE 567 contribution 322). We would like to express special thanks to B. Dupré for measurements of 568 trace elements in Toulouse University. We also would like to acknowledge B. Dupré, C. 569 Gariépy and D. Lemarchand for their help in collecting river samples. We thank E. 570 571 Lemarchand and E.T. Tipper for fruitful discussions. RM would like to thank Ph. Négrel and T.D. Bullen for comments in an earlier version of the manuscript and the Research Division 572 of BRGM is also acknowledged for funding. We thank R. James, P. Pogge van Strandmann 573 and B. Reynolds for providing critical comments that improved this manuscript. D. Vance is 574 575 also thanked for editorial handling and constructive comments. Finally, we also would like to thank P. Burnard and R.H. Hilton for comments and English corrections. This is IPGP 576 contribution n° 2581 and BRGM contribution n° 6327. 577

578

- 580 **References**
- 581
- 582 Anderson M.A., Bertsch P.M., and Miller W.P. (1989) Exchange and apparent fixation of 583 Lithium in selected soils and clay minerals. Soil Sci., 148, 46-52.
- Anghel I., Turin H.J., and Reimus P.W. (2002) Lithium sorption to Yucca mountain tuffs.
 Appl. Geochem., 17, 819-824.
- Calmels D., Gaillardet J. Brenot A., France-Lanord C. (2007) Sustained sulfide oxidation by
 physical erosion processes in the Mackenzie River basin: Climatic perspectives. Geology,
 35: 1003-1006.
- Carignan J., Cardinal D., Eisenhauer A., Galy A., Rehkämper M., Wombacher F., Vigier N.
 (2004) A reflection on Mg, Ca, Cd, Li and Si isotopic measurements and related reference
 materials. Geostandards and Geoanalytical Research, 28: 139-148.
- Carignan J., Vigier N., Millot R. (2007) Three secondary reference materials for Li isotopic
 measurements: ⁷Li-N, ⁶Li-N and LiCl-N. Geostandards and Geoanalytical Research, 31: 7 12.
- 595 Chan L.H., Edmond J.M. Thompson G., Gillis K. (1992) Lithium isotopic composition of
 596 submarine basalts: implications for the lithium cycle to the ocean, Earth Planet. Sci. Lett.,
 597 108: 151-160.
- Chan L.H., Hein J.R. (2007) Lithium contents and isotopic compositions of ferromanganese
 deposits from the global ocean. Deep-sea research part II topical studies in
 Oceanography, 54: 1147-1162.
- 601 Flesch G.D., Anderson A.R., Svec H.J. (1973) A secondary isotopic standard for ⁶Li/⁷Li
- 602 determinations. International Journal of Mass Spectrometry and Ion Physics, 12: 265-272.
- Gaillardet J., Dupré B., Louvat P., Allègre C.J. (1999) Global silicate weathering and CO₂
 consumption rates deduced from the chemistry of large rivers. Chemical Geology, 159: 330.
- Gaillardet J., Millot R., Dupré B. (2003) Chemical denudation rates of the western Canadian
 orogenic belt: the Stikine terrane. Chemical Geology, 201: 257-279.

- Georg R.B., Reynolds B.C., West A.J., Burton K.W., Halliday A.N. (2007) Silicon isotope
 variations accompanying basalt weathering in Iceland. Earth and Planetary Science
 Letters, 261: 476-490.
- Hall J.M., Chan L.H. (2004) Li/Ca in multiple species of benthic and planktonic foraminifera:
- 612 thermocline, latitudinal, and glacial-interglacial variation. Geochimica et Cosmochimica613 Acta, 68: 529-545.
- Hall J.M., Chan L.H., McDonough W.F., Turekian K.K. (2005) Determination of the lithium
 isotopic composition of planktic foraminifera and its application as a paleo-seawater proxy.
 Marine Geology, 217: 255-265.
- Hathorne E.C., James R.H. (2006) Temporal record of lithium in seawater: A tracer for
 silicate weathering? Earth and Planetary Science Letters, 246: 393-406.
- Hathorne E.C., James R.H. Lampitt R.S. (2009) Environmental versus biomineralization
 controls on the intratest variation in the trace element composition of the planktonic
 foraminifera G. inflata and G. scitula. Paleoceanography, 24: PA4204.

Holland H.D. (1978) The chemistry of oceans and atmosphere, Wiley and Sons, New York.

- Huh Y., Chan L.C., Zhang L., Edmond J.M. (1998) Lithium and its isotopes in major world
 rivers: implications for weathering and the oceanic budget. Geochimica et Cosmochimica
 Acta, 62: 2039-2051.
- Huh Y., Chan L.C., Edmond J.M. (2001) Lithium isotopes as a probe of weathering
 processes: Orinoco River. Earth and Planetary Science Letters, 194: 189-199.
- Huh Y., Chan L.C., Chadwick O.A. (2004) Behavior of lithium and its isotopes during
 weathering of Hawaiian basalt. Geochemistry, Geophysics, Geosystems, 5: 1-22.
- James R.H., Palmer M.R. (2000) The lithium isotope composition of international rock
 standards. Chemical Geology, 166: 319-326.
- Jeffcoate A.B., Elliott T., Thomas A., Bouman C. (2004) Precise, small sample size
 determinations of lithium isotopic compositions of Geological Reference Materials and
 modern seawater by MC-ICP-MS. Geostandards and Geoanalytical Research, 28: 161 172.

- Kisakürek B., Widdowson M., James R.H. (2004) Behaviour of Li isotopes during continental
 weathering: the Bidar laterite profile, India. Chemical Geology, 212: 27-44.
- 638 Kisakürek B., James R.H., Harris N.B.W. (2005) Li and δ^7 Li in Himalayan rivers: Proxies for 639 silicate weathering? Earth and Planetary Science Letters, 237: 387-401.
- Lear C.H., Rosenthal Y. (2006) Benthic foraminiferal Li/Ca: insights into Cenozoic seawater
 carbonate saturation state. Geology, 34: 985-988.
- Lemarchand D., Gaillardet J. (2006) Transient features of the erosion of shales in the
 Mackenzie basin (Canada), evidences from boron isotopes. Earth and Planetary Science
 Letters, 245: 174-18.
- Lemarchand E., Tipper E.T., Hindshaw R., Wiederhold J.G., Reynolds B.C., Bourdon B.,
 Kretzchmar R. (2009) Li isotope fractionation in surface waters of an alpine granitic
 catchment. Goldschmidt Conference Abstract A742.
- Lemay T.G. (2002) Geochemical and isotope data for formation water from selected wells,
 cretaceous to quaternary succession, Athabasca oil sand (in situ) area, Alberta, EUB/AGS
 Geo-Note 2002-02: 1-30.
- Millot R., Gaillardet J., Dupré B., Allègre C.J. (2002) The global control of silicate weathering
 rates and the coupling with physical erosion: new insights from the Canadian Shield. Earth
 and Planetary Science Letters, 196: 83-98.
- Millot R., Gaillardet J., Dupré B., Allègre C.J. (2003) Northern latitude chemical weathering
 rates: clues from the Mackenzie River Basin, Canada. Geochimica et Cosmochimica Acta,
 67: 1305-1329.
- Millot R., Guerrot C., Vigier N. (2004) Accurate and high-precision measurement of lithium
 isotopes in two reference materials by MC-ICP-MS. Geostandards and Geoanalytical
 Research, 28: 153-159.
- Négrel Ph., Allègre C.J., Dupré B., Lewin E. (1993) Erosion sources determined by inversion
 of major and trace element ratios and strontium isotopic ratios in river water: the Congo
 Basin case. Earth and Planetary Science Letters, 120: 59-76.

- Pistiner J.S., Henderson G.M. (2003) Lithium isotope fractionation during continental
 weathering processes. Earth and Planetary Science Letters, 214: 327-339.
- Pogge von Strandmann P.A.E., Burton K.W., James R.H., van Calsteren P., Gislason S.R.,
- 666 Mokadem F. (2006) Riverine behaviour of uranium and lithium isotopes in an actively 667 glaciated basaltic terrain. Earth and Planetary Science Letters, 251: 134-147.
- 668 Reeder S.W., Hitchon B., Levinson A.A. (1972) Hydrogeochemistry of the surface waters of
- the Mackenzie River drainage basin, Canada-I. Factors controlling inorganic composition.
 Geochimica et Cosmochimica Acta, 36: 825-865.
- Rudnick R.L., Tomascak P.B., Njo H.B., Robert Gardner L. (2004) Extreme lithium isotopic
 fractionation during continental weathering revealed in saprolites from South Carolina.
 Chemical Geology, 212: 45-57.
- Taylor S.R., Urey H.C. (1938) Fractionation of the lithium and potassium isotopes by
 chemical exchange with zeolites, J. Chem. Phys. 6: 429-438.
- Taylor S.R., MacLennan S.M. (1985) The continental crust: its composition and evolution,
 Blackwell Scientific Publications, 312 p.
- Teng F.Z., McDonough W.F., Rudnick R.L., Dalpé C., Tomascak P.B., Chappell B.W., Gao
- S. (2004) Lithium isotopic composition and concentration of the upper continental crust.
 Geochimica et Cosmochimica Acta, 68: 4167-4178.
- Teng F.Z., McDonough W.F., Rudnick R.L., Walker R.J. (2006) Diffusion-driven extreme
 lithium isotopic fractionation in country rocks of the Tin Mountain pegmatite. Earth and
 Planetary Science Letters, 243: 701-710.
- Teng F.Z., Rudnick R.L., McDonough W.F., Gao S., Tomascak P.B., Liu Y. (2008) Lithium
 isotopic composition and concentration of the deep continental crust. Chemical Geology,
 255: 47-59.
- Teng F.Z., Rudnick R.L., McDonough W.F., Wu F.Y. (2009) Lithium isotopic systematics of
 A-type granites and their mafic enclaves: Further constraints on the Li isotopic
 composition of the continental crust. Chemical Geology, 262: 370-379.

- Tomascak P.B. (2004) Developments in the understanding and application of lithium
 isotopes in the Earth and Planetary Sciences. Reviews in Mineralogy & Geochemistry, 55:
 153-195.
- Vigier N., Bourdon B., Turner S., Allègre C.J. (2001) Erosion timescales derived from U-decay
 series measurements in rivers, Earth and Planetary Science Letters, 193: 549-563.
- Vigier N., Rollion-Bard C., Spezzaferri S. and Brunet F. (2007) In-situ measurements of Li
 isotopes in foraminifera. Geochemistry, Geophysics, Geosystems Q01003.
- Vigier N., Decarreau A., Millot R., Carignan J., Petit S., France-Lanord C. (2008) Quantifying
 Li isotope fractionation during smectite formation and implications for the Li cycle,
 Geochimica et Cosmochimica Acta, 72: 780-792.
- Vigier N., Gislason S.R., Burton K.W., Millot R., Mokadem F. (2009) The relationship
 between riverine lithium isotope composition and silicate weathering rates in Iceland.
 Earth and Planetary Science Letters, 287: 434-441.
- White A.F., Blum A.E., Schulz M.S., Vivit D.V., Larsen M., Murphy S.F. (1998) Chemical
 weathering in a tropical watershed, Luquillo Mountains, Puerto Rico: I. Long-term versus
 short-term chemical fluxes. Geochimica et Cosmochimica Acta, 62: 209-226.
- Wimpenny J., James R.H., Burton K.W., Gannoun A., Mokadem F., Gíslason S.R. (2010)
 Glacial effects on weathering processes: New insights from the elemental and lithium
 isotopic composition of West Greenland rivers. Earth and Planetary Science Letters, 290:
 427-437.
- Zhang L., Chan L.H., Gieskes J.M. (1998) Lithium isotope geochemistry of pore waters from
 Ocean Drilling Program Sites 918 and 919, Irminger Basin, Geochim. Cosmochim. Acta
 62: 2437-2450.

713 **Figure captions** 714 715 Figure 1 Map showing the sample locations within the Mackenzie River Basin and adjacent basins 716 (Yukon, Stikine, Nass, Skeena and Fraser Basin). 717 718 719 Figure 2 720 Inverse correlation of Li isotopic composition and Li concentration in the dissolved load. In this diagram, the geomorphorphic provinces of the Mackenzie river basin are well 721 distinguished. 722 723 Figure 3 724 Histogram for Li contribution coming from silicate weathering (%) in Mackenzie river water 725 726 samples. 727 Figure 4 728 Histogram of Li isotopic composition in the dissolved and suspended loads of the Mackenzie 729 730 rivers. Bedrock values from this study (4 values) and from a compilation of literature data on 731 shale, granite and loess (Teng et al. 2004) are added for comparison. This figure clearly shows the ⁷Li enrichment of the dissolved phase compared to bedrocks, while suspended 732 733 sediments and bedrocks are essentially similar. 734 Figure 5 735 Lithium isotopic composition in the suspended sediments of the Mackenzie rivers (main 736 tributaries, Rocky and Mackenzie Mountains, Interior Platform and transition mountain/plain) 737 and glacial till sample (Canadian Shield) as a function of Na/Al ratio. This ratio can be seen 738

can be attributed either to present day weathering processes or to previous episodes of

739

27

as a proxy for Na loss compared to AI, and is therefore a weathering intensity proxy. Na loss

741 weathering that would have occurred during the geological history of the sedimentary742 bedrocks.

743

744 **Figure 6**

Lithium isotopic composition in the dissolved load as a function of f_{Li} . f_{Li} represents the 745 relative mobility of Li with respect to Na, an element not generally known to be 746 747 reincorporated into secondary phases. This parameter is defined as the ratio of Li/Na in the 748 dissolved load to that in the suspended sediments or, in the case of the Shield province, to 749 the Prosperous Lake moraine. Na* denotes here that the Na concentrations are corrected for atmospheric input and, for a couple of rivers, are corrected for evaporite dissolution 750 according to the results of Millot et al. (2003). Ratios in suspended sediments are considered 751 to represent bedrock ratios. For the Shield rivers, if the glacial till (sample CAN96-31) is used 752 for normalizing the Li/Na data from the shield rivers then fLi values higher that 1 would be 753 obtained for these rivers (not shown on the figure), but lithium isotopic composition and fLi are 754 755 still correlated.

756

757 Figure 7

Relationship between boron and lithium concentrations in the dissolved load of the rivers sampled in this study. Data from drift aquifers reported by Lemay et al. (2002) have been added to show that the contribution from a groundwater input is likely. Boron data are from Lemarchand and Gaillardet (2006). Boron isotopic data strongly suggest that boron isotopic composition was acquired during reactive transport of boron in shallow groundwaters of relatively high water residence time, and that, boron is therefore a good tracer of the groundwater contribution.

765

766 **Figure 8**

Li isotopic composition as a function of Silicate Weathering Rate (Figure 8a) and Silicate
Weathering Load (Figure 8b) for the river samples. Silicate weathering rates were calculated

in Millot et al. (2003) based on the inversion of major element concentration and Sr isotope data in the dissolved load, and by Gaillardet et al. (2003) for the Stikine province. In Figure 8b, the observed feature is consistent with the idea developed here that the Li isotopic composition of water in the Mackenzie basin rivers reflects mixing between waters modified by weak (incipient or surficial) weathering (weathering regime #1) and waters dominated by more intense water-rock interactions with the formation of secondary phases in the lowlands of the Mackenzie River Basin (weathering regime #2). 776 **Table captions**

777

778 Table 1

Lithium isotopic composition δ^7 Li (‰) and concentrations (µmol/L) for rivers waters of the Mackenzie Basin and the W. Cordillera. Analytical precision for each δ^7 Li measurement is reported in this table and range from 0.1 to 0.3‰ ($2\sigma_m$). As mentioned in the analytical section, the total reproducibility is ± 0.5‰ (2σ) and has been determined by long-term repeated measurements of a seawater sample (Millot et al. 2004). Surface area (km²), river discharge (km²/year), runoff (mm/year) and Silicate Weathering Rates (tons/km²/yr and mg/L) have also been added in this table (from Millot et al. 2003).

786

787 Table 2

Lithium isotopic composition δ^7 Li (‰) and concentrations (µg/g) in suspended sediments for rivers from the Mackenzie Basin and adjacent basins draining the Canadian orogenic belt (W. Cordillera). Na and Al concentrations are also given in µg/g.

791

792 Table 3

 δ^{7} Li (‰) and Li concentrations (µg/g) in sand samples from the Mackenzie River at the river mouth (CAN96-6), as well as in carbonate, blackshales, and two glacial tills sampled on the Canadian Shield.

796

798 Figure 1



























838 Figure 8



Tributaries	sample number	River name	8 ⁷ ⊔i 2 o	5	Area	Discharge	Runoff	Silicate Weat	nering Rate	I.
			%	hmol/L	km²	km³/year	mm/year	tons/km²/year	_ mg/L	
Main tributaries	CAN96-5	Peel (FortMcPherson)	+ 13.3 0.1	0.71	70600	21.65	307	1.73	5.63	1
	CAN96-6	Mackenzie (Tsiigehtchic)	+ 15.0 0.2	0.57	1680000	283.70	169	1.03	6.12	
	CAN96-7	Red Arctic (Tsiigehtchic)	+ 13.8 0.1	0.89	18600	5.02	270	1.37	5.08	
	CAN96-15	Peace (PeaceRiver)	+ 15.2 0.2	0.23	186000	57.75	310	0.92	2.96	
	CAN96-25	Mackenzie (FortSimpson)	+ 16.5 0.3	0.53	1270000	137.90	139	1.05	7.57	
	CAN96-26	Liard (FortSimpson)	+ 14.3 0.1	0.63	275000	77.32	281	1.19	4.24	
	CAN96-37	Peace (PeacePoint)	+ 15.4 0.1	0.52	305500	67.01	219	1.30	5.92	
	CAN96-38	Slave (FortSmith)	+ 15.5 0.2	09.0	616400	100.98	164	1.14	6.94	
	CAN96-42	Athabasca (FortMcMurray)	+ 12.7 0.2	0.86	131000	20.86	159	1.94	12.19	
Rocky and Mackenzie Mountains	CAN96-4	Yukon (Dawson)	+ 17.9 0.2	0.32						I I
	CAN96-19	Racing	+ 12.2 0.1	0.43	1900	2.30	1209	0.65	0.53	
	CAN99-5	Athabasca (Hinton)	+ 15.4 0.2	0.45	9780	5.46	558	0.49	0.88	
	CAN96-21	Liard (LiardRiver)	+ 12.9 0.1	0.36	33400	11.71	351	0.79	2.24	
	CAN99-21	Fraser	+ 17.5 0.1	0.11	•					
	CAN99-46	Liard (UpperLiard)	+ 20.5 0.1	0.19	33400	11.71	351	0.64	1.83	
	CAN99-54	Smith	+ 12.6 0.1	0.19	3740	0.77	207	0.13	0.62	
	CAN99-56	Liard (LiardRiver)	+ 18.3 0.2	0.19						
	CAN99-60	Toad	+ 15.8 0.1	0.50	2570	1.37	532	0.55	1.04	
	CAN99-64	Racing	+ 13.0 0.1	0.36	1900	2.30	1209	0.65	0.54	
	CAN99-74	Peace (Hudson'sHope)	+ 17.0 0.2	0.18	00669	34.71	497	0.89	1.79	
Interior Platform	CAN96-14	Little Smoky	+ 11.8 0.2	0.69	3010	0.77	257	2.42	9.43	1
	CAN96-33	Hay (mouth)	+ 16.3 0.1	1.29	47900	3.60	75	1.34	17.84	
	CAN96-40	Lesser Slave	+ 10.1 0.1	1.21	13600	1.32	97	1.28	13.21	
	CAN99-1	Pembina (Evansburg)	+ 16.8 0.1	0.53	2900	1.12	386	4.33	11.21	
	CAN99-2	Mc Leod	+ 15.4 0.1	0.85	9100	2.11	232	2.71	11.68	
	CAN99-71	Doig	+ 13.3 0.1	1.02			,			
	CAN99-72	Beatton	+ 11.4 0.1	0.51			,			
	CAN99-79	Simonette	+ 11.4 0.1	1.02	5050	1.16	229	3.94	17.17	
	CAN99-80	Little Smoky	+ 12.9 0.1	0.58	3010	0.77	257	2.38	9.26	
Transition mountain/plain	CAN96-13	Athabasca	+ 14.8 0.2	0.29						1
	CAN96-16/1	Smoky	+ 12.1 0.1	0.56	51300	11.83	231	1.90	8.22	
	CAN96-23	Fort Nelson	+ 11.8 0.1	0.89			,		5.98	
	CAN99-78	Smoky	+ 9.3 0.1	0.41	51300	11.83	231	0.80	3.48	
Canadian Shield	CAN96-31	Yellowknife	+ 26.9 0.1	0.17	16300	1.16	71	0.27	3.80	
	CAN99-84	Wecho	+ 26.0 0.2	0.20	3400	0.31	06	0.39	4.28	
	CAN99-86	Wopmay	+ 25.2 0.1	0.07		·	·	0.33	2.83	
	CAN99-90	Yellowknife	+ 29.0 0.1	0.19	16300	1.16	71	0.28	3.87	
W. Cordillera	CAN99-28	Skeena	+ 24.9 0.2	0.05	42200	30.36	719	4.20	5.84	1
	CAN99-30	Nass	+ 21.6 0.1	0.09	19200	24.68	1285	5.00	3.89	
	CAN99-39	Stikine	+ 21.7 0.1	0.10	29300	23.83	811	4.10	5.06	

Table 1

Table 2

Tributaries	sample number	River name	δ ⁷ Li	$2\sigma_{m}$	Li µg/g	Na µg/g	Al µg/g
Main tributaries CAN96-6		Mackenzie (Tsiigehtchic)	- 0.9	0.2	57.8	3710	70200
	CAN96-7	Red Arctic (Tsiigehtchic)	- 1.6	0.1	56.8	2819	73535
	CAN96-26	Liard (FortSimpson)	- 1.3	0.2	46.1	3339	71206
	CAN96-38	Slave (FortSmith)	- 0.2	0.1	41.2	5268	67447
Rocky and Mackenzie Mountains	CAN99-5	Athabasca (Hinton)	+ 2.6	0.2	28.9	-	-
	CAN99-16	Athabasca (Falls)	+ 3.2	0.1	19.4	-	-
	CAN99-21	Fraser	+ 1.4	0.1	48.3	10387	92276
	CAN99-44	French	+ 1.7	0.2	17.9	4377	46059
	CAN99-46	Liard (UpperLiard)	+ 0.9	0.2	35.1	8235	65224
	CAN99-51	Hyland	+ 0.3	0.1	44.3	8458	71206
	CAN99-53	Coal	+ 0.3	0.1	46.2	7939	70306
	CAN99-57	Trout	+ 2.8	0.2	17.8	3042	32082
Interior Platform	CAN96-14	Little Smoky	- 0.5	0.1	37.8	5713	79888
	CAN99-72	Beatton	- 1.7	0.2	51.4	2968	77294
	CAN99-73	Halfway	+ 0.6	0.1	30.5	2968	50029
Transition mountain/plain	CAN96-16	Smoky	- 0.4	0.1	38.9	5565	76024
	CAN99-78	Smoky	- 0.1	0.1	35.3	4674	64376
	CAN96-23	Fort Nelson	- 1.2	0.2	56.1	3190	74965
W. Cordillera	CAN99-30	Nass	+ 1.0	0.2	37.6	14690	73376
	CAN99-39	Stikine	+ 1.2	0.1	24.5	15358	75071

853	Table 3
854	
855	
856	
857	
858	
859	

Tributaries	Sample type sample number		sample name/location	δ ⁷ Li	$2\sigma_{m}$	Li µg/g
Main tributaries	Sand	CAN96-6	Mackenzie @Tsiigehtchic	- 0.5	0.1	47.4
Rocky and Mackenzie Mountains	Black Shale	CAN99-5	Athabasca @Hinton	- 1.0	0.2	29.1
	Carbonate	CAN99-65	Mc Donald	+ 7.2	0.1	1.5
Transition mountain/plain	Black Shale	CAN99-70	Sikanni Chief	- 1.1	0.1	64.4
Canadian Shield	Glacial Till	CAN96-31	Yellowknife	+ 4.8	0.2	16.2
	Glacial Till		Prosperous Lake	+ 5.2	0.2	54.6