

# Buried fluvial incisions as a record of Middle–Late Miocene eustasy fall on the Armorican Shelf (Bay of Biscay, France)

Fabien Paquet, David D. Menier, Guilhem Estournès, Jean-François Bourillet, Pascal P. Le Roy, François Guillocheau

# ► To cite this version:

Fabien Paquet, David D. Menier, Guilhem Estournès, Jean-François Bourillet, Pascal P. Le Roy, et al.. Buried fluvial incisions as a record of Middle–Late Miocene eustasy fall on the Armorican Shelf (Bay of Biscay, France). Marine Geology, 2010, 268 (1-4), pp.137-151. 10.1016/j.margeo.2009.11.002 . hal-00563648

# HAL Id: hal-00563648 https://brgm.hal.science/hal-00563648

Submitted on 26 Mar 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#### Marine Geology January 2010, Volume 268, Issues 1-4, Pages 137-151 http://dx.doi.org/10.1016/j.margeo.2009.11.002 © 2009 Elsevier B.V. All rights reserved.

## Buried fluvial incisions as a record of Middle–Late Miocene eustasy fall on the Armorican Shelf (Bay of Biscay, France)

Fabien Paquet<sup>a, e, \*,</sup> David Menier<sup>a</sup>, Guilhem Estournès<sup>a</sup>, Jean-François Bourillet<sup>b</sup>, Pascal Leroy<sup>c</sup> and François Guillocheau<sup>d</sup>

<sup>a</sup> Université de Bretagne-Sud, EA 2219 GeoArchitecture, Campus de Tohannic — ENSIbs, rue Yves Mainguy, 56017 Vannes, France

<sup>b</sup> IFREMER, Centre de Brest, 29280 Plouzané, France

<sup>c</sup> IUEM, Technopôle Brest-Plouzané, 29280 Plouzané, France

<sup>d</sup> Université de Rennes 1, UMR-CNRS 6118 Géosciences Rennes, Campus de Beaulieu, CS 74205, F-35042 Rennes Cedex, France

<sup>e</sup> BRGM, Geology Division — Geology of Sedimentary Basins, 3 rue Guillemin, BP 36009, 45060 Orléans Cedex 2, France

\*: Corresponding author : Fabien Paquet, Tel.: + 33 2 38 64 47 51; fax: + 33 2 38 64 33 33, email address : f.paquet@brgm.fr

#### Abstract:

High-resolution seismic data have been acquired in June 2008 on the Armorican Shelf (AS), in the northern Bay of Biscay, in order to reassess its stratigraphic architecture in detail and to study the impact of eustasy, tectonic and sediment delivery on the margin sedimentary record. Several profiles show fluvial-type incisions of several tens of meters (up to 54 m) associated to a widespread erosion surface. Several hypotheses are proposed for the stratigraphic position of this surface and incisions. We suggest Middle to Late Miocene age. Considering the relatively quiescent tectonic activity of the margin, we infer that the relative sea-level fall responsible of aerial incision on the Miocene shelf is eustasy-related. We propose the attested Serravallian–Tortonian eustatic lowstand (c. 11.6 Ma) as the key event responsible of such erosion and incision. This event marks the early beginning of the high-amplitude sea-level fluctuations that culminated during the Pleistocene and significantly controlled the present day AS morphology. The variability of vertical incision observed along single reaches can be explained by the confluence of several tributaries, the sinuosity of the channel and can be amplified by the unconsolidated nature of the Miocene substratum. The main pathways of the fluvial network corresponding to these buried valleys have been reconstructed and connections to other existing networks are proposed.

Keywords: Armorican Shelf; incised valleys; Miocene; eustasy; passive margin

#### 1. Introduction

In low sedimentation rate and quiescent passive margins, eustasy has a predominant effect on the stratigraphic architecture ([Vail et al., 1977] and [Posamentier and Vail, 1988]). In such settings, the development of fluvial channel incisions over the shelf is considered as an indicator of a significant global sea-level fall and lowstand, that expose the entire shelf below or close to the breakpoint (Dalrymple et al., 1994) and is associated sometimes to submarine canyon incisions (Vail et al., 1991). Several examples are documented worldwide and are used to date or constrain the amplitude and effects of eustasy changes through time ([Fulthorpe et al., 1999], [Fulthorpe et al., 2000] and [Lofi and Berné, 2008]) or are the subject of ongoing studies (e.g. IODP expeditions 313 and 317). In the northern Bay of Biscay, numerous Quaternary incised valleys described over

46 the Armorican Shelf illustrate the impact of the high-amplitude Pleistocene glacio-47 eustasy falls (c. -120m) (Pinot, 1974; Menier et al., 2006; Chaumillon et al., 2008). 48 Important sea-level falls of several tens of meters are also suspected for the Neogene 49 period (Hag et al., 1987) and may have resulted in the generation of a fluvial networks 50 over the emerged AS. Despite the presence of widespread erosion surfaces, related 51 Neogene channel incisions are not attested and clearly described directly offshore 52 southern Brittany. This is mainly due to the scarcity and the average quality of data 53 (Bourillet et al., 2005). However, the –Quaternary" network drawn by Pinot (1974) 54 was partly based on the detection of incisions on seismic data, c. 100 km from the 55 present day coastline. Surprisingly, the Pleistocene valley network as described in 56 recent studies (Menier, 2004; Menier et al., 2006; Thinon et al., 2008) does not extend 57 that far over the outer shelf. A re-evaluation of the stratigraphic position of the outer-58 shelf incisions of Pinot (1974) was therefore needed to verify a pre-Quaternary age 59 attribution. Evidence of potential Neogene channel incisions in surrounding areas is 60 located in the Western Approaches and are associated with the Fleuve Manche' 61 paleoriver system (Reynaud et al., 1999; Bourillet et al., 2006) where Neogene series 62 are well-documented (Evans and Hughes, 1984), and to the South, over both the 63 Vendean-Armorican Platform (Huerta et al., in press) and the North Aquitaine shelf 64 (Bellec et al., 2009).

This study presents the interpretation of recent high-resolution seismic sparker
profiles acquired on the Armorican Shelf that show clear incisions within the Miocene
series. We discuss the origin and the age of these incised valleys, and describe their
morphology and potential significance in terms of eustasy.

69

#### 70 GEOLOGICAL SETTING

71 The Armorican Shelf (AS) corresponds to a segment of the European Atlantic 72 margin in the northern Bay of Biscay (Fig. 1). Its formation initiated during the Early 73 Cretaceous (Barremian-Aptian, c. 125 Ma) consequently to the rotation of the Iberian 74 Peninsula and the contemporaneous opening of the Bay of Biscay (Montardet et al., 75 1979; Olivet, 1996; Thinon, 1999). Its present day morphology corresponds to a large 76 polygenetic erosion surface (wave-planed) resulting from significant Plio-Pleistocene 77 sea-level fluctuations (Imbrie et al., 1984; Waelbroeck et al., 2002). The Cenozoic 78 was dominated by post-rift mixed carbonaceous-silicoclastic sedimentation as 79 revealed by exploration well PENMA-1 (Preux, 1978), and cores (Bouysse et al., 80 1974; Thomas, 1999; Guillocheau et al., 2003). The knowledge of the stratigraphic 81 architecture of the AS is provided by interpretation of seismic data from the Bir 82 Hakeim survey (Bouysse et al., 1968) and correlations with PENMA-1 well 83 (Guillocheau et al., 2003). Post-rift deformation occurred mostly in response to the 84 Pyrenean collision that reactivated strike-slip Variscan structures (Montardet et al., 85 1979). This subsequent Oligocene compressive phase did not affect obviously this 86 area such as it did in the English Channel, with the modification of the Hurd deep 87 (Lericolais et al., 2003). The limited Cenozoic tectonic activity in the study area is 88 evidenced by sub-vertical faults that suggest a dextral strike-slip movement that faded 89 during the Miocene (Fig. 4 and 5). The margin is also affected by an apparent seaward 90 regional tilt documented onshore (Bonnet et al., 2000) and offshore (Vanney et al., 91 1972; Bourillet et al., 2003). Several regional erosion surfaces resulting from sea-level 92 changes and/or phases of regional tilt are recorded and form angular unconformities 93 (Fig. 1C, 4 and 5). One of these surfaces is interpreted to be middle to late Miocene in 94 age (Vanney et al., 1972; Guillocheau et al., 2003, Bourillet et al., 2003). The

95 Miocene-recent deposits of the AS margin are mostly located on the median and outer 96 parts of the shelf where they form a stack of wedges (Fig. 1C and 5). Incised valleys 97 are present over the inner shelf and the coastal area (Boillot et al., 1971; Bouysse et 98 al., 1974; Menier, 2003; Menier et al., 2006; Chaumillon et al., 2008). The origin of 99 both these incisions and their valley fills is linked to Late Pleistocene sea-level 100 fluctuations. Unfortunately, due to coarse superficial sediments, no attempts at coring 101 have yet succeeded in recovering paleovalley fill (Bourillet and Turon, 2003). 102 Recently acquired high-resolution seismic data detail the geometry of the 103 sedimentary succession and associated erosion surfaces. One of these surfaces 104 revealed several incisions, c. 100 km from the present day coastline.

105

#### 106 **METHODOLOGY:**

107 This study is based on the interpretation of high-resolution Sparker seismic data 108 acquired onboard CNRS-INSU -Côtes de la Manche" vessel in June 2008 (GeoEtel-109 2008 survey,) as well as the re-interpretation of lower-resolution seismic profiles from 110 older surveys (Bir Hakeim 1967, Job-ha-Zelian 1971, 1973 and 1974). Seismic units 111 and bounding surfaces are defined by describing reflection continuity, amplitude, 112 frequency, configuration and terminations (Mitchum et al., 1977a and 1977b). The 113 seismic stratigraphy obtained from our interpretations have been geometrically 114 correlated to cores and dredges samples and to PENMA-1 exploration well, which 115 provides the only regional access to the Meso-Cenozoic sedimentary record (Fig. 5). These correlations have been made assuming a mean velocity of  $2250 \pm 250 \text{ m.s}^{-1}$  for 116 the Miocene-recent interval and  $2910 \pm 10 \text{ m.s}^{-1}$  for the Oligocene one (velocity from 117 118 Garlizenn-1 exploration well; Maillard, 1982).

#### 120 **RESULTS:**

#### 121 Seismic stratigraphy

122 Sparker seismic profiles image the sedimentary succession from Paleogene to Present 123 from the median shelf to the shelf break. Six seismic units and one sub- unit have 124 been identified (U1 to U6 and U6b) separated by six bounding surfaces (S1 to S6). 125 Seismic characteristics of units and surfaces are described in Table 1. Each unit is 126 composed of one seismic facies that does not vary significantly laterally. In the 127 following section, we describe the seismic stratigraphy in terms of geometry and 128 stacking pattern. We also propose both lithology and stratigraphic position for units 129 when they are documented by subsurface samples, by correlations to surfaces at 130 PENMA-1 well (Preux, 1978), and on the basis of former studies (e.g. Vanney et al., 131 1972; Guillocheau et al., 2003).

132 Seismic unit 1 (U1):

U1 is the lower most unit identified on seismic data. As a result, no basal boundary is
visible on seismic data. Its seismic facies shows chaotic and locally steep reflections
(Table 1). It crops out along the present day coastline where it as been sampled (Fig.
5). It consists of metamorphic (micaschists) and crystalline (granites) Paleozoic rocks
deformed during the Variscan orogeny (Audren and Lefort, 1977). It therefore forms
the basement of the AS. The upper boundary of U1 is a clear erosion surface (S1) and
represents the base of the sedimentary succession of the AS.

140 Seismic unit 2 (U2):

141 U2 lies directly on the Variscan basement where it presents few onlap terminations on

- 142 S1. Its thickness is unclear but probably exceeds 400 ms (Fig. 5). U2 is made up of
- 143 parallel and continuous reflections (Table 1) characteristic of well-bedded marine
- sediments. U2 is sampled in PENMA-1 and at several locations over the inner shelf
- 145 where it corresponds to Paleogene chalks, calcareous mudstones and sandstones with
- rare terrigeneous-rich beds (Andreieff et al., 1968; Barbaroux et al., 1971; Bouysse et
- 147 al., 1974; Delanoë and Pinot, 1974; Delanoë et al., 1975; Preux, 1978). U2 is
- bounded above by erosion surface S2 highlighted by toplap terminations (Fig. 4 and
- 149 5) that reflects a slight regional tilt ( $< 0.5^{\circ}$ ). U2 is affected by brittle deformation with
- 150 NW trending sub-vertical faults and gentle folds (Fig. 1, 4 and 5).
- 151 The presence of the thick Cretaceous section (c. 500 m) identified in the lower third of
- 152 PENMA-1 (Fig. 3) is not attested at the seafloor, over the inner shelf, between
- 153 basement and U2. This may be due to a landward thinning and/or onlap of the series
- as visible in petroleum conventional seismic data (unpublished petroleum data).
- 155 Seismic unit 3 (U3):
- 156 U3 is a 40-50 ms-thick unit characterized by discontinuous and slightly wavy very
- 157 high-amplitude reflections (Table 1). It is bounded below by unconformity S2 and
- above by S3 that shows rather little erosion except on line 20-670 (Fig. 5). No
- seafloor samples have been collected in the area where the unit may crops out.
- 160 Nevertheless, U3 correlates geometrically to the upper-part of the Oligocene series
- 161 (Chattian?) at PENMA-1 (Fig. 5) where it is made up of shallow marine coral-rich
- 162 bioclastic sandstones (Fig. 3). Discontinuity and wavy forms of reflections within U3
- 163 may correspond to small coral reefs or mounds. The mapping extent of U3 is unclear
- 164 but should consist in a narrow corridor seaward from U2. Tectonic deformation style
  - 7

165 affecting U3 is similar to U2 but in an evident lesser proportion (Fig. 4 and 5). This 166 observation and the tilting of U2 suggest that a deformation phase occurred during 167 Paleogene with a probable culmination between Late Eocene and Late Oligocene. 168 Such a phase is described by several regional studies and syntheses and corresponds 169 to the Pyrenean compressive tectonic phase s.l. (Gély and Sztràkos, 2001). It is 170 supposedly responsible for 1) the inversion in the English Channel documented 171 between the Mid-Late Eocene and Early-Oligocene (Ziegler, 1987; Evans, 1990), 2) 172 the sedimentary hiatus in the Bay of Biscay (Hailwood et al., 1979), and coincides 173 with the contemporaneous uplift of North Atlantic margins (Anell et al., 2009 and 174 references herein). This phase may have partly triggered the development of S2 175 unconformity.

176 Seismic unit 4 (U4):

177 U4 is a 100 to 150 ms-thick unit made up of sub-parallel, continuous and low-178 amplitude reflections (Table 1). It shows downlap terminations on its lower boundary 179 S3 and an overall progradation pattern interrupted by transgressive phases of a lesser 180 order evidenced by onlap surfaces (Fig. 4 and 5). The seismic facies suggest well-181 bedded fine grained deposits (mudstones, siltstones). U4 is topped by a widespread 182 erosion unconformity that is characterized by toplap terminations of the underlying 183 reflections (Fig. 4 and 5). This erosion surface is more or less penetrative into the 184 underlying sediments and locally shows clear deep incisions on several profiles (Fig. 185 4, 5 and 6). Very few seafloor samples have been collected in the area (Fig. 1) and 186 mainly consist in Miocene calcareous marls and Aquitanian bioclastic limestones 187 (Andreieff et al., 1968). At PENMA-1, S3 and the lower part of U4 respectively 188 correlate with a thin poorly-sorted terrigeneous sandstone bed and a Miocene mixed 189 marine terrigeneous-calcareous mudstone. This unit may find lateral equivalents in

the Miocene -faluns" (shallow marine mixed bioclastic and terrigeneous coarse sands)
that are well documented onshore (Brittany, Anjou, and Tourraine areas; Lécuyer et
al., 1996; Néraudeau, 2003)

193 These observations suggest that S3 developed at the transition between a documented 194 Oligocene sea level low and a Miocene sea level high (Hag et al., 1987) evidenced by 195 the change from shallow marine limestones with corals to transgressive glauconitic 196 sandstone then prograding marine mudstones (Preux, 1978). The upper part of U4 is 197 not sampled or described in PENMA-1 (no recovery), it is thus lacking reliable 198 lithology and age calibrations (see discussion). The upper limit S4 presents evident 199 erosion features including irregular toplap/truncation surface with channel-like 200 incisions (Fig. 4, 5 and 6). As for S2, the toplap terminations evidenced a slight 201 seaward tilt ( $< 0.5^{\circ}$ ) of the sedimentary section (Fig. 5). The age calibration of S4 can 202 not be directly determined from existing samples and different hypotheses are 203 addressed in the discussion section.

### 204 Buried channel incisions of S4

205 Among several unconformities identified in the seismic stratigraphy of the AS, S4 is

the oldest to exhibit a deeply incised surface. Careful study of available seismic

207 profiles allows a better understanding of the processes resulting in the generation of

208 S4 and associated erosion features (Fig. 4 and 6). Incisions present (1) major and

- 209 deeper channels with (2) several associated secondary channels (tributaries), separated
- by (3) strath terraces developing in (4) a wider valley. These observations are in
- agreement with a fluvial and thus subaerial origin for the incisions (Posamentier,
- 212 2001). A possible karstic origin for these features is neglected here because the
- 213 incised lithologies contain a terrigeneous fraction (PENMA-1, Preux, 1978). The

| 214 | incision depth of the major channel and the valley shape along transversal profiles                             |
|-----|---|
| 215 | vary dramatically (Fig. 6). On profiles 7 and 26 (Fig. 6a and 6c), the vertical incision,                       |
| 216 | from the valley edges to the channel bottom, reaches 30 ms and 60 ms respectively                               |
| 217 | $(30 \text{ m and } 60 \text{ m at } 2000 \text{ m.s}^{-1})$ and is strongly localized, whereas, in between, on |
| 218 | profile 22, incision is less deep (10 ms; 10 m at 2000 m.s <sup>-1</sup> ) (Fig. 6b). Likewise, the             |
| 219 | shape of channels on seismic profile shows a clear variability from V-shape (e.g.                               |
| 220 | Profile 26, Fig. 6a) to U-shape (e.g. Profile 7, Fig. 6c) with more or less symmetry.                           |
| 221 | This is in agreement with seismic profiles crossing sinuous valley reaches with                                 |
| 222 | variable azimuth angles (Fulthorpe et al., 1999). The density of seismic profiles                               |
| 223 | allows the mapping of a paleo-fluvial network in that area (Fig. 7 and 8). We                                   |
| 224 | tentatively trace four distinct valleys, labeled A, B, C and D, from west to east, by                           |
| 225 | correlating incisions between profiles (Fig. 8). Valleys strike to the NE-SW,                                   |
| 226 | orthogonally to the margin. As incision depth fades seaward and as S4 and associated                            |
| 227 | valleys are eroded landward by S6, the network can not be traced over more than 30-                             |
| 228 | 40 km (Fig. 8). Our interpretations suggest a connection of both valleys B and C                                |
| 229 | downstream. The valley fill deposits (base of U5) are made up of several sub-units                              |
| 230 | separated by U-shaped erosion surfaces that laterally incise and widen the preexisting                          |
| 231 | main channel (e.g. profile 26, Fig. 6a). On profile 7, younger channelized incisions                            |
| 232 | are visible and one of them truncates the main erosion surface S4 (Fig. 6c). Therefore,                         |
| 233 | generation and reshaping of S4 erosion surface ends after deposition of U5 starts.                              |
| 234 | Seismic unit 5 (U5):  |
| 235 | U5 is a 40-50 ms-thick lens-shaped unit that thins both landward and seaward (Fig. 5).                          |

236 It shows very chaotic and discontinuous reflections with channels and onlap

237 terminations on S4 (Table 1). Such seismic facies suggests sandy sediments. Deposits

238 in the basal part of U5 fill the S4 incisions and present U-shaped erosion surfaces that

239 locally incise U4. Above, the upper-boundary S5 shows unusual characteristics with 240 possible channel incisions (Fig. 4 and 5) and a seaward truncation of several U5 241 reflections (Fig. 5). The age of U5 is not defined as but intersection between S5 and 242 sea floor corresponds to the base of the Pliocene-Pleistocene sedimentary succession. 243 Seismic Unit 6 (U6): 244 U6 is a >200 ms-thick unit that corresponds to the seaward most seismic unit 245 individualized in this study. It lies conformably over S5 and reflections are 246 discontinuous, chaotic with possible channels (Table 1) and show a seaward dip of c. 247 5° (Fig. 5). The seismic facies is similar to U5 suggesting sandy material. The upper 248 boundary of U6 corresponds to the -flat" sea floor that forms the morphology of the 249 shelf. U6, as well as all the other seismic units (U1 to U5) is strongly eroded at the sea 250 floor. On GeoEtel-2008 seismic data (eg. Fig. 4, 5 and 6), this erosion surface is 251 resolved as 1) a proper erosion surface topped by 2) a thin and discontinuous sediment 252 cover (sub-unit U6b). This sediment cover is made up of unconsolidated mixed 253 terrigeneous and bioclastic clayey sands of Holocene age. The age calibration of U6 254 can not be attested with certainty as no sea floor sample exists in the area except for 255 U6b. Nevertheless, U6 being the last and thus youngest unit of the AS we propose a 256 Pliocene-Pleistocene age. This is also supported by the complex internal stacking 257 pattern with several unconformities that may illustrate the impact of the Plio-258 Pleistocene eustasy variations. S6 may therefore correspond to the last member of a 259 series of coincident erosion surfaces resulting from the dramatic high-amplitude 260 Pleistocene sea level variations.

261

262 Summary:

263 Our seismic interpretation allows the identification of six seismic units separated by 264 major unconformities. The first unit U1 corresponds to the Paleozoic basement and 265 the following ones represents the Cenozoic sedimentary cover of the Armorican Shelf. 266 Lithology, calibrated from core samples and PENMA-1 well progressively evolves 267 from a biogenic carbonate-dominated sedimentation to mixed bioclastic-terrigeneous 268 deposits throughout the whole sediment section. Several unconformities show erosion 269 features and both S2 and S4 record phases of Cenozoic seaward tilting of the margin. 270 S4 is the oldest erosion surface to exhibit deeply incised fluvial valleys. It seems to 271 mark an important step in the transition between the two sedimentation styles stated 272 above. In the following section we discuss the age calibration of S4, the deciphering 273 of the various controlling parameters that lead to its generation, the morphology of the 274 valley reaches, and its significance in terms of paleogeography and evolution of the 275 shelf morphology.

276

#### 277 **DISCUSSION:**

278 Existence of fluvial channels on the AS continental margin indicates that rivers were 279 once flowing over the emerged shelf but the stratigraphic position of the incised 280 erosion surface S4 within sedimentary succession is still unclear. These channel-like 281 incisions had already been described by Pinot (1974) and associated to a continuous 282 late Pliocene-Pleistocene fluvial network extending over the AS and connecting 283 onshore valleys to the slope canyons. Nevertheless, careful study at the newly 284 acquired seismic data reveals that 1) the channels are associated with the tilted 285 Neogene S4 erosion surface and that 2) no direct connections to the inner shelf 286 paleovalleys and/or slope canyons can be traced.

#### 287 Age calibration of S4:

288

Seismic units (U4 and U5) on both side of S4 have not been properly dated making

age calibration of this surface rather speculative. Our interpretation and correlations

- to PENMA-1 indicate that U3 and the base of U4 are of Late Oligocene and Early
- 291 Miocene age respectively, and that a Pliocene-Pleistocene age is confidently
- attributed to U6. Therefore, the age of S4 is comprised between Early Miocene and at
- some stage in Pliocene. By analogy to the documented stratigraphy of the nearby
- 294 Western Approaches (Fig. 9) where incised valleys are also identified (Evans and
- Hughes, 1984; Powell, 1988; Reynaud et al., 1999, Bourillet et al., 2003; Stewart and
- 296 Davies, 2007), two hypotheses can be formulated as follow. These hypotheses are

then tested using the event timing of regional and global controlling parameters

- 298 (tectonic deformation, eustasy, and climate) that have been recorded in Western
- Europe and in other remote margins.

#### 300 *Hypothesis 1 – "Late Miocene-Pliocene" age model:*

301 In this model, 1) U4 is the lateral equivalent of both Jones and Cockburn formations

- 302 that are of early to mid-Miocene (Aquitanian-Serravallian) and of mid- to late
- 303 Miocene age respectively (Evans and Hughes, 1984; Powell, 1988), and 2) U5 and
- 304 U6-U6b correspond to both the Plio-Pleistocene Little Sole formation and the
- 305 Pleistocene-Holocene Melville formation respectively. In such context, S4 and
- 306 associated valleys are likely of late Miocene Pliocene age. This is in agreement with
- both the Messinian age preferred by Bourillet et al. (2003) and the late Pliocene age
- 308 proposed by Reynaud et al. (1999) for the lateral equivalent of S4 in the Western
- 309 Approaches.

#### 310 *Hypothesis 2 – "Mid-Late Miocene" age model:*

| 311 | In this model, U4 has only the Jones formation as lateral equivalent whereas the      |
|-----|---|
| 312 | Cockburn formation correlates with U5, and both U6 and U6b are equivalent to the      |
| 313 | Little Sole and the Melville formations. Thus S4 developed earlier, close to the mid- |
| 314 | late Miocene boundary. Such an attribution is also consistent with the study of       |
| 315 | Bourillet et al. (2003) that indeed proposed an age ranging from c. 12 Ma             |
| 316 | (Serravallian-Tortonian) to c. 5 Ma (Messinian). Late-Pliocene incised valleys        |
| 317 | (Reynaud et al., 1999) would find more suitable equivalents within U6 where           |
| 318 | channels are described.   |
|     |   |

319 Without reliable ages, discussing S4 age models requires to determine the possible

320 control parameters that lead to S4 development and the related timing of events.

321 Fluvial incisions into marine deposits evidence the emergence of the shelf that implies

an important relative sea-level (base-level) fall. Such a fall find its origin in either (1)

323 a tectonic deformation (local uplift) and/or (2) an eustatic fall (global).

#### 324 Cenozoic tectonics:

325 Tectonic deformation within the sedimentary succession of the AS and identified on

326 seismic data is characterized by sub-vertical faults and associated folds that affect

327 mostly the seismic unit U2 (Paleocene-Eocene). Angular unconformity S2 seals most

328 of the brittle deformation and only few faults traverse the following seismic units

329 (Fig. 4 and 5). S4 corresponds to another angular unconformity that shares several

features with S2, including toplap terminations of U4 reflections.

331 Over the Bay of Biscay area and over the AS, the Cenozoic tectonic deformation is

attributed to the alpine orogeny s.l. (Fig. 9). Two major phases are recognized in the

- 333 literature (Ziegler, 1990; Bourrouilh et al., 1995). The Pyrenean phase *s.l.* extends
- from Eocene to mid-Oligocene (Gély and Sztràkos, 2000 and 2001). It is expressed on

335 the AS as the series of sub-vertical faults (reactivated Variscan structures) and folds 336 visible in the contemporaneous seismic unit U2 (Bouysse et Horn, 1971; Delanoë, 337 1988), and as the Oligocene angular unconformity S2 that mostly seals this 338 deformation (Fig. 4 and 5). Contemporaneously with the late Pyrenean phase is the 339 first basin inversion of the Western Approaches Trough (Ziegler, 1987). The second 340 phase of deformation is related to the Miocene Alpine contractional phase. Its record 341 on the AS may correspond to the play of part of the Paleogene sub-vertical faults that 342 affect seismic units from U3 to U6 (Oligocene to recent). Similarly to S2, erosion 343 surface S4 is an angular unconformity that also highlights and seals a Neogene tilt of 344 the AS toward the SW (Fig. 4 and 5). This phase is also contemporaneous with a new 345 phase of inversion in the Western Approaches Trough, and both are of mid-late 346 Miocene to Pliocene age (Ziegler, 1987, 1990, Guillocheau et al., 2000). Thus, 347 Neogene tectonic deformation phases provide a good framework to explain the 348 various features identified on seismic data. Unfortunately, the duration of these phases 349 and the poor age control on their pulses preclude any accurate age calibration of S4. 350 Alternatively, northwest European passive margins undergone both tilting and 351 sagging attributed to mantle-driven processes during Cenozoic (Praeg et al., 2005) but 352 the timing and signature of deformation differs from what is recorded along the AS. 353 Finally, looking at the present day architecture of the AS margin (Fig. 4), the angular 354 erosion unconformity S6 attests a Plio-Pleistocene tilt of the margin that has increased 355 the subsidence of the outer shelf and the burial and preservation of S4 incised valleys.

356 Cenozoic eustasy:

357 During Cenozoic, sea level has undergone important and abrupt changes (Fig. 9)

accompanying the overall transition from the Mesozoic early-Cenozoic greenhouse to

the Quaternary icehouse conditions (Haq et al., 1987; 1988; Miller et al., 1998; Miller

360 et al., 2005). One example is the postulated mid-Oligocene eustatic lowstand visible 361 on the eustatic curve of Hag et al. (1987; 1988) close to the Rupelian-Chattian 362 boundary, or alternatively attributed to an earlier Oligocene cooling event by Miller et 363 al. (2008). According to different studies the estimated amplitude of the fall varies 364 from -160 m (Haq et al., 1987) to -55 m (Miller et al., 2005). On the AS, this 365 documented early- or mid-Oligocene eustatic fall correlates fairly well with the S2 366 erosion unconformity and is also illustrated by the transition from the outer-shelf 367 Eocene marine deposits from shallower inner-shelf Oligocene deposits described at 368 PENMA-1 (Preux, 1978; Guillocheau et al., 2003) (Fig. 3). Thus, the generation of S2 369 unconformity and its characteristics would result from the combined effects of both 370 tectonic deformation and a eustatic sea level fall.

371 Considering the similarity between S2 and S4 unconformities (Fig. 5), and especially

the existence of incised fluvial valleys (Fig. 4, 5 and 6), the generation of S4 would

373 have required a relative sea level fall following the Oligocene fall event with

amplitude similar or greater. One other major climatically-induced sea level fall

375 documented worldwide for this period starts during the late Langhian - early

376 Serravallian (c. 14 Ma) and culminates around the Serravallian-Tortonian boundary

377 (mid-late Miocene boundary at 11.608 Ma – GSSP; Hilgen et al., 2005) (Haq et al.,

378 1987, 1988; Kominz et al., 1998; Miller et al., 1998; John et al., 2004; Miller et al.,

379 2005). The Serravallian-Tortonian boundary correlates to the glacio-eustatic lowstand

380 T3.1 (Haq et al., 1987, 1988) associated with the permanent establishment of the East

- 381 Antarctic Ice Sheet (Zachos et al., 2001). The amplitude of the Middle Miocene sea-
- level fall estimated in former studies is still debated with values ranging from -25 m
- 383 to -180 m (Haq et al., 1987; Pigram et al., 1992; Miller et al., 1998; John et al., 2004).
- 384 We propose that S4 and associated channel incisions were completed close to the

385 Serravallian-Tortonian boundary at c. 11.6 Ma with a probable initiation occurring 386 earlier during late-Langhian early-Serravallian. The onset of the Icehouse and long-387 term sea-level lowstand conditions would also explain the presence of several 388 younger (e.g. Tortonian and Messinian) channelized erosion surfaces identified within 389 U5 and U6 (Fig. 4 and 5). In addition, Mio-Pliocene fluvial network remnants have 390 been described onshore in Brittany (Guillocheau et al., 1998; Van Vliet-Lanoë et al., 391 1998) in several places including Lauzach and Réguigny (Fig. 8). The basal fluvial 392 deposits filling the oldest incised valleys have been dated by ESR at  $8.7 \pm 1.5$  Ma 393 within the Tortonian (Van Vliet-Lanoë et al., 1998; Brault et al., 2004). This suggests 394 a possible Serravallian or early-Tortonian age for the incision phase. The 395 reconstruction of the Tortonian network provided by Brault et al. (2004) and based on 396 paleocurrent directions shows a southwestward direction for both the onshore 397 Tortonian and the supposedly contemporaneous offshore S4 fluvial networks (Fig. 7). 398 In addition, onshore outcrops exhibit several erosion surfaces and associated fluvial 399 networks (Brault et al., 2004). This configuration is shared by the offshore buried 400 valleys that show several younger incisions (Fig. 6). Thus, both offshore and onshore 401 networks could be genetically linked. Unfortunately, due to the joint effects of (1) the 402 Neogene seaward regional tilting of the AS and (2) the wave abrasion that occurred 403 during the Plio-Pleistocene sea-level fluctuations and gave its shape to the present day 404 AS (polygenetic wave-planed surface), the channelized erosion surface is only 405 preserved 100 km offshore the present day coastline. Such configuration avoids direct 406 correlations between this distal buried fluvial network and the onshore one. In the 407 same way, the question could be asked about the age of the so-called Quaternary 408 inner-shelf incised valleys. If a Plio-Pleistocene age for the sedimentary fill sounds

409 reasonable (Menier et al., 2006; Proust et al., 2009), the incision phase could be

410 underestimated (Menier, 2004) and be reassessed to the mid-Miocene time.

411 Compared to surrounding areas, a Serravallian-Tortonian boundary age attribution for 412 S4 seems reasonable. Over the North Aquitaine Shelf, Bellec et al. (2009) describe 413 several unconformities showing incised valleys. They propose that these surfaces 414 formed during the successive Langhian to early-Tortonian sea level falls, the last one 415 being the most erosive and correlating T3.1 sea level fall of Haq et al. (1987) as 416 proposed for S4 on the AS. The absence of earlier incisions below S4, within U4, may 417 be explained by (1) lower subsidence and sedimentation rates of the AS area that 418 resulted in the stacking of all surfaces in only one (S4) or (2) because incisions 419 occurred landward and have been eroded later (Fig. 5). Over the Western Approaches, 420 the age of the incised valleys is still debated but Bourillet et al. (2003) propose an age 421 ranging from 12 to 5 Ma (Serravallian to Messinian) based on a range of subsidence 422 rates. Such an age range is compatible with our age estimation for S4. Likewise, 423 major erosion unconformities attributed to mid- to late Miocene sea level falls are 424 described in other remote passive margins such as the New Jersey shelf (Fulthorpe et 425 al., 1999), the northern Java shelf (Posamentier, 2001), and the Gulf of Lions (Besson 426 et al., 2005). The sea level fall at the Serravallian-Tortonian boundary is specifically 427 preferred as the triggering event for several of these surfaces such as in the Gulf of 428 Papua (Tcherepanov et al., 2008) or in the Gulf of Lions (Lofi and Berné, 2008). 429 We thus favor the second hypothesis, and consequently attribute a Tortonian age to 430 U5 (equivalent of the WA Cockburn formation and the first onshore -Red Sands" 431 (Fig. 9). Erosion surface S5 could be of Messinian or basal Pliocene age (Hag events 432 TB3.4 or 3.6). U6 would therefore correspond to the Pliocene-Pleistocene outer shelf sedimentary prism and is a lateral equivalent of the Little Sole formation. 433

The Serravallian eustatic fall would have thus progressively exposed the shelf of the AS, thus creating the major unconformity S4 and associated channels. This event took place while the whole AS undergone a regional tilt related to the Alpine orogenic phase. It results that S4 developed also as an angular unconformity. This tilt has induced a local uplift of the area thus increasing the effect of eustasy on the relative sea level fall, and enhancing the stratigraphic signature S4.

#### 440 Significance of the valley morphology

441 At its lowest, the Serravallian – Tortonian sea level may have reached depth close to

442 or even below the smooth and rounded clinoform breakpoint. Potential connection

between S4 incised valleys and hypothetical mid-Miocene slope canyons, is not yet

444 demonstrated. This is mainly due to the scarcity of data and their rather low

445 penetration where the mid-late-Miocene strata are buried below the thick Plio-

446 Pleistocene wedge. Nevertheless, when looking at the vertical incision depth of the

447 four valleys, it tends to decrease south-westward (seaward) indicating that valleys and

448 canyons are not directly connected.

449 Alternatively, depth of vertical fluvial incision on shelves is often considered as a

450 proxy for estimating the depth reached during the sea-level lowstand. Concerning the

451 valleys described in this study, the use of the depth of channels, to estimate the

452 contemporaneous sea-level appears problematic. The observed variability of depth

453 incision of one valley between seismic sections corresponds to large-scale undulations

454 of the river bottom longitudinal profile and suggests a riffle-pool sequence

455 morphology developing on a sinuous channel (Richards, 1982; Lofthouse and Robert,

456 2008). Testing this hypothesis would require additional data in order to constrain the

457 three-dimensional morphology of the channels, the length of the riffle-pool sequence,

458 and finally, to discuss the potential river characteristics (e.g. water discharge, 459 sediment load, grain-size). The -pool" areas may also be controlled by the confluence 460 of one or more tributaries with the main channel. In addition, we consider that the 461 young and unconsolidated lower to middle Miocene series, incised by the Mid-462 Miocene rivers, has played a significant role by facilitating the incision in the -pools" 463 areas. Such variability of the depth of incision thus precludes accurate and reliable 464 estimation of the depth reached by the sea-level during the late Serravallian – early 465 Tortonian lowstand. Moreover, the tectonic tilt of the margin and the associated uplift 466 of the area, evidenced by the angular unconformity S4, have also played a role in the 467 relative amplitude of the sea-level fall. Additional data are needed to better 468 understand the valley morphology and its relation with the shelf edge.

469

#### 470 CONLUSIONS

471 High-resolution seismic data recently acquired on the Armorican Shelf reveals the 472 presence of buried channel incisions developed on a widespread erosion surface S4, c. 473 100 km from the present day coastline. Reconstruction of the paleo-fluvial network 474 reveals the sinuosity of the valleys and the potential riffle-pool morphology. These 475 incisions are the same described by Pinot (1974) but considering their stratigraphic 476 position, a Quaternary age is unlikely. Based on the seismic interpretation of the AS 477 stratigraphy, correlation to the eustatic sea-level curve of Hag et al. (1987), and from 478 comparison to other margins, we propose that the unconformity S4 is rather of mid-479 Neogene age. We also assume that the relative sea-level fall required for creating 480 such fluvial incisions on an exposed shelf was predominantly controlled by eustasy. 481 As an angular unconformity, S4 developed contemporaneously with a tilting of the

482 margin related to the Miocene Alpine orogenic phase. The associated uplift may have 483 increased the relative sea-level fall in the area, thus favoring the emersion of the shelf. 484 The most suitable eustatic fall event in terms of timing and amplitude initiated in the 485 Serravallian and culminated at the Serravallian-Tortonian boundary. Therefore we 486 proposed an age of c. 11.6 Ma for the completion of these incisions. Such an 487 attribution needs to be validated by new samples as other potential ages are proposed 488 in other studies for surrounding analogues. These valleys correlate well in both age 489 and direction with mid-Miocene incised valleys described onshore, over Brittany. The 490 question remains on the age of the incision phase of the inner shelf Quaternary incised 491 valleys. This improves our knowledge of the local Miocene paleogeography. Together 492 with recent results on the North Aquitaine shelf, these results provide new arguments 493 that tend to confirm the significance of the eustatic fall at the Serravallian-Tortonian 494 transition and its record on the northern margin of the Bay of Biscay. The expression 495 of the Serravallian-Tortonian boundary on the AS margin marks the beginning of high 496 amplitude eustatic fluctuations (icehouse) culminating during Pleistocene and creating 497 its present day plateau morphology (polygenetic wave-planed surface). The variation 498 in the depth of incision characteristic of riffle-pool morphology limits the ability to 499 quantify the relative and/or absolute sea-level fall. We finally point out that the 500 detailed knowledge of the valley three-dimensional morphology (riffle-pool, 501 meander) is requested before interpreting the significance of the incision depth in 502 terms of local and global controls. Additional data acquisition and interpretation are 503 requested to better constrain the geometry of the network and the relationship with the 504 shelf edge and potential interactions with Miocene slope canyons.

ACKNOWLEGMENTS: This study has been funded by société Lafarge and by the
Université de Bretagne-Sud, within the framework of the PERI-ARMOR project,

| 507 | (Chief scientist: Dr. David Menier). The authors would like to thank Craig Fulthorpe  |
|-----|---|
| 508 | and an anonymous reviewer for their pertinent comments on the earlier versions of the |
| 509 | manuscript. The seismic acquisition material has been provided by the University of   |
| 510 | Rennes 1 – Géosciences Rennes (Dr. Jean-Noël Proust). The authors also wish to        |
| 511 | thank Alexandre Dubois, Florent Scalliet, and Camille Traini for their collaboration, |
| 512 | discussions, and assistance during the GeoEtel-2008 survey. We are also indebted to   |
| 513 | the crewmembers of the CNRS-INSU vessel —Côt de la Manche".                           |
| 514 | <b>REFERENCES:</b>  |
| 515 | Andreieff, P., Bouysse, P., Horn, R., L'Homer, A., 1968. Données récentes sur         |
| 516 | l'Eocène au large de la Bretagne méridionale. Compte Rendu Sommaire des Séances       |
| 517 | de la Société Géologique de France, v. 5, 161-162.                                    |
| 518 |   |

- 519 Anell, I., Thybo, H., Artemieva, I.M., 2009. Cenozoic uplift and subsidence in the
- 520 North Atlantic region: Geological evidence revisited. Tectonophysics. 474, 78-105.

522 Audrun, C., Lefort, J-P., 1977. Géologie du plateau continental sud armoricain entre

523 les îles de Glénan et de Noirmoutier. Implications géodynamiques. Bulletin de la

524 Société Géologique de France. 7, 395-404.

525

526 Barbaroux, L., Blondeau, A., Margerel, J-P., 1971. Présence d'Yprésien fossilifère sur

527 le plateau continental à l'Ouest du plateau du Four (Loire-Atlantique). Comptes

528 Rendus de l'Académie des Sciences. 273, 12-15.

| 531 | Bellec, V.K., Cirac, P., Faugères, J-C., 2009. Formation and evolution of paleo-   |
|-----|--|
| 532 | valleys linked to a subsiding canyon, North Aquitaine shelf (France). Comptes      |
| 533 | Rendus de l'Académie des Sciences, Paris, doi:10.1016/j.crte.2008.07.016.          |
| 534 |  |
| 535 | Besson, D., Parize, O., Rubino, J-L., Aguilar, J-P., Aubry, M-P., Beaudoin, B.,    |
| 536 | Berggren, W.A., Clauzon, G., Crumeyrolle, P., Dexcote, Y., Fiet, N., Iaccarino, S. |
| 537 | Jime'nez-Moreno, G., Laporte-Galaa, C., Michaux, J., Von Salis, K., Suc, J-P.,     |
|     |  |

- .,
- 538 Reynaud, J-Y., Wernli, R., 2005. Latest Burdigalian network of fluvial valleys in
- 539 southeast France (western Alps), characteristics, geographic extent, age, implications:
- 540 Comptes Rendus Geosciences, 337, 1045–1054.

541

- 542 Boillot, G., Bouysse, P., Lamboy, M., 1971. Morphology, sediments and Quaternary
- 543 history of the continental shelf between the Straits of Dover and Cape Finisterre.
- 544 ICSU/SCOR Working Party 31 Symposium. Cambridge, U.K. In: Delay, F.M. (Ed.),
- 545 The Geology of the East Atlantic Continental Margin, Institute of Geological
- 546 Sciences. 70, 75–90.

- 548 Bonnet, S., Guillocheau, F., Brun, J-P., Van Den Driessche, J., 2000. Large-scale
- 549 relief development related to Quaternary tectonic uplift of a Proterozoic-Paleozoic

| 550 | basement: the Armorican Massif, NW France. Journal of Geophysical Research. 105 | 5, |
|-----|---|----|
| 551 | 19,273–19,288.  |    |

| 553 | Bourillet, J-F., | Revnaud, J- | Y., Baltzer, A | ., Zaragosi, S., | 2003. The | Fleuve Manche': |
|-----|------------------|-------------|----------------|------------------|-----------|-----------------|
|     |                  | , .,,.      |                |                  |           |                 |

- the submarine sedimentary features from the outer shelf to the deep-sea fans. Journal
- 555 of Quaternary Science. 18, 261–282.

556

| Bourillet, J-F., Turon, J-L., 2003. Rapport scientifique de la | mission |
|--|---------|
|--|---------|

558 MD133/SEDICAR. Les rapports de campagne à la mer, IPEV, Brest, OCE/2003/04,

559 150 pp.

560

| 561 | Bourillet, . | J-F., Menier | , D., | Gaborit, 1 | К., | 2005. | Architecture | des | sédiments | quaternaires |
|-----|--------------|--------------|-------|------------|-----|-------|--------------|-----|-----------|--------------|
|-----|--------------|--------------|-------|------------|-----|-------|--------------|-----|-----------|--------------|

562 et vallées incises de la marge sud-armoricaine. In: Abstract Volume 10ème Congrès

563 Français de Sédimentologie. ASF, Paris, Presqu'île de Giens, 11-13 October 2005,.

564 42.

565

| 566 | Bourillet, J-F., | Zaragosi, S., | Mulder. T. | . 2006. Th | e French A | tlantic marg | in and the |
|-----|------------------|---------------|------------|------------|------------|--------------|------------|
|     |                  |               |            | ,          |            |              |            |

deep sea submarine systems. Geo-Marine Letters. 26, 311-315.

568

- 569 Bourrouilh, R., Richert, J-P., Zolnaï, G., 1995. The North Pyrenean Aquitaine basin,
- 570 France: evolution and hydrocarbons. AAPG Bulletin. 79, 831–853.)

| 572 | Bouysse, P., Horn, R., 1971. Etude structurale du plateau continental au large des  |
|-----|---|
| 573 | côtes méridionales de la Bretagne. Cahiers Océanographiques. 23, 497-517.           |
| 574 |   |
| 575 | Bouysse, P., Horn, R., Le Gorgeu, J-P., 1968. Résultats d'une prospection sismique  |
| 576 | réflexion continue en Bretagne méridionale, entre Penmac'h et Belle-Île. Comptes    |
| 577 | Rendus de l'Académie des Sciences, Paris. 267, 568-571.                             |
| 578 |   |
| 579 | Bouysse, P., Chateauneuf, J-J., Ters, M., 1974. Présence d'Yprésien, niveau         |
| 580 | transgressif et taux de sédimentation flandriens en baie de Vilaine (Bretagne       |
| 581 | méridionale). Comptes Rendus de l'Académie des Sciences, Paris. 279, 1421-1424.     |
| 582 |   |
| 583 | Brault, N., Bourquin, S., Guillocheau, F., Dabard, M-P., Bonnet, S., Courville, P., |
| 584 | Estéoule-Choux, J., Stepanoff, F., 2004. Mio-Pliocene to Pleistocene                |
| 585 | paleotopographic evolution of Brittany (France) from a sequence stratigraphic       |
|     |   |

analysis: relative influence of tectonics and climate. Sedimentary Geology. 163, 175-210.

588

| 589 | Chantraine, J., Autran, | A., Cavelier, | C. and others, 1996. | Carte géologique de la |
|-----|-------------------------|---------------|----------------------|------------------------|
|-----|-------------------------|---------------|----------------------|------------------------|

590 France à 1/1 000 000, Revised 2003. Editions BRGM, Orléans.

591

| 592 | Chaumillon, | Е., | Proust, | J-N., | Menier, | D., | Weber, | N., | 2008. | Incised | -vall | ey |
|-----|-------------|-----|---------|-------|---------|-----|--------|-----|-------|---------|-------|----|
|     | ,           |     | ,       |       |         |     |        |     |       |         |       | ~  |

593 morphologies and sedimentary-fills within the inner shelf of the Bay of Biscay

594 (France): A synthesis. Journal of Marine Systems. 72, 383-396.

595

| 596 | Dalrymple. | R.W | Bovd. | R., | Zaitlin, | B. | R., | 1994. | Histor | v of research. | types and                               |
|-----|------------|-----|-------|-----|----------|----|-----|-------|--------|----------------|---|
|     |            |     |       | ,   |          |    | ,   | - / / |        | /              | , |

- 597 internal organization of incised-valley systems: Introduction to the volume, *in*
- 598 Dalrymple, R.W., et al., eds., Incised-valley systems: Origins and sedimentary
- sequences. SEPM (Society for Sedimentary Geology) Special Publication. 51, 3–10.

600

601 Delanoë, Y., 1988. Les grands traits de la structure et de l'évolution géodynamique

602 des dépôts tertiaires du plateau continental sud-armoricain d'après les enregistrements

603 de sismique réflexion. Géologie de la France. 1988-1, 79-90.

604

605 Delanoë, Y., Pinot, J-P., 1974. Etude structurale du Tertiaire de la région du Banc

606 Bertin entre Belle-Ile et les Iles de Glénan (Bretagne méridionale). Union des

607 Océanographes de France. 6, 59-65.

608

- 609 Delanoë, Y., Lehébel, L., Margerel, J-P., Pinot, J-P., 1975. La Baie de Concarneau est
- 610 un bassin tectonique dans lequel d'épais dépôts du Lutétien supérieur ont été
- 611 conserves. Comptes Rendus de l'Académie des Sciences. 281, 1947-1950.

612

| 613 | Delanoë, Y., Margerel, J-P., Pinot, J-P., 1976. En Baie de Concarneau, l'Oligocène    |
|-----|---|
| 614 | marin est discordant sur un Eocène ondulé, faillé et érodé, et l'Aquitanien a voilé   |
| 615 | l'ensemble après une nouvelle pénéplanation. Comptes Rendus de l'Académie des         |
| 616 | Sciences. Paris. 282, 29-32   |
| 617 |   |
| 618 | Evans, C.D.R., 1990. United Kingdom Offshore Regional Report: the Geology of the      |
| 619 | Western English Channel and its Western Approaches. British Geological Survey,        |
| 620 | NERC, HMSO: London.   |
| 621 |   |
| 622 | Evans, C.D.R., Hughes, M.J., 1984. The Neogene succession of the South Western        |
| 623 | Approaches, Great-Britain. Journal of the Geological Society of London, 141(2), 315-  |
| 624 | 326.  |
| 625 |   |
| 626 | Fulthorpe, C.S., Austin, J.A., Jr., Mountain, G.S., 1999. Buried fluvial channels off |
| 627 | New Jersey: Did sea-level lowstands expose the entire shelf during the Miocene?.      |
| 628 | Geology. 27, 203–206.   |
| 629 |   |
| 630 | Fulthorpe, C.S., Austin, J.A., Jr., Mountain, G.S., 2000. Morphology and distribution |
| 631 | of Miocene slope incisions off New Jersey: Are they diagnostic of sequence            |
| 632 | boundaries?. Geology. 112, 817-828.   |

| () ( | $\Omega / 1$ ID | 0 1 1    | 17 2000            | T(1)         | 1/ /       | 1 .        | 1      | •       | 1      |
|------|-----------------|----------|--------------------|--------------|------------|------------|--------|---------|--------|
| 634  | (Telv L-P       | Nztrakos | к /()()()          | I 'evolution | nalengengr | anhiaile e | t geod | vnamiau | ie dii |
| 051  | Gery, 5 I .,    | oznakos, | <b>IX</b> ., 2000. |              | pullogeogn | upingue e  | i goou | ynannqu | ic uu  |

Bassin aquitain au Paléogène : enregistrement et datation de la tectonique pyrénéenne.

636 Géologie de la France. 2000-2, 31–57.

637

638 Gély, J-P., Sztràkos, K., 2001. La tectonique pyrénéenne à l'Oligocène : une phase

639 majeure de déformation en compression méconnue du Bassin aquitain (France).

640 Comptes Rendus de l'Académie des Sciences, Paris. 332, 507–512.

641

642 Guillocheau, F., Bonnet, S., Bourquin, S., Dabard, M-P., Outin, J-M., Thomas, E.,

643 1998. Mise en évidence d'un réseau de paléovallées ennoyées (paléorias) dans le

644 Massif armoricain: une nouvelle interprétation des sables pliocènes armoricains.

645 Comptes Rendus de l'Académie des Sciences, Paris. 327, 237–243.

646

- 647 Guillocheau, F., Robin, C., Allemand, P., Bourquin, S., Brault, N., Dromart, G.,
- 648 Friedenberg, R., Garcia, J-P., Gaulier, J-M., Gaumet, F., Grosdoy, B., Hanot, F., Le
- 649 Strat, P., Mettraux, M., Nalpas, T., Prijac, C., Rigollet, C., Serrano, O., Granjean, G.,
- 650 2000. Meso-Cenozoic geodynamic evolution of the Paris Basin : 3D stratigraphic
- 651 constraints. Geodynamica Acta. 13, 189-246.

652

- 653 Guillocheau F., Brault N., Thomas E., Barbarand J., Bonnet S., Bourquin S., Estéoule-
- 654 Choux J., Guennoc P., Menier D., Néraudeau D., Proust J-N., Wyns R., 2003. Histoire

- 655 géologique du Massif Armoricain depuis 140Ma (Crétacé-Actuel). Bulletin
- d'Information des Géologues du Bassin de Paris. 40, 13-28.

- Hailwood, E.A., Bock, W., Costa, L., Dupeuble, P.A., Müller, C., Schnitker, D., 1979.
- 659 Chronology and biostratigraphy of northeast Atlantic sediments, DSDP Leg 48. In:
- 660 Montardet, L., Roberts, D.G., et al. (Eds.), Initial Report of the Deep Sea Drilling
- 661 Project. Washington. 1119-1141.

662

- 663 Haq, B.U., Hardenbol, J., Vail, P.R., 1987. Chronology of fluctuatingsea levels since
- 664 the Triassic. Science. 235, 1156–1167.

665

- 666 Haq, B.U., Hardenbol, J., Vail, P.R., 1988. Mesozoic Cenozoic chronostratigraphy
- and eustatic cycles. In: Wilgus, C.K., Posamentier, H., Ross, C.K., Kendall, C.G.St.C.
- 668 (Eds.), Sea-level Changes: An Integrated Approach. Society of Economic
- 669 Paleontologists and Mineralogists Special Publication. 42, 71–108.

670

- Hilgen, F., Aziz, H.A., Bice, D., Iaccarino, S., Krijgsman, W., Kuiper, K., Montanari,
- A., Raffi, I., Turco, E., Zachariasse W.-J., 2005. The Global boundary Stratotype
- 673 Section and Point (GSSP) of the Tortonian Stage (Upper Miocene) at Monte Dei
- 674 Corvi. Episodes. 28, 6-17.

- 676 Imbrie, J., Hays, J.D., Martinson, D.G., McIntyre, A., Mix, A.C., Morley, J.J., Pisias,
- 677 N.G., Prell, W.L., Shackleton, N.J., 1984. The orbital theory of Pleistocene climate:
- support from a revised chronology of the marine  $\delta$  18O record. In: Berger A, Imbrie J,
- Hays J, Kukla G, Saltzman B. (Eds.), Milankovitch and climate. Reidel Publishing
- 680 Company, Dordrecht, 269-305.

- 682 John, C.M, Karner, G.D., Mutti, M., 2004.  $\delta^{18}$ O and Marion Plateau backstripping:
- 683 Combining two approaches to constrain late middle Miocene eustatic amplitude.
- 684 Geology. 32, 829-832.

685

Kominz, M.A., Miller, K.G., Browning, J.V., 1998. Long-term and short-term global
Cenozoic sea-level estimates. Geology. 26, 311–314.

688

- 689 Lécuyer, C., Grandjean, P., Paris, F., Robardet, M., Robineau, D., 1996. Deciphering
- 690 « temperature » and « salinity » from biogenic phosphates : the <sup>18</sup>O of coexisting
- 691 fishes and mammals of the Middle Miocene sea of western France. Paleogeography,
- 692 Paleoclimatology, Paleoecology. 126, 61-74.

693

- 694 Lericolais G., Auffret J-P. Bourillet J-F., 2003. The Quaternary Channel River :
- seismic stratigraphy of its palaeo-valleys and deeps. Journal of Quaternary Science.
- 696 18, 245**-**260.

| 698 | Lofi, J., and Berné, S., 2008. Evidence for pre-Messinian submarine canyons on the |
|-----|--|
| 699 | Gulf of Lions slope (Western Mediterranean). Marine and Petroleum Geology. 25,     |
| 700 | 804-817.   |

| 702 | Lofthouse, C., Robert, A., 2008. Riffle-pool sequences and meander morphology. |
|-----|--|
| 703 | Geomorphology. 99, 214-223.  |

704

| 705 | Maillard, J., | 1982. | Rapport d | e fin c | le sondage | Garlizenn-1 | (GEN | l), Mer | Celtique |
|-----|---------------|-------|-----------|---------|------------|-------------|------|---------|----------|
|-----|---------------|-------|-----------|---------|------------|-------------|------|---------|----------|

permit, S.N.E.A.P. (Société National Elf Aquitaine Production). BEPH (Bureau

Exploration-Production des Hydrocarbures) Open file report 14-3527. 20 pp.

708

| 709 | Menier, D. | , 2004. Mor | phologie et | remplissage | des vallées | fossiles sud | -armoricaines: |
|-----|------------|-------------|-------------|-------------|-------------|--------------|----------------|
|     | ,          | /           |             |             |             |              |                |

710 apports de la stratigraphie sismique. Ph.D. Thesis memoir, Université de Bretagne

711 Sud, Mémoires Géosciences Rennes. 110, 202 pp.

712

```
713 Menier, D., Reynaud, J-Y., Proust, J-N., Guillocheau, F., Guennoc, P., Bonnet, S.,
```

714 Tessier, B., Goubert, E., 2006. Basement control on shaping and infilling of valleys

715 incised at the southern coast of Brittany, France. Society of Economic Paleontologists

and Mineralogists Special Publication. 85, 37-55.

717

| 718 | Miller, K.G. | Mountain, | G.S., Bro | owning, | J.V., | Kominz, | M., \$ | Sugarman, | P.J., | Christie- |
|-----|--------------|-----------|-----------|---------|-------|---------|--------|-----------|-------|-----------|
|     | , , ,        | , ,       | ,         | 0,      | ,     | ,       | ,      | 0,        | ,     |           |

719 Blick, N., Katz, M. E., Wright, J.D., 1998. Cenozoic global sea level, sequences, and

| 720 | the New Jersey Transect: results from coastal plain and continental slope drilling. |
|-----|---|
| 721 | Reviews of Geophysics. 36, 569-601.   |

| 723 | Miller, K.G., Kominz, M., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E.,  |
|-----|--|
| 724 | Sugarman, P.J., Cramer, B.S., Christie-Blick, N., Pekar, S.F., 2005. The Phanerozoic |
| 725 | record of sea-level change. Science. 310, 1293-1298.                                 |
| 726 |  |
| 727 | Miller, K.G. Browning, J.V., Aubry, M-P., Wade, B.S., Katz, M.E., Kulpecz, A.A.,     |
| 728 | Wright, J.D., 2008. Eocene-Oligocene global climate and sea-level changes: St.       |
| 729 | Stephens Quarry, Alabama. Geological Society of America Bulletin. 120, 34-53.        |
| 730 |  |
| 731 | Montadert L., Roberts D.G., De Charpal O., Guennoc P., 1979. Rifting and             |
| 732 | subsidence of the northern continental margin of the Bay of Biscay. in : Usher X.    |
| 733 | (Ed.), Initial Reports of the Deep Sea Drilling Project, Washington. 48, 1025–1060.  |
| 734 |  |
| 735 | Néraudeau, D., 2003. Lithologies, Faunes et Paléogéographies des Dépôts de Type      |
| 736 | Falun. Bulletin d'Information des Géologues du Bassin de Paris. 40, 6-12.            |
| 737 |  |
| 738 | Olivet J-L., 1996. La cinématique de la plaque Ibérique. Bulletin des Centres de     |
| 739 | Recherche, d'Exploration et de Production d'Elf- Aquitaine. 21, 131–195.             |
| 740 |  |

| 741 | Pigram, C.J., Davies, P.J., Feary, D.A., Symonds, P.A., 1992. Absolute magnitude of  |
|-----|--|
| 742 | the second-order middle to late Miocene sea-level fall, Marion Plateau, Northeast    |
| 743 | Australia. Geology. 20, 858-862.   |
| 744 |  |
| 745 | Pinot, J-P., 1974. Le Précontinent Breton entre Penmac'h, Belle-Ile et l'escarpement |
| 746 | continental, Etude Géomorphologique, Lannion, Impram, 256 pp.                        |
| 747 |  |
| 748 | Posamentier, H.W., 2001. Lowstand alluvial bypass systems: incised vs. unincised.    |
| 749 | American Association of Petroleum Geologists Bulletin. 85, 1771-1793.                |
| 750 |  |
| 751 | Posamentier, H.W., P.R. Vail, 1988. Eustatic controls on clastic deposition II-      |
| 752 | sequence and systems tract models. In: C. K. Wilgus, B. S. Hastings, C. G. St. C.    |
| 753 | Kendall, H. W. Posamentier, C. A. Ross, J. C. Van Wagoner, eds., Sea level change-   |
| 754 | an integrated approach: Society of Economic Paleontologists and Mineralogists        |
| 755 | Special Publication. 42, 125–154.  |
| 756 |  |
| 757 | Powell, A.J., 1988. A preliminary investigation into the Neogene dinoflagellate cyst |
| 758 | biostratigraphy of the British Southwestern Approaches. Bulletin des Centres de      |
| 759 | Recherche, Exploration et Production Elf-Aquitaine. 12, 277–311.                     |
| 760 |  |
| 761 | Praeg, D., Stokerb, M.S., Shannona, P.M., Ceramicolac, S., Hjelstuend, B., Laberge,  |

762 J.S., Mathiesenf A., 2005. Episodic Cenozoic tectonism and the development of the

| 763 | NW European _pæsive' continental margin. Marine and Petroleum Geology. 22, |
|-----|--|
| 764 | 1007–1030.   |

| 766 | Preux. | R.,1978. | Rapport of | le fin de | sondage | Penma-1, | Loire | Maritime | permit. |
|-----|--------|----------|------------|-----------|---------|----------|-------|----------|---------|
|     | ,      |          |            |           |         |          |       |          | p       |

767 S.N.E.A.P. (Société National Elf Aquitaine Production). BEPH (Bureau Exploration-

768 Production des Hydrocarbures) Open file report 14-3435. 15 pp.

769

- 770 Proust, J-N., Renault, M., Guennoc, P., Thinon, I., 2009. Sedimentary architecture of
- the Loire drowned valleys of the French Atlantic shelf. Journal of Sedimentary
- 772 Research. In press.
- 773
- Reynaud, J-Y., Tessier, B., Proust, J.N., Dalrymple, R., Bourillet, J-F., De Batist, M.,
- 775 Lericolais, G., Berné, S. Marsset, T., 1999. Architecture and sequence stratigraphy of a
- 176 late neogene incised valley at the shelf margin, Southern Celtic Sea. Journal of
- 777 Sedimentary Research, 69, 351-364.

778

Richards, K.S., 1982. Rivers: Form and Process in Alluvial Channels: New York,Methuen, 361 pp.

781

- 782 Stewart, H.A., Davies, J.S., 2007. SW Approaches MESH Survey, R/V Celtic
- 783 Explorer Cruise CE0705, BGS Project 07/06, Operations Report. British Geological
- 784 Survey Commissioned Report, CR/07/123.

| 786 | Tcherepanov, E.N., Droxler, A.W., Lapointe, P., Mohn, K., 2008. Carbonate seismic       |
|-----|---|
| 787 | stratigraphy of the Gulf of Papua mixed depositional system: Neogene stratigraphic      |
| 788 | signature and eustatic control. Basin Research. 20, 185-209.                            |
| 789 |   |
| 790 | Thinon, I., 1999. Structure profonde de la Marge Nord Gascogne et du Bassin             |
| 791 | Armoricain. Ph.D. Thesis memoir, Université de Bretagne Occidentale.                    |
| 792 |   |
| 793 | Thinon, I., Menier, D., Guennoc, P., Proust, J-N., Guillocheau, F., Bonnet, S., Le Roy, |
| 794 | P., Augris, C., Bourillet, J-F., Baltzer, A., Tessier, B., Pastol, Y., Garlan, T.,      |
| 795 | Bechennec, F., Le Metour, J., Graviou, P., Alix, A.S., Cornu, S., Loget, N., Scalliet,  |
| 796 | F., 2008. Carte géologique de la France à 1 / 250 000 de la marge continentale,         |
| 797 | Lorient, Bretagne Sud. Éditions BRGM-CNRS.  |
| 798 |   |
| 799 | Thomas, E., 1999. Évolution Cénozoïque d'un domaine de socle : le Massif                |
| 800 | Armoricain. Ph.D. unpublished Thesis memoir, Université de Rennes 1. 148 p.             |
| 801 |   |
| 802 | Vail, P.R., Mitchum, J.R., Todd, R.G., Widmier, J.M., Thompson, S., Sangree, J.B.,      |
| 803 | Bubb, J.N., Hatlelid, W.G., 1977. Seismic stratigraphy and global changes of sea level  |
| 804 | in depositional sequences. In Seismic Stratigraphy – applications to hydrocarbon        |
| 805 | exploration, Payton, E. (Ed): American Association of Petroleum Geologist Memoir.       |
| 806 | 26, 49-212.   |

| 808 | Vail, P. R., Audemard, F., Bowman, S. A., Eisner, P. N., Perez-Cruz, C., 1991. The     |
|-----|--|
| 809 | stratigraphic signatures of tectonics, eustasy and sedimentology-An overview, in:      |
| 810 | Einsele, G., et al., eds., Cycles and events in stratigraphy: Berlin, Springer-Verlag. |
| 811 | 617–659.   |

- 813 Vanney J. R., Horn R., Martin G., 1972. La disposition des séquences sédimentaires
- 814 post-mésozoïques sous la partie externe de la plate-forme continentale sud-
- armoricaine (entre 45°40' et 46°40' lat. Nord). Compte Rendus de l'Académie des
- 816 Sciences. 275, 2829-2832.

817

- 818 Van Vliet-Lanoë, B., Laurent, M., Hallégouët, B., Margerel, J-P., Chauvel, J-J.,
- 819 Michel, Y., Moguedet, G., Trautman, F., Vauthier, S., 1998. Le Mio–Pliocène du
- 820 Massif armoricain. Données nouvelles. Comptes Rendus de l'Académie des Sciences,

821 Paris. 326, 333–340.

822

- 823 Waelbroeck, C. Labeyrie, L., Michel, E., Duplessy, J-C., McManus, J.F., Lambeck,
- K., Balbon, E., Labracherie, M, 2002. Sea-level and deep water temperature changes
- derived from benthic foraminifera isotopic records. Quaternary Science Reviews. 21,295-305.

| 828 | Zachos, J., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and |
|-----|--|
| 829 | aberrations in global climate 65 Ma to present. Science. 292, 686-693.                 |
| 830 |  |
| 831 | Ziegler, P.A., 1987. Evolution of the Western Approaches Trough. Tectonophysics,       |
| 832 | 137, 341-346.  |
| 833 |  |
| 834 | Ziegler, P.A., 1990. Collision related intra-plate compression deformations in western |

and central Europe. Journal of Geodynamic. 11, 357-388.

836

# 837 FIGURE CAPTIONS:

838

| 839 | <b>Figure 1:</b> A- Map showing the location of the study area in the Bay of Biscay, W.A.: |
|-----|--|
| 840 | Western Approaches, A.S.: Armorican Shelf, Aq.S.: Aquitaine Shelf; B – Simplified          |
| 841 | geological map covering the Armorican Shelf, showing the extent of the main                |
| 842 | geological units and the position of exploration well and offshore cores with retrieved    |
| 843 | Cenozoic material (Andreieff et al., 1968; Barbaroux et al., 1971; Bouysse et al.,         |
| 844 | 1974; Delanoë and Pinot, 1974; Delanoë et al., 1975; Delanoë et al., 1976). Thin           |
| 845 | dashed lines correspond to the position of geophysical data (see fig. 2 for details). The  |
| 846 | Bold lines correspond to the profiles displayed on figure 3; C – Cross section of the      |
| 847 | Armorican Shelf showing the main stratigraphic succession from Cretaceous to               |
| 848 | present. The vertical axis is in second two-way travel time (TWTT) (modified from          |

849 Vanney et al., 1972; Guillocheau et al., 2003, Thinon et al., 2008). See Fig. 1B for850 location.

851

| 050 |                    | 1              |                 | 1 0 0      | 1 /        | • •        |
|-----|--------------------|----------------|-----------------|------------|------------|------------|
| 857 | Figure 7. Position | mane chowing f | he extension of | the tour S | narker_tw  | ne ceicmic |
| 052 |                    | maps showing u |                 | inc ioui o | parker-typ |            |
|     | 0                  |                |                 |            |            |            |

surveys (Bir Hacheim 1967; Trophal 2002; GeoEtel 2008) used in this study,

including line numbers and the sections displayed in figures (bold lines).

855



857 description, biostratigraphy correlations established from foraminifer and nannofossil

determinations, and interpretation of depositional environments (adapted from Preux,

859 1978). Top Oligocene and intra Oligocene labels mark two important changes in the

860 depositional environment settings that may correlate to seismic unconformities.

861

862 Figure 4: Interpreted seismic profiles from BH67 (20-670; 20-790; 20-970) and 863 GeoEtel 2008 (A) surveys showing the stacking pattern of the main seismic units (U1 864 to U6b) and unconformities (S1 to S6) from the inner shelf to the shelf break. On 865 profile 20-670, seismic units and unconformities are correlated to the calculated 866 position of top- and intra-Oligocene markers in projected PENMA-1 (c. 3 km from 867 NW). The box on profile GeoEtel-2008 A refers to the section displayed in figure 4a. 868 869 Figure 5: Seismic image and interpreted section of Sparker seismic profiles GeoEtel-870 07 (A) and Job 1974 (3-VIII-74) (B) (see Figure 1 for location) showing the

interpreted seismic units (U2 to U6b) and unconformities (S2 to S6). Three buried
incised valleys are identified on S4 and labeled A, B and C. Valley B is displayed in
detail in figure 6 as indicated by boxes. See figures 2 and 4 for Location of profiles.
Note that the zoom on Fig. 6a. is a crop of GeoEtel 2008 – 26 but is indicated by a
box on Job 1974 (3-VIII-74) as both profiles are similar in location.

876

877 Figure 6: Details of high-resolution Sparker profiles from GeoEtel-2008 survey

878 crossing the incised fluvial valleys of S4 (valley B), from profiles 26 (A), 22 (B) and

879 07 (C). Profiles show the main incision channel (MC) and associated tributaries (Tr)

separated by terraces (Te) in a wider valley. On profile 07 (C) other younger

channelized incisions (YCI) are visible that cut the major erosion surface S4. Vertical

axis in ms (TWTT: Two Way Travel Time) and horizontal axis shows Shot Point

number (SP). See Figs. 1 and 3 for location.

884

Figure 7: Oblique three-dimensional view from North showing the interpreted
seismic profiles GeoEtel 07, 21, 22, 25 to 29 and Trophal 50 (labels) and the fluvial
incisions developed on the major erosion surface S4 (bolder line). The dotted lines
correspond to proposed reconstructions of the incised valleys B and C. Vertical axis in
ms (TWTT: Two Way Travel Time) and horizontal axis shows Shot Point number
(SP).

891

Figure 8: Map of the Armorican Shelf showing (1) the cartographic trace of S4within the Miocene series, (2) the reconstruction of the buried fluvial network with

| 894 | the four valleys (A, B, C and D), (3) the mid-Miocene (Serravallian-Tortonian) fluvial |
|-----|--|
| 895 | remnants identified onshore at Lauzach and Réguigny (Brault et al., 2004), and (4) the |
| 896 | -Quaternary incised valleys" (Menier, 2004; Menier et al., 2006; Thinon et al., 2008). |
| 897 | The Pleistocene incised fluvial network is preserved offshore until -70 m where        |
| 898 | incision starts to fade over the mid-shelf (Menier et al., 2006). Direct connection    |
| 899 | between networks is precluded by erosion onshore and by the major erosion surface      |
| 900 | S6 of the AS offshore.   |

| 902 | Figure 9: Correlation of the AS seismic stratigraphy to (1) the Western Approaches       |
|-----|--|
| 903 | seismic stratigraphy (e.g. Bourillet et al., 2003), (2) the onshore fluvial stratigraphy |
| 904 | (Van Vliet-Lanoë et al., 1997, Guillocheau et al., 1998; Brault et al., 2004), (3) the   |
| 905 | PENMA-1 section (Preux, 1978), (4) the global sea level curve (modified from Haq et      |
| 906 | al., 1987), (5) the main orogenic phases affecting the western Europe, and (6) the       |
| 907 | Stratigraphic chart (ICS, 2008).   |
| 908 |  |

Table 1: Characteristics of the seismic units (U1 to U6b) and bounding surfaces (S1 to

S6) recognized on the data covering the AS with seismic description, Lithological

911 interpretation and examples from GeoEtel-2008 survey.

Figure 1 Click here to download high resolution image





#### Figure 3 Click here to download high resolution image





# Table 1Click here to download high resolution image

| Seismic<br>units<br>U1 | Descriptions of seismic facies  | Interpretations  | Seismic image samples (crops from GeoEtel-2008)     |  |  |
|------------------------|---|--|---|--|--|
|                        | reflection terminations   | correlations   | Scale bars: horizontal $-250$ m ; vertical $=20$ ms |  |  |
|                        | Very low to average continuity,<br>high amplitude and frequency,<br>subparallel to chaotic<br>configuration. Highly deformed<br>Base: not determined<br>Top: eroded by S1 (top lap)   | Crystalline and<br>metamorphic<br>basement<br>(Paleozoic)  | A-35  |  |  |
| U2                     | Good continuity, average<br>frequency, average to<br>high amplitude, parallel<br>configuration. Faulted & folded<br><b>Base</b> : onlap on S1<br><b>Top</b> : eroded by S2 (toplap)   | Well-bedded<br>marine deposits<br>identified as Paleocene<br>and Eocene chalky<br>calcareous mudstones<br>in various cores and<br>in PENMA-1 well                          | 8-06  |  |  |
| U3                     | Average continuity, average<br>frequency, very high amplitude,<br>subparallel to wavy configu-<br>ration. Slightly faulted & folded<br>Base: concordant on 52<br>Top: concordant with 53  | Bedded marine deposits<br>with mounds (coral<br>patches?) correlated<br>to Oligocene bioclastic<br>sandstones with corals,<br>bivalves and bryozoans<br>in PENMA-1 well    | C-07  |  |  |
| U4                     | Good continuity, average<br>frequency, low amplitude,<br>subparallel configuration with<br>clinoforms, Rarely faulted &<br>folded<br>Base: downlap on 53<br>Top: eroded by S4 with incisions  | Well-bedded<br>marine deposits corres-<br>ponding to mixt terri-<br>geneous / calcareous<br>Miocene mudstones<br>in core samples and<br>PENMA-1 well                       | D- 07 0   |  |  |
| U5                     | Very low continuity, average<br>frequency, very low amplitude,<br>chaotic configuration with cha-<br>nnels. Rarely faulted & folded<br><b>Base</b> : onlap on S4<br><b>Top</b> : eroded upward and down-<br>ward by S5 with incisions | Pourly-bedded coarse<br>marine and channel<br>fill (fluvial) LST<br>deposits of probable<br>Late Miocene age.  | E · 20t   |  |  |
| U6                     | Very low continuity, average<br>frequency, low amplitude,<br>chaotic configuration with<br>channels & clinoforms. Rarely<br>faulted & folded<br><b>Base</b> : concordant/onlap on 55<br><b>Top</b> : eroded by 56 (toplap)            | Pourly-bedded coarse<br>marine deposits<br>with rare channels<br>corresponding to the<br>outer-shelf Pliocene -<br>Pleistocene sediment<br>wedge (Red sands)               | F-11  |  |  |
| U6b                    | Transparent to low continuity,<br>low amplitude. Not deformed<br>Base: concordant/downlap on S6<br>Top: sea floor   | Poorly-bedded muddy<br>sands of disconnected<br>sediment bodies of the<br>Pleistocene - Holocene<br>"Grande Vasière", depo-<br>sited and reworked<br>during LST and/or TST | G-07  |  |  |



Figure 6 Click here to download high resolution image







#### Figure 9 Click here to download high resolution image

