



Comparison of arsenic and mercury release linked to iron and sulfur biogeochemical processes

André Burnol, Jennifer Hellal, Claire Bény, Jacques-André Bourdoiseau, Catherine Crouzet, Catherine Jouliau, Fabienne Battaglia-Brunet

► To cite this version:

André Burnol, Jennifer Hellal, Claire Bény, Jacques-André Bourdoiseau, Catherine Crouzet, et al.. Comparison of arsenic and mercury release linked to iron and sulfur biogeochemical processes. European Geosciences Union General Assembly 2011, Apr 2011, Vienna, Austria. hal-00581251v1

HAL Id: hal-00581251

<https://brgm.hal.science/hal-00581251v1>

Submitted on 30 Mar 2011 (v1), last revised 13 May 2011 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Comparison of arsenic and mercury release linked to iron and sulfur biogeochemical processes

André Burnol (1), Jennifer Harris-Hellal (1), Jacques-André Bourdoiseau (2), Catherine Crouzet (1), Claire Beny (1), and Fabienne Battaglia-Brunet (1)

(1) BRGM, EPI, Orléans, France (a.burnol@brgm.fr), (2) Université de La Rochelle, LEMMA, La Rochelle, France

Understanding and predicting the mobility of redox-sensitive metals, such as arsenic (As) and mercury (Hg), in aquifers is important to reduce their adverse effects. Both these metals have a strong affinity for iron (hydr)oxides that control their sequestration and their bioavailability in soils and aquifers, in strong relation to microbial activity and physico-chemical conditions.

The aim of this work was to study the mineralogical transformations, induced by bacterial consortia collected in polluted environments, leading to arsenic or mercury release and speciation in controlled column systems under anoxic conditions.

The experimental setup consists of columns filled in the lower half with sand and in the upper half with a mixture of sand and iron oxides, initially doped with either arsenite As(III) or mercury Hg(II). The system is inoculated with the bacterial consortia of polluted aquifers and a nutritive medium is then circulated through the system in which biological and physico-chemical parameters were followed.

The effect of the activity of sulfate-reducing bacteria on arsenic mobilization was demonstrated at the interface between the Fe-rich zone and the zone without Fe: the biotic reduction of solid Fe(III) or the abiotic reduction by HS⁻ are both possible. The increase of the energy gain for both reducers (iron and sulfate reducers) is also suggested and this phenomenon could explain the elevated As concentrations seen in certain aquifers (e.g. Bengal delta aquifers).

Other column experiments are presently being carried out to elucidate mercury redox transformations in similar conditions: the reduction of Hg(II) to volatile Hg⁰ and methylation to HgCH₃⁻ are both likely to take place in such anoxic conditions and could increase the mobility of mercury.