



HAL
open science

Parallel simulation of three-dimensional convective mixing in long-term geological CO₂ storage in saline aquifers

Mohammed Adil Sbai

► **To cite this version:**

Mohammed Adil Sbai. Parallel simulation of three-dimensional convective mixing in long-term geological CO₂ storage in saline aquifers. EGU General Assembly 2007 - European Geosciences Union, Apr 2007, Vienna, Austria. hal-00589919

HAL Id: hal-00589919

<https://brgm.hal.science/hal-00589919>

Submitted on 2 May 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Parallel simulation of three-dimensional convective mixing in long-term geological CO₂ storage in saline aquifers

M. A. Sbai (1)

(1) BRGM, Water Division, Orléans, France (a.sbai@brgm.fr / Fax: +33 2-38643446)

Accurate numerical modelling of the effective dissolution rate of aqueous CO₂ solutions is important for analysis of long-term storage efficiency and risk assessments. Theoretical and field studies showed the possibility of switching from a diffusion-controlled mode of dissolution to a natural convection mode which is much more efficient for accelerating the dissolution mechanism. However, most of the previous investigations have been restricted to two-dimensional cross-section models similar to Elder (1947) problem. In this work, we developed a high resolution parallel simulator to investigate these mechanisms with grid blocks needed for accurate modelling of convective mixing in realistic three-dimensional configurations. The simulator sequentially couples groundwater flow and CO₂ transport equations by density and viscosity contrasts. Density and viscosity of the aqueous CO₂ solution is predicted by accurate equations of state module validated against previously published experimental data for a wide range of temperatures and pressures of interest. Numerical experiments show that two-dimensional modelling over-predicts the time to onset natural convection and under-predicts the time to achieve ultimate dissolution of the CO₂-free phase in a homogeneous porous medium. The role of aquifer heterogeneity in further enhancing CO₂ dissolution is illustrated for a generic case study. Heterogeneity leads to a disconnection of the otherwise continuous fingers to form CO₂ blobs and ganglia, increasing the effective surface area between CO₂-rich blobs and formation waters. Results suggest that high resolution three-dimensional modelling should be guided in CO₂ storage projects to (a) estimate the onset of natural convection, (b) determine the size of convective instabilities when they occur, and (c) calculating the maximum mixing achieved.