



**HAL**  
open science

## Dating tools applied to mineralized Eocene groundwater, for a sustainable management of the Entre-deux-Mers drinking water, Aquitaine Basin, South western France.

Eline Malcuit, Philippe Négrel, Christophe Innocent, Emmanuelle  
Petelet-Giraud, Pierre Durst

### ► To cite this version:

Eline Malcuit, Philippe Négrel, Christophe Innocent, Emmanuelle Petelet-Giraud, Pierre Durst. Dating tools applied to mineralized Eocene groundwater, for a sustainable management of the Entre-deux-Mers drinking water, Aquitaine Basin, South western France.. EGU General Assembly 2011, Apr 2011, Vienna, Austria. hal-00593470

**HAL Id: hal-00593470**

**<https://brgm.hal.science/hal-00593470>**

Submitted on 16 May 2011

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## **Dating tools applied to mineralized Eocene groundwater, for a sustainable management of the Entre-deux-Mers drinking water, Aquitaine Basin, South western France.**

Eline Malcuit (1), Philippe Negrel (2), Christophe Innocent (3), Emmanuelle Petelet-Giraud (3), and Pierre Durst (4)

(1) BRGM, Regional Geological Survey Service Bordeaux, France, e.malcuit@brgm.fr, (2) BRGM, Metrology Monitoring Analysis Department, Orleans, France, p.negrel@brgm.fr, (3) BRGM, Water Department, Orleans, France, e.petelet@brgm.fr, (4) BRGM, Regional Geological Survey Service Bordeaux, France, p.durst@brgm.fr

In the south-west of France, the Eocene aquifer is one of the main resources for irrigation, thermo-mineral water, and mainly for drinking water in the Bordeaux region.

This aquifer is characterized by the presence of a large mineralized area, centered on the Entre-deux-Mers region, between the Garonne and the Dordogne rivers, where the groundwaters show strong mineralization and anomalous levels of critical elements, such as sulphates and fluoride, leading to difficulties of resource exploitation for drinking water supply.

Initiated early 2009, the CARISMEAU 2 project, focuses on the geochemical, multi-isotopic and hydrogeological characterization of this mineralized groundwater sector of the Entre-deux-Mers area. The main objectives of this project are to improve the understanding of the origin of the salinity of this mineralized area and to investigate how these mineralized waters circulate in the Eocene aquifer and/or in this multi-layer aquifer system.

The deposit sequences characterizing the Eocene aquifer system are progradational westward, from detrital deposits to carbonates. The Eocene sands and the Eocene limestones are hydraulically connected, the limit of their extension is located under the city of Bordeaux. The groundwater recharge may occur through the Eocene outcrops located in the north and north-east of this mineralized area of the Entre-deux-Mers, and also by vertical leakage from the Oligocene aquifer.

Furthermore, the second aspect to consider in this saline area concerns the piezometric evolution of the Eocene aquifer years after years. In fact, a trough in the potentiometric surface is noticeable for the Eocene Aquifer, centred under the city of Bordeaux. For years, the decline of the piezometric surface is roughly one meter per year in the center. The cone of pressure relief in this confined aquifer stretches to the east year after year in the same direction, toward the Garonne and the Dordogne rivers.

In autumn 2009, the first investigation of over 50 wells and springs gave preliminary characterizations of the aquifer. Groundwaters from more than 50 boreholes have been sampled. Others groundwater samples come from continuously monitoring, in 2010 - 2011, during long-term pumping of several boreholes for drinking water.

Combined geochemical analysis (major and trace elements) and classical isotopic methods using  $\delta^{18}\text{O}_{\text{H}_2\text{O}}$  and  $\delta^2\text{H}_{\text{H}_2\text{O}}$ ,  $\delta^{34}\text{S}_{\text{SO}_4}$  and  $\delta^{18}\text{O}_{\text{SO}_4}$  is carried out. In addition, an innovative set of isotopic methods using strontium isotopes and more exploratory isotopic methods like boron, lithium, uranium/thorium and radium isotopes are applied on the mineralized area. U/Th results are compared to former  $^{14}\text{C}$  results. Moreover several  $^3\text{H}$  analyses have been performed on 11 samples, including one spring which age and original aquifer were unknown.

These dating tools may help to better understand the complex circulations in the Eocene aquifer, of primary importance for drinking water, and its connections to the others aquifer layers. The results of this dating approach coupled with hydrodynamics models, hydrogeological knowledge and a better understanding of the deposit conditions and mineralization may help to better manage this strategic resource of drinking water.