



HAL
open science

Insight of past hydrological conditions in Holocene fluvial sediments in the Loire River (France) through signatures of stables isotopes

Philippe Négrel, Wolfram Kloppmann

► To cite this version:

Philippe Négrel, Wolfram Kloppmann. Insight of past hydrological conditions in Holocene fluvial sediments in the Loire River (France) through signatures of stables isotopes. EGU General Assembly 2012, Apr 2012, Vienna, Austria. pp.XX. hal-00661115

HAL Id: hal-00661115

<https://brgm.hal.science/hal-00661115>

Submitted on 18 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Insight into past hydrological conditions through signatures of stable isotopes in Holocene Loire River fluvial sediments (France)

Philippe Négrel

Wolfram Kloppmann

BRGM, Orléans, France

Multi-proxy indices ($\delta^{18}\text{O}$, $\delta^{13}\text{C}$, granulometry, mineralogy) in the sediments from a channel infill in the Middle Loire alluvial plain are used to highlight some aspects of the basin evolution over the period from 0 to 10000 yr. BP. The Loire River in central France is 1010 km long and drains an area of 117,800 km². The catchment is composed of plutonic rocks (500 - 300 Ma), a large volcanic area in the upstream section and carbonate deposits (200 - 6 Ma old) in the central part. During the Late-glacial and Holocene, the lacustrine carbonate substratum in the Middle Loire alluvial plain was incised by the Loire River and numerous oxbows and channels related to meander migration have been identified. We investigated a channel fill that spans part of the Boreal, the Atlantic, the Subboreal and probably a large part of the Subatlantic through sediment coring.

The channel fill consists mainly of fine clayey sediments deposited during flooding of the river; there is a total absence of coarse clastics and sandy materials in the 5m core excepted in the lower part. The distribution of particle sizes shows an up-core decrease with a coarser fraction occurring in the first 2/3th of the record. Semi-quantitative XRD analysis of the sediment showed that quartz, K-feldspar, plagioclase and calcite are the main mineral components and that the <2 μm fraction is dominated by illite-smectite and kaolinite. The distribution of the clays highlights an up-core increase of the illite-smectite fraction.

The historical record of isotope ratio variations together with the distribution of particle sizes allows to constrain the evolution of the river dynamics. The strong decrease of the $\delta^{13}\text{C}$ of carbonates in the upper part of the record has been ascribed to the progressive closure of the meander and thus the control of the C isotope signature by biological activity in a local environment. Furthermore, we can postulate that the variations in the carbonate $\delta^{18}\text{O}$ are related to the respective contribution of river water and groundwater, being less dependent of the meander evolution than carbon isotope signatures. The isotope record of the river dynamics also agrees with the variations in the clay-mineralogy.