



HAL
open science

Use of two new Na/Li thermometric relationships for geothermal fluids in volcanic environments

Bernard Sanjuan, Ragnar Asmundsson, Romain Millot, Brach Michel

► To cite this version:

Bernard Sanjuan, Ragnar Asmundsson, Romain Millot, Brach Michel. Use of two new Na/Li thermometric relationships for geothermal fluids in volcanic environments. Goldschmidt Conference 2012, Jun 2012, Montréal, Canada. hal-00662009

HAL Id: hal-00662009

<https://brgm.hal.science/hal-00662009>

Submitted on 22 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Use of two new Na/Li thermometric relationships for geothermal fluids in volcanic environments

BERNARD SANJUAN^{1*}, RAGNAR ASMUNDSSON²,
ROMAIN MILLOT³ AND MICHEL BRACH³

¹ BRGM, Department of Geothermal Energy, Orléans, France,
b.sanjuan@brgm.fr (* presenting author)

² Tiger Energy Services, Taupo, New Zealand,
ragnar.asmundsson@tigerhd.com

³ BRGM, Department of Metrology, Monitoring and Analysis,
Orléans, France, r.millot@brgm.fr, m.brach@brgm.fr

Thermometers such as Silica, Na/K, Na/K/Ca, Na/K/Ca/Mg or $\delta^{18}\text{O}$ ($\text{H}_2\text{O}-\text{SO}_4$), based on empirical or semi-empirical laws derived from chemical equilibrium reactions between water and minerals in the deep reservoirs, are commonly used in geothermal exploration in order to estimate the reservoir temperatures. Unfortunately, these estimations are not always concordant because of processes which can perturb the chemical composition of the fluids during their ascent up to the surface (water mixing, fluid cooling, etc.). Given these discordances, auxiliary thermometers such as Na/Li, based on statistical relationships, were also developed. As Li is rather lowly reactive, the use of this thermometer can give more reliable temperature estimations. Presently, three different Na/Li relationships ([1], [2]) are mainly available according to the fluid salinity and the geological environment (volcanic/granitic and sedimentary rocks).

This study carried out in the framework of the European HITI project (High Temperature Instruments for supercritical geothermal reservoir characterization and exploitation) with the collaboration of ISOR Iceland Geosurvey proposes two new Na/Li thermometric relationships. The first concerns the fluids derived from high-temperature seawater-basalt interaction processes existing in the oceanic ridges and rises as well as in the emerged rifts such as those of Iceland (Reykjanes, Svartsengi and Seltjarnarnes geothermal fields) and Djibouti (Asal-Ghoubbet and Obock geothermal areas). It can be expressed as follows:

$$\text{Log (Na/Li in mol/l)} = 920/(T^\circ\text{K}) + 1.105 \quad (r^2 = 0.994).$$

The second relationship, developed using dilute fluids collected only from Icelandic geothermal wells in the 100-325°C range and surprisingly close to that determined by Fouillac and Michard (1981) for volcanic saline fluids at temperatures higher than 200°C, is:

$$\text{Log (Na/Li in mol/l)} = 1786/(T^\circ\text{K}) - 0.936 \quad (r^2 = 0.976).$$

The uncertainty on temperature estimation is $\pm 25^\circ\text{C}$ for both relationships. These results confirm that the Na/Li ratios not only depend on temperature but also on other parameters. The nature of the reservoir rocks and fluid seems to be the most influent one. Some case studies in the literature and thermodynamic considerations suggest that the Na/Li ratios could be controlled by chemical equilibrium reactions involving different mineral assemblages where illite and micas would be, however, always present.

[1] Fouillac Ch. and Michard G. (1981) *Geothermics*, **10**, n°1, 55-70.

[2] Kharaka and Mariner (1989) *Naeser and McCulloch Eds, Springer-Verlag, New York*, 99-117.