



## **Influence of temperature on the geochemistry of Callovo-oxfordian argillite: batch experiments and modelling at 80°C**

Hélène Gailhanou, Christophe Tournassat, Valérie Laperche, Sylvain Grangeon, Philippe Blanc, Nicolas C.M. Marty, Catherine Crouzet, Benoît Henry, Jean-Marc Greneche, Myriam Kars, et al.

### **► To cite this version:**

Hélène Gailhanou, Christophe Tournassat, Valérie Laperche, Sylvain Grangeon, Philippe Blanc, et al.. Influence of temperature on the geochemistry of Callovo-oxfordian argillite: batch experiments and modelling at 80°C. International meeting "Clays in Natural and Engineered Barriers for Radioactive Waste Confinement", Oct 2012, Montpellier, France. hal-00701811

**HAL Id: hal-00701811**

**<https://brgm.hal.science/hal-00701811>**

Submitted on 18 Dec 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Influence of temperature on the geochemistry of Callovo-Oxfordian argillite: batch experiments and modelling at 80°C**

H. Gailhanou<sup>1</sup>, C. Tournassat<sup>1</sup>, V. Laperche<sup>1</sup>, S. Grangeon<sup>1</sup>, P. Blanc<sup>1</sup>, N. Marty<sup>1</sup>, C. Crouzet<sup>1</sup>, B. Henry<sup>1</sup>, J.M. Grenèche<sup>2</sup>, M. Kars<sup>3</sup>, B. Madé<sup>4</sup>, S. Altmann<sup>4</sup> and E.C. Gaucher<sup>1</sup>.

1. BRGM – 3 av. Claude Guillemin BP6009 45060 Orléans cedex 2, France ([h.gailhanou@brgm.fr](mailto:h.gailhanou@brgm.fr))
2. LUNAM –Université du Maine, Institut des Molécules et Matériaux du Mans, UMR CNRS 6283, 72085 Le Mans cedex ([jean-marc.greneche@univ-lemans.fr](mailto:jean-marc.greneche@univ-lemans.fr))
3. Laboratoire des Fluides Complexes et leurs Réservoirs, Université de Pau et des Pays de l'Adour, Avenue de l'Université, BP1155, 64013 PAU cedex
4. ANDRA – Parc de la Croix Blanche, 1-7 rue Jean Monnet, 92298 Châtenay-Malabry Cedex ([Benoit.Made@andra.fr](mailto:Benoit.Made@andra.fr)) .

## **Introduction**

A repository of high-level nuclear wastes in the Callovo-Oxfordian (COx) clay rock will lead to a temporary increase in temperature locally around the disposal cells, up to about 80°C. Thermally induced modifications of the pore-water chemistry of the clay rock must be assessed since the pore water chemistry can affect the rates of corrosion of metallic materials used in the disposal facility (Gaucher et al., 2009; Altmann, 2008). The chemistry of COx pore water has been studied at ambient temperature (~ 25°C) using modelling approaches constrained by parameter values measured on core samples (Gaucher et al., 2006, 2009) and by measurements of pore water collected by means of specially designed in-situ experiments (Vinsot et al., 2008). In particular, the predictive model from Gaucher et al. (2009) reproduces the measured, in situ pore-water composition without any estimated parameters.

In the present study, carried out as part of a collaborative BRGM/Andra research program, the results of long-term batch experiments performed at 80°C, are compared with compositions predicted using a model based on the one developed for 25°C.

## **Batch experiments**

Two batch experiments have been performed with COx rock samples in contact with synthetic water, for respectively 6 and 20 months. The experiments were prepared at 25°C under anoxic atmosphere in closed reactors, with solution compositions close to the pore water composition. Afterwards, each system was heated at 80°C. During the experiments, measurements of  $P_{CO_2}$  were carried out as this parameter is considered as an indicator of the reactivity of this carbonate-dominated system. At the end of the experiments, the suspension was filtered at 0.1 µm at 80°C, under anoxic atmosphere. The pH was also measured at 80°C. Major cations (Ca, Mg, K, Na, Sr) and anions (notably Cl, SO<sub>4</sub>), TIC and minor and trace elements (notably Si, Al, Fe, Zn, Pb) were analysed.


Concerning the solids, some minor changes were detected in the mineralogy between initial and final solids, essentially partial dissolution of celestite and a small increase in the amount of dolomite. Besides, an interesting change was observed by TEM with the formation of needles of nanocrystalline goethite on pyrite particles for the solid from the 20 months experiment (see figure). Such needles were not observed on the initial solid, although magnetic measurements at low temperature on the sample revealed the presence of nanocrystalline goethite. Moreover, an increasing of the  $Fe^{2+}/Fe^{3+}$

ratio in the octahedral environment of clay minerals, compared with initial ratio, was evidenced by  $^{57}\text{Fe}$  Mössbauer spectrometry.

### Modelling of batch experiments

The predictive model of composition as a function of temperature was developed, based on the Gaucher et al. (2009) model and using the PHREEQC code and Thermoddem database (Blanc et al., submitted). As a first approximation, the model considers a primary mineral assemblage with quartz, pyrite, carbonates (calcite, dolomite and siderite), sulfates (celestite), aluminosilicates (chlorite and illite). Moreover, the model takes into account exchange reactions and dissolution reaction kinetics for the potentially more reactive minerals.

For both cases, at 6 and 20 months, the comparison between estimated and observed concentrations of ions in solution are in rather good agreement for pH, Si, Al, major cations (Ca, Mg, Na, K, Sr) and for sulphates and chlorides. The calculated concentrations of Fe are, on the other hand, too high compared with the measured concentrations, for both experiments. The control on the Fe(II) solubility by siderite was already questioned in Gaucher et al. (2009). This problem is being studied, notably by considering the presence of nanogoethite detected in the initial sample and in the solid after 20 months experiment.



**TEM image of needles of nanocrystalline goethite on a pyrite particle, after 20 months experiment at 80°C.**

### References

- Altmann S. (2008) 'Geo'chemical research: a key building block for nuclear waste disposal safety cases. *J. Contam. Hydrol.* 102, 174–179.
- Blanc P., Lassin A., Piantone P., Azaroual M., Jacquemet N., Fabbri A. and Gaucher E. C. (submitted) Thermoddem: a geochemical database focused on low temperature water/rock interaction and waste materials. *Appl. Geochem.*
- Gaucher E. C., Blanc P., Bardot F., Braibant G., Buschaert S., Crouzet C., Gautier A., Girard J.-P., Jacquot E., Lassin A., Negrel G., Tournassat C., Vinsot A. and Altmann S. (2006) Modelling the porewater chemistry of the Callovian-Oxfordian formation at a regional scale. *C. R. Geosci.* **338**, 917–930.
- Gaucher, E. C., Tournassat, C., Pearson, F. J., Blanc, P., Crouzet, C., Lerouge, C., and Altmann, S., (2009) A robust model for pore-water chemistry of clayrock: *Geochim. Cosmochim. Acta* **73**, 6470–6487.
- Vinsot A., Mettler S. and Wechner S. (2008) In situ characterization of the Callovo-Oxfordian pore water composition. *Phys. Chem. Earth* **33**, S75–S86.