



Extending the prediction of the thermodynamic properties of clay minerals to the trapping of trace elements

Philippe Blanc, Philippe Vieillard, Hélène Gailhanou, Stéphane Gaboreau, Arnault Lassin, Eric C. Gaucher, C.I. Fialips, Eric Giffaut

► To cite this version:

Philippe Blanc, Philippe Vieillard, Hélène Gailhanou, Stéphane Gaboreau, Arnault Lassin, et al.. Extending the prediction of the thermodynamic properties of clay minerals to the trapping of trace elements. International meeting "Clays in Natural and Engineered Barriers for Radioactive Waste Confinement", Oct 2012, Montpellier, France. hal-00701813

HAL Id: hal-00701813

<https://brgm.hal.science/hal-00701813>

Submitted on 26 May 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXTENDING THE PREDICTION OF THE THERMODYNAMIC PROPERTIES OF CLAY MINERALS TO THE TRAPPING OF TRACE ELEMENTS

P. Blanc¹, P. Vieillard², H. Gailhanou¹, S. Gaboreau¹, A. Lassin¹, E.C. Gaucher¹, C. Fialips³, E. Giffaut³

1. BRGM – 3 av. Claude Guillemin BP6009 45060 Orléans cedex 2, France (p.blanc@brgm.fr)
2. CNRS-IC2MP-UMR7285 , HydrASA, 5 Ave Albert Turpain, F-86022 Poitiers Cedex, France
3. ANDRA – Parc de la Croix Blanche, 1-7 rue Jean Monnet, 92298 Châtenay-Malabry Cedex 2, France

Introduction

The thermodynamic properties of clay minerals, which control the stability of these minerals in solution, are still a matter of debate in spite of recent advances (Gailhanou et al., submitted). This is especially the case for the minerals that may structurally include trace elements and potential radionuclides such like Ni, Cd, Co, Cr, Mn, Pb, ... The usual methods developed in order to predict thermodynamic properties are parameterised using a given set of minerals. For clay minerals, the latter are mainly composed by Si, Al, Fe and Mg, apart from the alkalis elements (Chermak and Rimstidt, 1989), which means that predictions are limited to minerals whose layers are composed by Si, Al, Fe and Mg. At the vicinity of H&ILW disposal cells, the possible interactions between clay rock or engineered barrier and waste degradation products can result in the appearance of clay minerals that may structurally include radionuclides within an irreversible trapping process. This work aims at proposing a method for predicting the thermodynamic properties of such minerals.

Theoretical principle and selection of calibration phases

Vieillard (1994) has developed a methodology of estimation based on the difference of electronegativity by considering three scales of values of the parameter $\Delta_{\text{H}}\text{O}^-(\text{M}_{\text{clay}}^{z+})$ in the three sites of phyllosilicates. We have considered the work of Vieillard (1994) that originally applies to the estimate of ΔH_f^0 and extended it to the estimate of $\text{Cp}(\text{T})$, S^0 and V . Some popular estimate methods (Chermak and Rimstidt, 1989) are based on the hypothesis that the thermodynamic property of a mineral can be obtained by combining the properties of its components. An improvement of this principle had consisted in decomposing minerals into their polyhedral components (Chermak and Rimstidt, 1989). Now, we can write the fictive solution equilibrium with a basic polyhedral component M_xO_y as: $x\text{M}_{\text{aq}}^{z+} + y\text{O}^- = \text{M}_x\text{O}_y$ and assuming the entropy of this fictive reaction is zero, we can define a $\Delta_{\text{S}}\text{O}^-$ parameter as:

$$\Delta_{\text{S}}\text{O}^-(\text{M}_{\text{clay}}^{z+}) = \text{S}^0(\text{O}^-) = \frac{\text{S}^0(\text{M}_x\text{O}_y) - x \cdot \text{S}^0(\text{M}_{\text{aq}}^{z+})}{y}.$$

The value for the oxide analog of the polyhedral unit is obtained by implementing S^0 of the oxide in the $\text{S}^0(\text{M}_x\text{O}_y)$ term. We have also defined, from the same reasoning, similar parameters for heat capacity and volume of the basic polyhedral components:

$$\Delta_{\text{Cp}}\text{O}^-(\text{M}_{\text{clay}}^{z+}) = \frac{\text{Cp}(\text{M}_x\text{O}_y) - x \cdot \text{Cp}(\text{M}_{\text{aq}}^{z+})}{y}; \Delta_{\text{V,Cp}}\text{O}^-(\text{M}_{\text{clay}}^{z+}) = \frac{\text{V}(\text{M}_x\text{O}_y) - x \cdot \text{Cp}(\text{M}_{\text{aq}}^{z+})}{y}.$$

Results and discussion

On Figure 1, we have displayed, for entropy, the correlation obtained between calculated values of $\Delta_s O^{\circ} = M^{Z+}$ for the polyhedral unit and for the oxide analog. A straight line and a second-order function are obtained, for the interlayer and octahedral cations, respectively, with a good correlation coefficient.




Fig. 1 - Development of predictive capacity for entropy estimates

The implementation of the derived semi-empirical, first or second order relations allows to estimate the thermodynamic properties of a clay mineral, MX80 ($Na_{0.409}K_{0.024}Ca_{0.009}(Si_{3.738}Al_{0.262})(Al_{1.598}Mg_{0.214}Fe^{3+}_{0.173}Fe^{2+}_{0.035})O_{10}(OH)_2$) in the present case, loaded by 6 radionuclides and to compare the values with the results obtained by Gailhanou et al. (submitted). The results can be expressed in terms of the concentrations for the elements Ni, Cd, Co, Cr, Mn and Pb and in terms of energetic potential with respect to the measurements performed by Gailhanou et al. (submitted).

References:

- Chermak J.A. and Rimstidt J.D. (1989) Estimating the thermodynamic properties (ΔG°_f and ΔH°_f) of silicate minerals at 298K from the sum of polyhedral contributions. *Am. Miner.*, **74**, 1023-1031.
- Gailhanou, H., Blanc, P., Rogez, J., Mikaelian, G., Kawaji, H., Olives, J., Amouric, M., Denoyel, R.S.B., Montouillout, V., Vieillard, P., Fialips, C. I., Giffaut, E., Michau, N., Gaucher, E.C., (Submitted). Thermodynamic properties of illite IMt-2, smectite MX-80 and beidellite SBld-1 by calorimetric methods: Enthalpies of formation, heat capacities, entropies and Gibbs free energies of formation. *Geochimica et Cosmochimica Acta*.
- Vieillard, P., 1994a. Prediction of enthalpy of formation based on refined crystal structures of multisite compounds.1. Theories and examples. *Geochimica et Cosmochimica Acta* 58, 4049-4063.