



**HAL**  
open science

## Modeling the dioctahedral smectites layer charge variation versus structural Iron reduction level

Jebril Hadi, Christophe Tournassat, Stéphanie Betelu, Ioannis Ignatiadis,  
Laurent Charlet

### ► To cite this version:

Jebril Hadi, Christophe Tournassat, Stéphanie Betelu, Ioannis Ignatiadis, Laurent Charlet. Modeling the dioctahedral smectites layer charge variation versus structural Iron reduction level. International meeting "Clays in Natural and Engineered Barriers for Radioactive Waste Confinement", Oct 2012, Montpellier, France. hal-00703576

**HAL Id: hal-00703576**

**<https://brgm.hal.science/hal-00703576>**

Submitted on 3 Jun 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Modeling the dioctahedral smectites layer charge variation versus structural Iron reduction level

J. Hadi<sup>1,\*</sup>, C. Tournassat<sup>1</sup>, S. Betelu<sup>1</sup>, I. Ignatiadis<sup>1</sup>, L. Charlet<sup>2</sup>

<sup>1</sup>BRGM, Environment and Processes Division, 3 Avenue Claude Guillemin, BP 36009, 45060 Orléans Cedex, France (\*corresponding author: j.hadi@brgm.fr)

<sup>2</sup>ISTERRE, BP 53, 38041 Grenoble, Cedex 9, France

Iron is one of the most common redox species in soils and sedimentary rocks. Amongst iron-bearing phases, phyllosilicates might play key roles in various bio-geochemical processes involving redox reactions, where structural Fe ( $Fe_{str}$ ) can act as a renewable source/trap of electron.

A large set of data from kinetics, spectroscopic or electrochemical studies on dioctahedral smectites demonstrates that reduction of  $Fe_{str}$  impacts many clay properties such as colour, layer charge, swelling pressure, colloidal properties that are linked to layer structural changes. Experiments also suggest that this mechanism is partly reversible, depending on type and properties of the primary oxidized clay, on how the reduction is induced (chemically and/or biologically) and on extent of iron reduction level.

The complexity of the involved mechanisms makes the prediction of  $Fe_{str}$  redox properties challenging. For instance, only empirical models are currently available to quantify structural changes as a function of reduction level. However, a predictive and mechanistic model of these changes is a prerequisite to develop a thermodynamic model for  $Fe_{str}$  redox properties.

In this contribution, we propose a mechanistic statistical model to explain 2:1 layer excess negative charge changes induced by structural Fe(III) to Fe(II) chemical reduction (by dithionite). This model completes this published by Drits and Manceau (2000) and was calibrated on data from our own and from the literature.

Actually, a large number of studies on  $Fe_{str}$  redox properties ( $E_h$  and kinetics) neglects the major structural changes that occur during redox reactions of this material and that are partially reversible, and are focused in measuring a single  $E_h$  value. Actually, the complex relationship that exist between the different structural iron sites should lead to consider that not only one but several Fe(II)/Fe(III) poles (classes) must exist in the structure, thus exhibiting gradually decreasing  $E_h$  values. Hence, further developments of our model will include Crystal Field Theory (CFT) calculation to identify the variety of  $Fe_{str}$  redox potential, which arises from the varying  $Fe_{str}$  neighbouring inside the same structure and along the redox processes.

## Acknowledgement

*The research leading to these results has received funding from i) the European Union's European Atomic Energy Community's (Euratom) FP7/2007-2011 under grant agreement n° 212287 (RECOZY project) and ii) the ANDRA under the BRGM-ANDRA partnership (CAPTANDRA project, No N°045731SOS).*

## References

Drits, V.A. and Manceau, A. (2000) *A model for the mechanism of  $Fe^{3+}$  to  $Fe^{2+}$  reduction in dioctahedral smectites*. Clays and Clay Minerals **48**:185-195.