



HAL
open science

GRACE mass balance estimates for all the Earth's ice-covered regions

Thomas Jacob, J. Wahr, T. W. Pfeffer, S. Swenson

► **To cite this version:**

Thomas Jacob, J. Wahr, T. W. Pfeffer, S. Swenson. GRACE mass balance estimates for all the Earth's ice-covered regions. Grace Science team meeting 2012, Sep 2012, Potsdam, Germany. hal-00721697

HAL Id: hal-00721697

<https://brgm.hal.science/hal-00721697>

Submitted on 30 Jul 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

GRACE mass balance estimates for all the Earth's ice-covered regions

We have used GRACE satellite gravity measurements to construct 2003-2010 mass loss estimates for all the world's mountain glacier systems and ice caps (GIC's), as well as for Greenland and Antarctica. We find that the GIC's, excluding the Greenland and Antarctic peripheral glaciers and ice caps (PGIC's), lost mass at a rate of 148 ± 30 Gt yr⁻¹ from January 2003 to December 2010, contributing 0.41 ± 0.08 mm yr⁻¹ to global sea level rise (SLR) during this period. This rate is about 30% smaller than previously published ground-based estimates that most nearly matched this time span. Our results for the high mountains of Asia, in particular, show a mass loss of only 4 ± 20 Gt yr⁻¹, compared to 47-55 Gt yr⁻¹ in previously published estimates.

We also estimate that the Greenland and Antarctic ice sheets, including their PGIC's, lost 384 ± 71 Gt yr⁻¹, contributing 1.06 ± 0.19 mm yr⁻¹ to SLR, over this same time period. The total SLR contribution from all ice-covered regions was then 1.48 ± 0.26 mm yr⁻¹, agreeing with independent estimates of SLR from new water to within the respective error bars.