



# Structure of hydrothermal convection in the Upper Rhine Graben as inferred from corrected temperature data and basin-scale numerical models

Laurent Guillou-Frottier, Clément Carré, Bernard Bourguin, Vincent Bouchot, Albert Genter

## ► To cite this version:

Laurent Guillou-Frottier, Clément Carré, Bernard Bourguin, Vincent Bouchot, Albert Genter. Structure of hydrothermal convection in the Upper Rhine Graben as inferred from corrected temperature data and basin-scale numerical models. AGU Fall Meeting 2012, Dec 2012, San Francisco, United States. hal-00746045

**HAL Id: hal-00746045**

**<https://brgm.hal.science/hal-00746045>**

Submitted on 26 Nov 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Structure of hydrothermal convection in the Upper Rhine Graben as inferred from corrected temperature data and basin-scale numerical models

L. Guillou-Frottier<sup>1</sup>, C. Carré<sup>1</sup>, B. Bourguine<sup>1</sup>, V. Bouchot<sup>1</sup>, A. Genter<sup>2</sup>

1. BRGM, GeoResources Division, Orléans, France

2. GEIE Exploitation Minière de la Chaleur, Kutzenhausen, France

Geothermal anomalies in sedimentary basins are strongly controlled by fluid circulation within permeable zones. This study presents a new compilation of newly corrected bottom-hole temperature data in the French part of the Upper Rhine Graben, where the Soultz-sous-Forêts temperature anomaly is locked beneath a local horst structure. After a geostatistically constrained interpolation procedure, maps and cross-sections are extracted from the 3D thermal block, together with the associated standard deviations. Thermal anomalies are preferentially associated with the thickest zones of the permeable fractured Buntsandstein (sandstones) formation, in apparent contradiction with previous models where two major fault zones were suggested to control fluid flow. The underlying fractured granitic basement hosts fluid circulation patterns which are apparently controlled at large-scale by the inclined basement-sediments interface.

Based on these observations, numerical models of hydrothermal convection including an inclined basement-sediments interface, a local horst structure, and realistic petrophysical properties have been carried out. The depth-decrease of permeability, the inclination of the interface and the fixed heat flow condition at the base of the model, explain why only a few upwellings can be triggered. Thermal anomalies and a measured temperature profile can be reproduced when fault permeability equals  $10^{-14}$  m<sup>2</sup>. Interestingly, structure of convective patterns also exhibits a second hotter upwelling, 8 km east of the Soultz-sous-Forêts upwelling zone, where another geothermal exploration project is now underway. The understanding of hydrothermal convection with realistic fluid and rock properties clearly appears as a predictive tool for geothermal exploration strategies.

Abstract accepted for 2012 AGU Fall meeting, San Francisco.