



HAL
open science

Ground motion scenario based on multi-scale mapping of fault heterogeneity

Hideo Aochi, Satoshi Ide

► **To cite this version:**

Hideo Aochi, Satoshi Ide. Ground motion scenario based on multi-scale mapping of fault heterogeneity. Joint Assembly IAHS - IAPSO - IASPEI, Jul 2013, Gothenburg, Sweden. hal-00809382

HAL Id: hal-00809382

<https://brgm.hal.science/hal-00809382v1>

Submitted on 9 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Presenting author corresponding details

Family Name: Aochi
First Name: Hideo
Institution/company: BRGM
City: Orléans
Country: France
Email: h.aochi@brgm.fr
Verify E-mail: h.aochi@brgm.fr

Abstract title

Ground motion scenario based on multi-scale mapping of fault heterogeneity

Abstract text

It has been proposed that earthquake complexity can be described by multi-scale heterogeneity in fracture energy of the fault interface (Ide and Aochi, JGR, 2005). The 2011 Tohoku earthquake revealed us such multi-scalability of the fault heterogeneity, seemingly a cascade-rupture mode (Aochi and Ide, EPS, 2011; Ide and Aochi, accepted in Tectonophysics, 2013). We are then interested in the effect on the ground motion through systematic works on the model parameters. This study presents the ground motion simulations based on the multi-scale dynamic rupture simulations in the regional scales for both shallow moderate earthquakes and subduction large earthquakes. In all cases, the synthetic seismograms in time vary according to the variation in dynamic rupture process (rupture directivity, rupture directivity, and slip distribution, source time function). The frequency content changes significantly in the near field, while it is quite homogeneous for a receiver located far away.

Symposium

075 IASPEI: S201c - Earthquake scenarios

Presentation preference

No preference

Grant application**Author details**

Aochi, H., BRGM, France (Presenting); Ide, S., University of Tokyo, Graduate School of Science, Japan