



Pressure transient tests on geothermal wells in the Dogger aquifer, Paris Basin

Elodie Giuglaris, Virginie Hamm

► To cite this version:

Elodie Giuglaris, Virginie Hamm. Pressure transient tests on geothermal wells in the Dogger aquifer, Paris Basin. EGU General Assembly 2014, Apr 2014, Wiens, Austria. hal-00944880

HAL Id: hal-00944880

<https://brgm.hal.science/hal-00944880>

Submitted on 19 Apr 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Pressure transient tests on geothermal wells in the Dogger aquifer, Paris Basin

Giuglaris Elodie, Hamm Virginie ¹

¹ BRGM, 3 Avenue Claude Guillemin, BP 36009 - 45060 Orléans Cedex 2, France

In this work, we propose a new interpretation of pressure transient tests conducted in more than sixty wells targeting the Dogger formation in the Paris basin. The most part of these wells were drilled during the eighties over a very short period of time in an urban area. The transient tests were only recorded during a pressure buildup and show that contrary to what has long been assumed the Dogger aquifer behavior is not homogeneous.

The interpretation of the obtained data was mainly done with Horner's method, independently for each well, and assuming generally that the aquifer is homogeneous. Available data from pressure transient tests are limited because of the precision of the tools used, and because pressure buildup was only recorded during eight to twelve hours. The objective was to determine the average transmissivity and the skin factor in order to obtain the productivity of the wells.

We gather all available data, including those obtained on the thirteen recent wells, and propose a new interpretation in terms of aquifer model. We use the pressure derivative method developed in the nineties for oil reservoirs which allows a better visualization of the model and of the heterogeneity of the aquifer.

Most of the well tests have a derivative pressure curve with a slope equal to zero, that does not change after the well bore effects, confirming that the aquifer is relatively homogeneous. However on a dozen of wells, data from pressure transient tests display singular behaviors that can be related to a double porosity or double permeability model for some wells and to a composite model for others (figure 1). These observations question the validity of a homogeneous model for the Dogger aquifer in the Paris basin and the currently used parameters for the thermal and hydrodynamic predictive model of geothermal exploitation.

Finally, this work, supported by the French agency of environment and renewable energy (ADEME), increases our understanding of the Dogger aquifer in the Paris basin and will allow the optimization of the ongoing hydraulic tests and the future exploitation of geothermal energy in the area.




Figure 1 : Pressure and derivative responses during the transient test on the well "Aulnay 2"

