



**HAL**  
open science

## Multi-isotopes constraints on the origins and processes of groundwater salinization in coastal aquifers. Example of Recife, Northeast of Brazil

Lise Cary, Emmanuelle Petelet-Giraud, G Bertrand, Wolfram Kloppmann, Luc Aquilina, Helene Pauwels, Veridiana Martins, Ricardo Hirata, Suzana Montenegro

### ► To cite this version:

Lise Cary, Emmanuelle Petelet-Giraud, G Bertrand, Wolfram Kloppmann, Luc Aquilina, et al.. Multi-isotopes constraints on the origins and processes of groundwater salinization in coastal aquifers. Example of Recife, Northeast of Brazil. EGU General Assembly 2015, Apr 2015, Vienne, Austria. pp. EGU2015-10320. hal-01103181

**HAL Id: hal-01103181**

**<https://brgm.hal.science/hal-01103181>**

Submitted on 14 Jan 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## **Multi-isotopes constraints on the origins and processes of groundwater salinization in coastal aquifers. Example of Recife, Northeast of Brazil**

Lise Cary (1), Emmanuelle Petelet-Giraud (1), Guillaume Bertrand (2), Wolfram Kloppmann (1), Luc Aquilina (3), Helène Pauwels (1), Veridiana Martins (2), Ricardo Hirata (2), and Suzana Montenegro (4)

(1) BRGM, 3 av. C. Guillemin, BP36009, 45060 Orléans, France (e.petelet@brgm.fr), (2) USP, Instituto de Geociências - Rua do Lago, 562 ; Butantã - 05508-080 Sao Paulo, Brazil, (3) Université Rennes1 - CNRS, OSUR-Géosciences Rennes, campus Beaulieu Av. du Gal Leclerc, F-35000 Rennes, France, (4) UFPE, Civil Engineering Department, 50740 Recife, Brazil

The Recife Metropolitan Region (PE, Brazil) is a typical "hot spot" illustrating the problems of southern countries on water issues inducing high pressures on water resources both on quantity and quality in the context of global social and environmental changes. This study focuses on the groundwater geochemistry in a coastal multilayer aquifer and aims at investigating the sources and processes of salinization. The RMR basement is constituted by two different Precambrian blocks separated by a large lineament area. The sedimentary fillings of the two basins present different origins that can be distinguished by the Sr isotope composition. The northern deep Beberibe aquifer displays very high strontium isotope ratios with a large range of values ( $87\text{Sr}/86\text{Sr} = 0.7102$  to  $0.7233$ ) illustrating the main continental origin of sediments whereas the southern deep Cabo aquifer showed lower  $87\text{Sr}/86\text{Sr}$  values ( $87\text{Sr}/86\text{Sr} = 0.7097$  to  $0.7141$ ) indicating the contribution of the marine sedimentation dating from the Atlantic opening. Although sulfate isotopes, Electric Conductivity and Cl contents indicate a clear mixing with seawater for some samples of the deep Cabo and Beberibe aquifers, all  $87\text{Sr}/86\text{Sr}$  values are above the present-day seawater composition. This can be related to the complex local history of transgression/regression phases that induced alternately salinisation and freshening with gains and losses of cations and Sr, together with water-rock interactions. Stable isotopes of the water molecule clearly evidence the local present day recharge especially within the surficial aquifer, whereas some samples are affected by in situ evaporation processes and/or recharge with evaporated water originating from dam used for water supply. The two deep aquifers display a high range of B concentrations ( $\sim 20$  to  $600 \mu\text{g.L}^{-1}$ ) and B isotope composition ( $\delta^{11}\text{B} = 6.7$  to  $68.5 \text{‰}$ , with the highest values known to date ( $63$ - $68.5 \text{‰}$ ). This suggests multiple sources and processes affecting B behavior, among which mixing with saline water, B sorption on clays/organic matter and mixing with wastewater. Indeed, the Tertiary and Quaternary surficial aquifers are highly contaminated with wastewater and locally salinized; here salinisation is possibly due to present seawater intrusion.