

Impact of pesticides on denitrification in groundwaters

Caroline Michel, Nicole Baran, Laurent André, Catherine Joulian

▶ To cite this version:

Caroline Michel, Nicole Baran, Laurent André, Catherine Joulian. Impact of pesticides on denitrification in groundwaters. 1st International Conference on Microbial Ecotoxicology. Ecotoxicomic 2017, Nov 2017, Lyon, France. 2017. hal-01586077

HAL Id: hal-01586077 https://brgm.hal.science/hal-01586077v1

Submitted on 12 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Impact of pesticides on denitrification in groundwaters

Caroline Michel*¹, Nicole Baran¹, Laurent Andre¹, and Catherine Joulian¹

¹BRGM – Bureau de Recherches Géologiques et Minières (BRGM) – 3 avenue Claude Guillemin, B.P. 36009, Orléans cedex 2, France

Abstract

Denitrification processes are involved in the decrease of nitrate (NO3-) concentrations in groundwaters impacted by human activities. Non-target/side effects of pesticides on soil microbial denitrification have been demonstrated. It is however now well established that groundwater is an ecosystem in itself with specific microbial communities different from soils' ones. If the impact of pesticides on groundwater bacterial communities has already been studed, the impact of pesticides on the denitrification activity in groundwater has still to be investigated. Groundwater samples are collected from several wells on a catchment submitted to agricultural practices with fertilizers and pesticides uses. Groundwaters contain nitrate (lower than limit of quantification up to 80 mg/l), pesticides and some of their metabolites in low concentrations. The abundance of genes involved in denitrification (narG, napA, nirS) shows the occurrence of bacteria able to perform, at least, the two first steps of denitrification. Three groundwaters distributed within the catchment and exhibiting different denitrification signatures based on genes identification and physic-chemical properties are selected. They are used in laboratory to perform batch experiments in conditions boosting microbial denitrification (addition of KNO3 and acetate, anaerobic conditions, 25°C). The microbial ecotoxicology study is conducted in the presence or absence of pesticides spiked as single or cocktail pesticides. Several parameters (pH, concentrations of acetate, NO3, NO2, N2O, biomass) are recorded over time to quantify denitrification rate according to experimental conditions. This experimental work constitutes a first step in the understanding of the inhibitor/catalyzer role of pesticides on denitrification process.

Keywords: denitrification, groundwater microbial community, pesticides, side effects

^{*}Speaker