



HAL
open science

Understanding b-value of Gutenberg-Richter relation from dynamic rupture simulations

Hideo Aochi

► **To cite this version:**

Hideo Aochi. Understanding b-value of Gutenberg-Richter relation from dynamic rupture simulations. EGU General Assembly 2019, Apr 2019, Viennes, Austria. hal-01972914

HAL Id: hal-01972914

<https://brgm.hal.science/hal-01972914>

Submitted on 8 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Understanding b -value of Gutenberg-Richter relation from dynamic rupture simulations

Hideo Aochi (1,2)

(1) BRGM, Orleans, France (h.aochi@brgm.fr), (2) Laboratoire de Géologie, Ecole Normale Supérieure/CNRS UMR 8538, PSL Research University, Paris, France (aochi@biotite.ens.fr)

It is requested to understand the mechanism of b -value in Gutenberg-Richter relation of seismicity to improve the probabilistic seismic hazard study. Therefore, we study a simple fault system described by multi-scale heterogeneity (Ide and Aochi, JGR, 2005) and carry out dynamic rupture simulations of earthquakes subject to different stress accumulations, namely a uniformly, constantly loading (natural seismicity) and a localized stress propagating from a point (induced seismicity). We also test different distributions of the heterogeneity (its density, scaling relation of heterogeneity, etc.). The variation of b -value has been briefly reported previously (e.g. Aochi and Ide, JGR, 2009; Rohmer and Aochi, GJI, 2015), but not systematically studied yet. From our simulations, we observe that the b -value increases when the heterogeneity is sparse or when the stress accumulation is spatially heterogeneous. This is comprehensible, as a small rupture initiation difficultly extends to large scale, and this can be a key in the difference of natural and induced seismicity. The system also provides a very wide range of b -values, larger than 2.