



**HAL**  
open science

# Autochthonous greenstone belts of Madagascar : Implication for the archean tectonics in the frame of the greater darwhar craton

Jean-Yves Roig, Delphine Bosch, Claude Delor

## ► To cite this version:

Jean-Yves Roig, Delphine Bosch, Claude Delor. Autochthonous greenstone belts of Madagascar : Implication for the archean tectonics in the frame of the greater darwhar craton. IGCP 6148 SUPERCONTINENT CYCLES & GLOBAL GEODYNAMICS, Jun 2019, Antananarivo, Madagascar. hal-02080459

**HAL Id: hal-02080459**

**<https://brgm.hal.science/hal-02080459>**

Submitted on 26 Mar 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# AUTOCHTHONOUS GREENSTONE BELTS OF MADAGASCAR: IMPLICATION FOR THE ARCHEAN TECTONICS IN THE FRAME OF THE GREATER DARWHAR CRATON

Roig J.Y.<sup>1</sup>, Bosch D<sup>2</sup> and Delor C<sup>1</sup>

1-Bureau des Recherches Géologiques et Minières, 3 avenue C. Guillemin, BP 6009, 45060 Orléans, Cedex 2, FRANCE

2- Géosciences Montpellier, UMR 5243 Université Montpellier 2, Place E. Bataillon 34095 Montpellier cedex 5

The new geochronologic, petrologic and structural data acquired during the mapping project of the Programme de Gouvernance des Ressources Minérales de Madagascar, completed by academic works, lead to the re-definition of eight geological domains which make up the metamorphic Malagasy basement. The Center of the Island only consists of the Antananarivo and Antongil/Masora domain respectively. The Paleo-Mesoarchean Antongil/Masora domain (3.32-3.15 Ga) is interpreted as a fragment of the wider Dharwar craton of India. Its adjacent Neoproterozoic Antananarivo domain (2.7-2.5 Ga) is well known because it comprises a critical component of mafic-ultramafic gneisses which crop out as five N-striking belts interpreted as Archaean metamorphosed greenstone belts referred hereafter as “Tsaratana Complex”.

In the conventional view, the Tsaratana complex is interpreted as a single allochthonous unit thrust over granitoid gneisses of the Antananarivo block during the Panafrican orogeny. Two aspects of this notion are controversial: (i) the age of the allochthon's emplacement and (ii) the very nature of the allochthon itself

We present new structural data and U-Pb isotopic ages arguing that neither the structural evolution nor the age constraints of the identified geological events are consistent with an allochthonous model of nappe tectonics. On the contrary, the “Tsaratana Complex” is likely an autochthonous domain or was juxtaposed with the Antananarivo Domain in the late Archaean / Early Palaeoproterozoic times. In this way, the Antananarivo Domain and the associated “Tsaratana Complex” does not differ from the Eastern Dharwar Craton. Consequently, the reliability of the “Betsimisaraka suture” is discussed and we propose that Neoproterozoic rocks of the Eastern Dharwar Craton and the Antananarivo domain were symmetrically disposed around the Mesoarchean nucleus formed by the Western Dharwar Craton and the Antongil/masora Domain as early as Neoproterozoic times.