HAL
open science

Data analysis of pXRF measurements on soil samples from the Brouzils Sb deposit (France)

Bruno Lemiere, Jérémie Melleton, Pascal Auger

To cite this version:

Bruno Lemiere, Jérémie Melleton, Pascal Auger. Data analysis of pXRF measurements on soil samples from the Brouzils Sb deposit (France). 2nd International Conference "Mines of the Future", Jun 2019, Aachen, Germany. hal-02121075

HAL Id: hal-02121075

https://brgm.hal.science/hal-02121075v1

Submitted on 6 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Data analysis of pXRF measurements on soil samples from the Brouzils Sb deposit (France)

Bruno LEMIERE, Jeremie MELLETON, Pascal AUGER (BRGM, 45060 ORLEANS, FRANCE)

Aachen International Mining Symposia 2019

The UpDeep project aims at developing low footprint strategies, in order to facilitate exploration programs in either populated areas or environmentally sensitive areas, especially in Europe where the social acceptance of mining is particularly low. The impact of weak leach soil and plants analysis is much smaller than classical drilling or trenching programs. Dynamic sampling and on-site decision making on survey orientation using pXRF soil surveys achieve also the objective of low footprint and minimise sample collection. Field work was performed both in Nordic countries and in France, the present study at the Les Brouzils antimony deposit, a critical element.

The Brouzils vein field occurs in was detected by bedrock Sb geochemistry and electromagnetic survey. This site was briefly mined and is considered as a potential Sb resource.

For UpDeep, 146 soil samples were analysed by pXRF. They belong to Ah and B soil horizons at five Sb prospects, spread on six profiles. Results were processed as independent data sets in order to identify independently geochemical trends and spatial patterns. pXRF data sets are usually consistent with quasi-linear relationships with laboratory analyses. The most significant variables and their relations (Sb-As-Mn association) were deduced from a general PCA and CA factor analysis.

Data were then converted to centered log ratios. This separates clearly one lithology-related group (Rb, K, Ti, Sr, Th, Zr) and a mineralisation-related group (Sb, As) which includes Mn and Fe in heat map format. A second group (S, Pb, Ba) seems to be unrelated with (Sb, As, Mn) and closer to the lithology group. Similar but slightly different element associations and enrichment factors were observed on Ah and B data.

Spatial Sb and As anomalies on profiles are well resolved, located closeby the known position of vein occurrence and of mine workings, with little or no anomalies recorded in the host formations. Mn anomalies are slightly broader.

High density pXRF measurements can be successfully used for precise location of anomalies and for optimisation of laboratory analyses.