



Impact of chalk karst on river flow in Normandy : comparison of the behavior of three rivers and quantification of karst flow, France

Pierre-Yann David, Didier Pennequin, Hélène Bessiere

► To cite this version:

Pierre-Yann David, Didier Pennequin, Hélène Bessiere. Impact of chalk karst on river flow in Normandy : comparison of the behavior of three rivers and quantification of karst flow, France. 46th IAH Congress Malaga - IAH 2019, Sep 2019, Malaga, Spain. hal-02137033

HAL Id: hal-02137033

<https://brgm.hal.science/hal-02137033>

Submitted on 22 May 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Impact of chalk karst on river flow in Normandy : comparison of the behavior of three rivers and quantification of karst flow, France. #603

Withdraw

Accepted

Didier Pennequin submitted this abstract and it was finally accepted for track

Topic 7 - Karst Hydrogeology

as Oral.

[Go to contribution](#)

Author:

Pierre-Yann David

Co-authors:

Didier Pennequin, H  l  ne Bessiere

In 1758, Jean-Etienne Guettard mentioned in a publication of the French Royal Academy of Sciences the particular behavior of three rivers in Normandy that "gradually disappear and then reappear downstream". Guettard noticed "holes in the river bed that absorb river flow". These three rivers are respectfully the Iton, the Avre and the Risle rivers, all flowing on the Upper Cretaceous chalk, and located in the eastern part of Normandy, in northwest France. The Risle and Avre rivers are both constituted of 3 segments showing changing hydraulic behaviors: a first section upstream where the river and the groundwater (GW) are hydraulically connected and are in equilibrium, a middle portion where the river and the GW are disconnected, with sinkholes forming in the minor bed, progressively decreasing the river flowrate and sometimes leading to complete river dry up, and a third section downstream, representing converging outlets of the karst aquifer system, where river and GW undergo reconnection, starting up with a strong resurgence area. The Iton river is a more complex hydrosystem with two successive karst sections resulting in five segments showing changing hydraulic behaviors (river and WT level being alternatively connected and disconnected). Several monitoring networks for groundwater level and flowrate do exist in these rivers since the seventies or eighties. A more dense and better located monitoring network was implemented in the Risle river basin as part of the Risle Observatory which operated over a 3 year period (2013-2016). These monitoring networks allowed for using lumped hydrological modelling with GARDENIA to notably quantify groundwater karstic flowrate as well as the groundwater components of the river flow. Eleven (11) lumped modelling processes were carried out with calibration using both piezometric levels and riverflow : 2 models were generated in the Iton basin, 4 in the Risle basin and 3 in the Avre basin. Two (2) other models were also elaborated in less karstic chalk basins (Austreberthe and Cailly Rivers) for comparison.

Regarding the groundwater component of the river flow, results obtained showed that for less karstic chalk basins, the average net GW input accounts for 74% to 98% of the mean river flow. In karstic basins, the net groundwater component of the mean river flow can decrease to around 30% at the end of the karst sections(i.e; 34 % for the Risle and 29% for the Avre). In the karst sections of the rivers, GW karst flow can be highly significant, up to 36% of the effective rainfall in the second karst segment of the Iton river and 56% in the middle portion of the Risle river). Acquiring a good understanding of these GW and river flow interaction processes is essential for floodrisk assessment in karstic chalk environments.

D

Didier Pennequin submitted this abstract 29 Apr 2019

As contribution type: Oral

For tracks:

Topic 5 - Tools, methods and models to study groundwater

Topic 7 - Karst Hydrogeology

Author:

Co-authors:

- Pierre-Yann David (BRGM)

- Didier Pennequin (BRGM)
- H  l  ne Bessiere (BRGM)

Speaker:

- Pierre-Yann David (BRGM)

J

This abstract was accepted 20 May 2019
The abstract was accepted as Oral for track
Topic 7 - Karst Hydrogeology
and a [contribution](#) was created in the event.