



HAL
open science

Statistical analysis of microseismicity during the stimulations at Soultz-sous-Forêts (France) EGS site

Hideo Aochi, Sabine Ben Rhouma, Thomas Le Guenan, Julie Maury

► To cite this version:

Hideo Aochi, Sabine Ben Rhouma, Thomas Le Guenan, Julie Maury. Statistical analysis of microseismicity during the stimulations at Soultz-sous-Forêts (France) EGS site. DEEPEGS Symposium 2019, Nov 2019, Orléans, France. hal-02290848

HAL Id: hal-02290848

<https://brgm.hal.science/hal-02290848>

Submitted on 18 Sep 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

For Symposium DEEPEGS (7-8 November 2019). Poster preferred.

Statistical analysis of microseismicity during the stimulations at Soultz-sous-Forêts (France) EGS site.

H. Aochi, S. Ben Rhouma, T. Le Guenan, J. Maury (BRGM)

Monitoring the microseismicity according to the fluid injection is necessary for assessing better the reservoir behavior at Enhanced Geothermal System or other subsurface exploitation. Mechanical modeling is helpful, but it is still difficult to consider all the aspects due to the limits of numerical frameworks and poor knowledge of certain model parameters. We then alternatively aim to analyze the statistically microseismicity to follow the temporal evolution linked to the injection operation. Earthquakes occur due to any loading forces (regional stress accumulation and/or pore pressure increase). On the other hand, once an earthquake occurs, it redistributes the stress in the surrounding and leads to other earthquakes (called 'aftershock'). We need to distinguish them. In seismology, statistical models have been developed and successfully applied in various seismicity in the world, including Epidemic-Type Aftershock Sequence (ETAS) model (e.g. Ogata, Ann. Inst. Statist. Math, 50:379-402, 1998). We apply the ETAS model on the microseismicity for the 1993 and 2000 stimulation experiments at Soultz-sous-Forêts EGS site (Alsace, France). Thousands earthquakes in the published catalogue (Data Center for Deep Geothermal Energy; <https://cdgp.u-strasbg.fr/>) allow us to analyze the temporal change of the seismicity rate. Among the five model parameters in ETAS model, some parameters are too sensitive to decide reliably, however our analyses show an important launch of the seismicity rate at the beginning and a gradual decrease with time. This is qualitatively consistent with the observations.