



# Micropollutants and boron isotopes as tracers of domestic wastewater effluents in surface and groundwaters

Emmanuelle Petelet-Giraud, Nicole Baran, Coralie Soulier

## ► To cite this version:

Emmanuelle Petelet-Giraud, Nicole Baran, Coralie Soulier. Micropollutants and boron isotopes as tracers of domestic wastewater effluents in surface and groundwaters. 2020. hal-02426439

**HAL Id: hal-02426439**

**<https://brgm.hal.science/hal-02426439>**

Preprint submitted on 2 Jan 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# 36TH INTERNATIONAL GEOLOGICAL CONGRESS

2 - 8 MARCH 2020, DELHI, INDIA

## Micropollutants and boron isotopes as tracers of domestic wastewater effluents in surface and groundwaters

PETELET-GIRAUD, E.<sup>1</sup>, Nicole BARAN<sup>1</sup>, Coralie SOULIER<sup>1</sup>

1. BRGM, Orléans, France, e.petelet@brgm.fr

Surface water and groundwater contamination by a chemical element or molecule may have several origins. For instance, AMPA can originate from agricultural practices as a degradation product of glyphosate, but also from wastewater effluents as a degradation product of phosphonates. The discrimination of the origins and vectors of contamination in the environment is both an environmental and a societal issue, and is required to define integrated water resources management at the catchment or water body scale by implementing appropriate measures to effectively struggle against pollution. The objective of this study is to define a methodology to identify a "domestic wastewater" component within surface waters and groundwater. As there is no ideal tracer, i.e. conservative, persistent in water compartments, present in quantity above the detection limit and originating from a single type of pollution source, we propose a multi-tracer approach (chemical and isotopic) to identify and validate the relevance of foreseen tracers, in order to overcome this limitation. Among the relevant tracers of wastewater, some may be used for their intrinsic or combined discriminant power: i.e. Tracer of detergents: boron concentration and isotopes; Pharmaceuticals tracers: e.g. carbamazepine, ibuprofen, gadolinium anomaly; Life-style tracers: e.g. caffeine, sweeteners.

The studied catchment is impacted both by diffuse agricultural pollution and punctual wastewater inputs from small capacities wastewater treatment plants without tertiary treatment process, representative of rural areas. Potential combination of suitable tracers is discussed combining various micropollutants and isotopic signatures.