



Modeling the thermodynamic properties of CO₂ aqueous solution using a combination of HKF and Pitzer equations

Marc Parmentier, Adeline Lach, Laurent André, Arnault Lassin

► To cite this version:

Marc Parmentier, Adeline Lach, Laurent André, Arnault Lassin. Modeling the thermodynamic properties of CO₂ aqueous solution using a combination of HKF and Pitzer equations. 43rd IAH Congress, Sep 2016, Montpellier, France. . hal-02457858

HAL Id: hal-02457858

<https://brgm.hal.science/hal-02457858>

Submitted on 28 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Abstract n°2017

1 Although the CO₂-H₂O geochemical system has been extensively studied, it is still difficult to find a model thermodynamically-consistent, able to explain with equivalent confidence the experimental data of density, heat capacity and reciprocal solubility in the conditions range of CO₂ sequestration.

2 Theoretical background

CO₂-H₂O system was modeled using a gamma-phi approach for the liquid-gas equilibrium

$$m_{CO_2} \gamma_{CO_2} K_{CO_2}^0 \exp\left(\frac{\bar{V}_{CO_2}(P - P^0)}{RT}\right) = y_{CO_2} \varphi_{CO_2} P$$

$$a_{H_2O} K_{H_2O}^0 \exp\left(\frac{\bar{V}_{H_2O}(P - P^0)}{RT}\right) = y_{H_2O} \varphi_{H_2O} P$$

With

γ : activity coefficient of aqueous CO₂ calculated with the Pitzer formalism

ψ : fugacity coefficient calculated with the Peng-Robinson equation of state

K: equilibrium constant at water saturation P⁰

V: partial molar volume average over P range

In the aqueous phase, HKF [1] and Pitzer [2] equations are implemented in PhreeSCALE [3] that allows to calculate standard, partial and apparent properties.

$$\frac{G^{ex}}{w_w RT} = f(I) + \sum_i \sum_j \lambda_{ij} m_i m_j$$

$$\frac{\partial G^{ex}}{\partial w_w} \quad \frac{\partial G^{ex}}{\partial n_i} \quad \frac{\partial^2 G^{ex}}{\partial T^2} \quad \frac{\partial G^{ex}}{\partial P}$$

$$\Phi - 1 = \dots$$

$$\gamma_i = \dots$$


$$C_{p,\Phi} = C_p^0 + \dots$$

$$V_\Phi = V_0 + \dots$$

3 Compilation of experimental data

TP range relevant for CO₂ sequestration:
Hydrate <T<200°C, 0<P<80MPa.

- Apparent molar volume V_Φ [4-6] (density)
- Apparent molar heat capacity Cp_Φ [4,7]
- Reciprocal solubility [large compilation]
m_{CO₂} in aqueous phase
γ_{H₂O} in CO₂-rich phase



4 Methodology

- Comparison of various sets of HKF parameters [11-13]

$$V_{CO_2}^0 = a_1 + \frac{a_2}{\psi + P} + \left(\frac{1}{T - \theta} \right) \left(a_3 + \frac{a_4}{\psi + P} \right) - \omega Q$$







$$Cp_{CO_2}^0 = c_1 + \frac{c_2}{(T - \theta)^2} - \left(\frac{2T}{(T - \theta)^3} \right) \left(a_3(P - Pr) + a_4 \ln \left(\frac{\psi + P}{\psi + Pr} \right) \right) + \omega TX$$

Species-specific parameters for HKF equations (units J.mol⁻¹[.K][.bar⁻¹])

	10.a ₁	10 ⁻² .a ₂	a ₃	10 ⁻⁴ .a ₄	c ₁	10 ⁻⁴ .c ₂	10 ⁻⁶ .ω
[11]	26.14	31.26	11.77	-12.92	167.50	36.82	-0.08
[12]	26.14	31.26	11.77	-12.92	160.89	22.40	-0.81
[13]	64	-42	-232	31	154	15	-1.31
This study	31.01	1.71	8.77	-5.12	60.28	61.04	-6.79

- Comparison with others formalisms when possible [9-10]
- New optimization of HKF parameters (with PEST® software) on the data compilation
- Test the use of a T-dependent λ_{CO₂-CO₂} interaction parameter.
 - γ_{CO₂} become different to unity
 - Cp_Φ become different to standard molar Cp

5 Preliminary results



6 Conclusions

No published set of HKF parameters was found to be able to correctly reproduce both V_m and Cp of aqueous CO₂. When using a T-dependent λ_{CO₂-CO₂}, the fit was improved. However further works are needed to avoid unrealistic extrapolation at higher temperatures.

- [1] Tanger, 1988 ; [2] Pitzer, 1973 ; [3] Lach, 2016 ; [4] Barbero, 1983 ; [5] Hnedkovsky, 1996 ; [6] McBride, 2015 ; [7] Hnedkovski, 1997 ; [8] Spycher, 2003 ; [9] Appelo, 2013; [10] Appelo, 2014 ; [11] Shock, 1989 ; [12] Shulte, 2001 ; [13] Plyasunov, 2001 ; [14] Drummond, 1981 ; [15] IAPWS-97