



Liassic age of an oceanic gabbro of the External Rif (Morocco): Implications for the Jurassic continent–ocean boundary of Northwest Africa

André Michard, Abdelkader Mokhtari, Philippe Lach, Philippe Rossi, Ahmed Chalouan, Omar Saddiqi, Ech-Cherki Rjimati

► To cite this version:

André Michard, Abdelkader Mokhtari, Philippe Lach, Philippe Rossi, Ahmed Chalouan, et al.. Liassic age of an oceanic gabbro of the External Rif (Morocco): Implications for the Jurassic continent–ocean boundary of Northwest Africa. *Comptes Rendus Géoscience*, 2018, 350 (6), pp.299-309. 10.1016/j.crte.2018.05.006 . hal-02860803

HAL Id: hal-02860803

<https://brgm.hal.science/hal-02860803>

Submitted on 6 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Tectonics, Tectonophysics

Liassic age of an oceanic gabbro of the External Rif (Morocco): Implications for the Jurassic continent–ocean boundary of Northwest Africa



André Michard ^{a,*}, Abdelkader Mokhtari ^b, Philippe Lach ^c, Philippe Rossi ^d, Ahmed Chalouan ^e, Omar Saddiqi ^f, Ech-Cherki Rjimati ^g

^a 10, rue des Jeûneurs, 75002 Paris, France

^b Université Moulay-Ismaïl, Faculté des sciences, Département de géologie, BP11201, Beni M'Hamed, Meknès, Morocco

^c BRGM, Laboratoire de géochronologie, BP 36009, 45060 Orléans cedex 2, France

^d CCGM, 77, rue Claude-Bernard, 75005 Paris, France

^e Université Mohammed-V, Faculté des sciences, Département des sciences de la Terre, avenue Ibn-Batouta, BP. 1014, Rabat, Morocco

^f Laboratoire Géosciences, Université Hassan-II-Casablanca, BP 5366 Madrif, Casablanca, Morocco

^g Ministère de l'Énergie et des Mines, Direction de la géologie, Rabat Institutes, BP 6208, Rabat, Morocco

ARTICLE INFO

Article history:

Received 22 March 2018

Accepted after revision 26 May 2018

Available online 16 August 2018

Handled by Isabelle Manighetti

Keywords:

Central Atlantic

Alpine Tethys

Transform zone

West African continental margin

Gibraltar Arc

ABSTRACT

This work concerns the northernmost limit of the West African Craton (WAC) and Variscan WAC-related terranes of NW Africa. Based on newly obtained radiometric age of an oceanic gabbro from the “Mesorif Suture Zone” of the External Rif Belt, we propose a revised interpretation of this puzzling lineament. We report on a 190 ± 2 Ma LA-ICP-MS U-Pb zircon age of a trondhjemite vein cross-cutting the Bou Adel gabbro, which is one of the largest oceanic units of the quoted suture zone. We previously interpreted the arcuate MSZ in terms of transported, hyper-extended margin of the Alpine Tethys, based on a K-Ar 166 ± 3 Ma age ascribed to the Bou Adel gabbro in the literature. The new, Early/Middle Liassic age coincides instead with the onset of oceanic floor formation in the Central Atlantic. We hypothesize that the Mesorif suture zone corresponds to the transported trace of the West African Atlantic margin surrounding the northwestern Moroccan Meseta promontory and connecting with the ENE-trending North African Transform/North African transform. The latter zone sharply bounded the North Africa margin and connected the Central Atlantic with the Alpine Tethys. We propose that transported elements from the North African transform constitute the “Mesorif Basalt–Brecias” lineament parallel to and more external than the Mesorif suture zone. If correct, this new interpretation provides an opportunity to develop detailed field and laboratory studies of an exhumed segment of the up-to-now conceptual Jurassic North African transform.

© 2018 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The 2-Ga-old West African Craton (WAC) is surrounded by deformed margins of different types and ages (Fig. 1).

Limiting our study to the northern half of the craton, we find to the east the Pan-African Trans-Saharan collisional belt, formed around 620–600 Ma (Bosch et al., 2016; Caby, 2003). To the west, the Pan-African units are incorporated in the Variscan Mauritanide Belt, built around 300 Ma during the Pangean collision (Bea et al., 2016; Gärtner et al., 2014; Le Goff et al., 2001; Montero et al., 2017;

* Corresponding author.

E-mail address: andremichard@orange.fr (A. Michard).




Fig. 1. The northwestern limits of WAC and WAC-related Variscan terranes. A. Atlantic margin geomorphology and salt basins, after Biari et al. (2017), modified. Salt basin from Tarfaya to Dakhla, after Davison and Daily (2010). Main geological boundaries around the WAC after Michard et al. (2010). B. Geotectonic map of the northwestern Africa–Iberia contact zone, showing the relationships of the Atlantic margin with the Azores–Gibraltar–West Mediterranean transform zone. Background map after Mascle and Mascle (2012). Triassic evaporite basins in pink. Offshore Triassic basin after Hafid et al. (2008).

Villeneuve et al., 2006) and extending northward up to latitude 28°N. This poly-orogenic border has been partly superimposed by the structures of the West African passive margin, formed basically during the Triassic–Liassic breakup of Pangea at ca. 190 Ma (Labails et al., 2010).

The northern border of the WAC is currently defined by the South Atlas Fault (SAF, Fig. 1), i.e. the southern limit of the High Atlas intracontinental Alpine belt at some distance north of the Pan-African suture zone of the central Anti-Atlas (Ennih and Liégeois, 2001, 2008; Frizon de Lamotte et al., 2008). However, the SAF is mostly inherited from the Variscan South Meseta Fault (SMF; Michard et al., 2010), which limits the Meseta orogen from its Anti-Atlas foreland. A Paleoproterozoic and Neoproterozoic crust of Gondwanan affinity has been recently recognized in the Meseta domain (El Houicha et al., 2018; Letsch et al., 2017; Ouabid et al., 2017; Pereira et al., 2015). The Meseta Paleoproterozoic crust was rifted off the WAC before colliding against it during the Neoproterozoic “WAC-Cadomian” event of the Pan-African cycle (Hefferan et al., 2014). During the Paleozoic, the Meseta domain formed the distal passive margin of northwestern Gondwana before being amalgamated along the WAC Anti-Atlas border during the Variscan collision (Hoepffner et al., 2017; Michard et al., 2010). Thus, the most distal margin of the WAC-related terranes corresponds to the northern limit of the Atlas–Meseta domain beneath the thrust units of the ENE-trending Rif–Tell Alpine orogen.

The present paper aims at defining the east-trending limit of northwestern Africa inside the Rif domain and to

recognize its connection with the northeast-trending Atlantic margin. The Moroccan Atlantic margin is currently described as preserving its overall northeast trend up to latitude 35°N (Biari et al., 2017; Hafid et al., 2008; Klingelhoefer et al., 2016), where it disappears beneath the accretionary prism of the Gulf of Cadiz (Crutchley et al., 2011; Gutscher et al., 2009). The implicit suggestion is that the Moroccan Atlantic margin should be crosscut there by the Newfoundland–Azores–Gibraltar fracture zone (Olivet et al., 1984; Sallarès et al., 2011), which broadens and coincides farther to the east with the Tell–Rif orogen (Galindo-Zaldívar et al., 2003; Meghraoui and Pondrelli, 2012). In contrast, we argue in the following that the Liassic Atlantic margin curved eastward at about latitude 35°N, entered the future Rif domain and connected there with the ENE-striking North African Transform (NAT) fault south of the Alpine Tethys domain (Frizon de Lamotte et al., 2011; Lemoine et al., 1987). This argument is based on a new, robust U–Pb zircon age obtained from an oceanic gabbro massif that crops out in the External Rif belt. This massif belongs to the so-called “Mesorif Suture Zone” (MSZ; Fig. 2; (Benzaggagh et al., 2014; Michard et al., 2007, 2014)). Based on the new radiometric date, we propose the working hypothesis that rock material from the Atlantic margin could in fact constitutes most of, if not all, the oceanic elements of the Mesorif suture tectonic lineament. Moreover, we emphasize that the Kimmeridgian–Berriasian tholeiites and carbonates breccias described in the Mesorif (Ben Yaïch et al., 1989; Benzaggagh, 2011; Michard et al., 2007, 2014) define a second lineament,




Fig. 2. A. Location of the Rif mountain belt within the Maghrebide Belt (Durand-Delga, 1980). B. Simplified geological map of the Rif Belt after Suter (1980b) and Benzaggagh et al. (2014), modified. Miocene directions of transport after Frizon de Lamotte (1987), Chalouan et al. (2006, 2014), and Poujol et al. (2017). MBB: Mesorif Basalts-Breccias lineament; MSZ: Mesorif Suture Zone.

here labeled the “Mesorif–Basalts–Breccias” lineament, which could represent a segment of the Jurassic NAT. Therefore, transported elements of the Central Atlantic–Alpine Tethys connection zone would be exposed in the External Rif domain.

2. Geological setting

The Rif mountain belt extends from north to south over a maximum width of ~150 km between the Mediterranean coast and the Atlas–Meseta domain, which forms its

autochthonous foreland (Fig. 2; Chalouan et al., 2008, and references therein). The Rif Mountains belong to the Maghrebide belt, which extends to the east up to the Algerian Kabylies (Kabylias), Peloritani Mountains (NE Sicily), and Calabria (Durand-Delga, 1980). Every Maghrebide segment and the Rif as such consist of imbricated nappes with metamorphic Internal Zones forming the backstop of the tectonic prism. The Rif Belt shares its Internal Zones with the Betic Cordilleras of southern Spain. These common internal zones are labeled the Alboran Domain, which is thought to have been rifted off eastern Iberia together with the other Maghrebides internal terranes ("AlKaPeCa"; see Fig. 5 below) during the Late Eocene–Oligocene northwestward subduction of the Alpine Tethys lithosphere (e.g., Frizon de Lamotte et al., 2000, 2011; van Hinsbergen et al., 2014, and references therein). The Alboran Domain units (from bottom to top, the Sebtides, Ghomarides, and Dorsale calcaire units) are thrust over the Maghrebian Flyschs nappes, which consist dominantly of Lower Cretaceous-Early Miocene turbiditic, deep-water deposits (El Talibi et al., 2014; Guerrera and Martín-Martín, 2014; Zaghloul et al., 2007). The Maghrebian Flyschs have been interpreted as the detached infilling of the Liguro-Maghrebian Tethys (Durand-Delga et al., 2000, and references therein), although they likely onlapped the most distal Dorsale margin during the Middle Jurassic (Chalouan et al., 2011, p. 36; Olivier et al., 1996). They are associated with Upper Jurassic ophiolites and high-pressure/low-temperature metamorphism in Calabria. Ophiolites are also known south of Lesser Kabylia (Petite Kabylie, Algeria) associated with the Flyschs (Bouillin et al., 1997). In the Rif Mountains, only tholeiitic pillow basalts are found as tectonic slivers associated with the Flyschs nappes south of the Bokoya Internal units (Durand-Delga et al., 2000). These basalts are associated with radiolarites and micrites paleontologically dated from the Middle–Late Jurassic. The Flyschs nappes overlie in turn the most internal units of the External prism, namely the Intrarif (Ketama and Tanger units). The root zone of the Maghrebian Flyschs between the Alboran Domain units and the Intrarif is considered to be the Tethyan suture.

The External Zones have been described for long as duplex-type imbricated units (Ketama and Tanger–Loukkos Intrarif units; Mesorif and Prerif units) and gravity nappes detached from the imbricated ones (Ouezzane, Aknoul, Tsoul), altogether derived from the African passive margin (Chalouan et al., 2008; Favre, 1992; Favre et al., 1991; Frizon de Lamotte, 1985; Leblanc, 1979; Suter, 1980a, 1980b; Zaghloul et al., 2005). A balanced cross-section of the tectonic stack of basement and cover units of the External Zone has been ultimately presented by Frizon de Lamotte et al. (2017).

In the eastern Rif, the Intrarif Ketama unit overlies the Beni Malek serpentinite massif, which exhibits a thin calcareous cover including detrital clasts of serpentinite and spinel. This remarkable massif was interpreted as derived from an exhumed mantle segment at the foot of the (hyper-extended) North-African margin (Michard et al., 1992, 2007). The Beni Malek massif and its hidden continuation at shallow depth (Elazzab et al., 1997) are the

easternmost elements of what has been described as the "Mesorif Suture Zone", which extends westward up to the Taounate region of central Rif and includes the Bou Adel gabbro here dated (Benzaggagh et al., 2014; Michard et al., 2014). The Mesorif suture zone consists of a string of kilometer-size slivers of oceanic crustal rocks with thin oceanic sedimentary cover rocks pinched between the Intrarif and Mesorif domains. The Beni Malek massif displays serpentinized spinel lherzolite including occasional pyroxenite layers. The other massifs display varied gabbro facies (cumulate troctolite, ferrogabbro) with E-MORB tholeiitic affinities crosscut by plagiogranite/trondhjemite veins or pockets. Altered basaltic flows and breccias including fragmented variolitic pillow lavas occur at Bou Adel and next to the Beni Malek serpentinites (Skifat metabasites). The sedimentary sequences that overlie these slivers are characterized by gabbroic/basaltic brecias, marbles with ophiolitic clasts, and locally radiolarites (Benzaggagh et al., 2014). We attempted to obtain a paleontological age from radiolarite samples, but the radiolarian fossils were poorly preserved and the age remained indeterminate (M. Chiari, Florence University, Italy, pers. comm.).

3. Dating the Bou Adel gabbro massif

3.1. Sampling and method

The Bou Adel gabbro massif (Fig. 3A) consists of cumulate troctolite and ferrogabbro crosscut in several places by veins, sills or pockets of plagiogranite/trondhjemite (Benzaggagh et al., 2014). Sample BA07 here dated has been collected in a several-meter-thick trondhjemite pocket cropping out along the Oued Azrou stream (Fig. 3B) 300 m to the southeast of the swimming pool of the Bou Adel touristic site (coordinates 34°32'2.85"N, 4°30'12.15"W). In thin section, the sampled rock shows large euhedral plagioclase crystals with minor interstitial quartz; it strictly compares with the trondhjemite illustrated by Benzaggagh et al. (2014) in their fig. 10C.

Zircons were separated from 1 kg of sample using standard heavy liquid and magnetic separation techniques. Seven grains were selected and handpicked to be mounted in an epoxy resin and then polished. Cathodoluminescence (CL) images (Fig. 3C) were obtained by Scanning Electron Microscopy (SEM) to reveal their internal structures. The selected zircon grains are generally broken and range from 50 to 100 µm in size. Only grain #1 is well faceted, prismatic, and pyramid-topped, and displays a core–rim structure.

In situ U-Pb isotope analyses of zircon grains were performed using an ICP-MS ELEMENT XR spectrometer coupled with a CETAC EXCITE 193-nm laser at the GEMOC laboratory facilities at Orsay (University of Paris-Sud). The zircon mount was located in a two-volume cell, with helium as the carrier gas. The laser was operated at a repetition rate of 8 Hz, with an ablation spot size of 40 µm and a fluence of 3.5 J·cm⁻². Calibration for zircon analyses was carried out using the geostandard 91500 zircon (Wiedenbeck et al.,




Fig. 3. A. Location of the dated sample (red star) in the Bou Adel massif (schematic geologic map from Benzaggagh et al., 2014, after the Geological map of Morocco, scale 1:50,000). B. Aspect of the sampled trondhjemite outcrop. C. Cathodoluminescence images of the analyzed zircon grains. D. Concordia age of the BA07 trondhjemite sample in a Wetherill diagram (analyses corrected from common lead).




Fig. 4. Plot of analyses 1 and 2 on grain #1 in the Wetherill diagram. The straight blue line was poorly defined by the two common lead corrected analyses going through close to the origin without being forced. The same analyses without lead correction plot as the two dotted gray ellipses.

1995), and quality control was achieved using the Plésovice standard (338 ± 1 Ma, Slama et al., 2008).

The sensitivity of the machine permits recording ^{204}Pb to check for possible lead contamination of the zircon grains. As the 204 mass could be influenced by the presence of ^{204}Hg , the ^{202}Hg (only candidate for this mass) was recorded in order to correct this interference. Then, the corrected mass 204 supposed to only represent common lead was used to correct the sample's analyses. The percentage of common lead revealed by the ^{204}Pb measurement in each analysis is given in Table SM 1. Five analyses among eleven were shown to be weakly contaminated by common lead (spots 3, 4, 7, 8 and 11) with $f^{204}\text{Pb} < 0.4\%$. Only those five analyses were used to calculate a concordant age. Data reduction was carried out with the GLITTER software package (van Achterbergh et al., 2001), but the common lead correction was performed with an in-house built program. The results are presented in Fig. 3D with common lead corrected data and generated using the ISOPLOT/EX (v. 3) software by Ludwig (2003).

3.2. Results

Eleven ablations were performed on seven zircon grains (Table SM 1). Analyses 1 and 2 were performed on inherited zircon grain #1, whose significance will be discussed further. These analyses were not used in the set of data to calculate the age of the sample. Analysis 6 was performed on a chemically disturbed zircon grain #4 (see ESM image), and analyses 5 and 9 were performed on the heterogeneous part of zircon grains. Spot 10 has been performed on an apparently fairly homogeneous zircon, but some evidence of cracks on grain #6 could have disturbed the corresponding measurement. Those four analyses, performed on heterogeneous areas, display too much common lead contamination to be effectively corrected; they are plotted as dotted ellipses in Fig. 3D. The five other analyses (spot 3, 4, 7, 8 and 11) were performed on apparently homogeneous areas showing a similar luminescence on each grain. Those five analyses are

less contaminated with common lead, and the corrected analyses provide a concordant age of 190 ± 2 Ma on the Wetherill diagram (Fig. 3D, full line). The fit of the analyses vs. the Concordia diagram provides robust constraint on the age of crystallization of these zircon grains.

Analyses 1 and 2 performed on grain #1, i.e. the inherited, well-faceted prismatic and pyramid-topped zircon grain revealed ^{204}Pb content and could have suffered lead loss. On the Wetherill diagram of Fig. 4, these two analyses after ^{204}Pb correction plot onto a straight line passing through the origin (75 ± 60 Ma) without any constraint (for comparison, the analyses without common lead correction are plotted as dotted gray ellipse). This could reveal that the lead-loss process was recent. Accordingly, this grain cannot be dated with precision, but an age of crystallization at about 2 Ga is likely.

4. Discussion

4.1. The previous interpretation of the Mesorif suture zone might be abandoned

In line with the early interpretation of the Beni Malek serpentinites as exhumed mantle from the Ocean–Continent Transition (OCT) of the Late Jurassic North-African margin (Michard et al., 1992), the curved alignment of gabbros massifs with similar oceanic sedimentary cover that forms the main part of the Mesorif suture zone (Fig. 2) has been regarded as the trace of the same OCT, displaced by the Miocene thrust tectonics (Benzaggagh et al., 2014; Michard et al., 2014). In other words, the gabbro slivers of the Mesorif suture zone were regarded as emplaced in the framework of the Jurassic opening of the Maghrebian-Ligurian Tethys, which is known to be basically Callovian-Oxfordian to Tithonian in age (ca. 166–150 Ma; Lemoine et al., 1987; Li et al., 2013) although the earliest isotopic dates are Bajocian (ca. 170 Ma; Bill et al., 2001). This hypothesis was consistent with a K–Ar whole-rock age of 166 ± 3 Ma (Bathonian–Callovian) obtained on the Bou Adel

gabbro by J.-M. Cantagrel at the Clermont-Ferrand Laboratory, and published by Asebriy (1994, p. 51). However, the limits of the K-Ar method when applied to low-grade metamorphic gabbros such as those of the Mesorif suture zone are well known (Ducrot and Lancelot, 1977). For this reason, we decided to obtain a robust, U-Pb zircon age from the Bou Adel trondhjemite veins that were emplaced by the end of the gabbro crystallization. Now, the Liassic age that we obtained contradicts our previous interpretation.

4.2. The Mesorif Basalts–Breccias (MBB) lineament and the new proposal for the MSZ

In the External Rif belt, an arcuate alignment of basalts and carbonate breccias, i.e. the Mesorif Basalts–Breccias lineament, almost parallels the MSZ in a more external position (Fig. 2). Along this lineament, the Kimmeridgian–Tithonian/Berriasian sequences of the external Mesorif–internal Prerif units are typified by tholeiitic lava flows and/or calcareous breccias, including tholeiitic basalt, dolerite, and fine-grained gabbro elements (Benzaggagh, 2011; Benzaggagh et al., 2014). This mix of carbonate and magmatic formations occurs everywhere on top of a kilometer-thick Callovian–Oxfordian turbidite formation (the “ferrysch”) that seals the Lower–Middle Jurassic tilted-block and half-graben structures of the North-African paleomargin (Suter, 1980a; Favre, 1992). Most of the elements of the breccias are shallow water carbonates reworked into deep waters by mass flows and turbidity currents at the expense of a more external carbonate platform. The oldest of the associated magmatic rocks are pyroclastic layers emplaced by the end of the Oxfordian, and the latest are disrupted basalt flows included in the Upper Tithonian–Berriasian breccias. The chaotic breccias also contain pebbles or boulders of ferrysch-like sandstones, whose abundance and size increase upward. Large olistoliths of typical ferrysch facies occur locally in the Berriasian sequence (Benzaggagh et al., 2014).

The tectonic significance of the reported magmatic-sedimentary association has been underestimated up to

now, although in their study of the western Mesorif (Ouezzane area), Ben Yaïch et al. (1989) noticed that it would indicate an abrupt thinning of the North-African margin. We think it necessary to interpret the association at the scale of the entire Mesorif considering two significant observations, (i) the parallelism of the Basalts–Breccias lineament with the Mesorif suture, and (ii) the slightly diverging thrust transport directions of the External Rif tectonic units during the Miocene (Fig. 2). These directions are typically oriented to the southwest in the eastern and central Rif, whereas they progressively turn to the WSW and the west in the western Rif (Chalouan et al., 2006, 2014; Frizon de Lamotte, 1985, 1987; Poujol et al., 2017).

Keeping in mind the above remarks, and based on the 190 ± 2 Ma age of the Bou Adel gabbro, we may explore a new avenue to explain both the Mesorif suture and Basalts–Breccias lineaments in the framework of the Central Atlantic–Alpine Tethys relationships. The connection between these two oceanic domains implies classically the activity of a broadly latitudinal North-African Transform (NAT; Fig. 5A) during the Jurassic–Early Cretaceous times (Lemoine et al., 1987; Rosenbaum et al., 2002) or that of an equivalent, east–west system of densely segmented oceanic ridges (Fig. 5B; Frizon de Lamotte et al., 2011; Sallarès et al., 2011). Schettino and Turco (2011) combine a North African transform-equivalent labeled the “Gibraltar Fault” with a narrow, segmented oceanic corridor in their geodynamic reconstruction for the Late Jurassic period. However, the opening of the Central Atlantic began at ca. 190 Ma (Sinemurian–Pliensbachian; Labails et al., 2010), whereas the Alpine Tethys opened not before 170 Ma (Bajocian) at the earliest (Bill et al., 2001; Lagabrielle and Lemoine, 1997). This diachronism, seldom considered up to now, makes less simple the interpretation of the crustal movements at the Rifian corner of North Africa. Moreover, in the studied area, the strike and kinematics of the major faults depend on the movement of the Iberian plate with respect to Africa and Europe, which is still discussed (Barnett-Moore et al., 2016; Vissers and Meijer, 2012). Plate tectonic reconstructions indicate that the North African




Fig. 5. Two published reconstructions of the connection between Central Atlantic and Tethys oceans during the Late Jurassic. A. Classical view involving a sinistral North African transform zone (NAT), after Rosenbaum et al. (2002). B. Alternative reconstruction involving a narrow, segmented oceanic corridor between Africa and Iberia, after Sallarès et al. (2011). In both proposals, the Central Atlantic northeast-striking ridge is sharply interrupted by the east-trending transform or oceanic corridor.

transform was a sinistral transform zone during the Late Jurassic, but that it was dextral during the Aptian–Albian, due to the anticlockwise rotation of Iberia (Vissers and Meijer, 2012, and references therein).

We suggest that the magmatic-sedimentary dramatic events recorded along the MBB lineament during the Late Jurassic–Berriasian are linked to the Late Jurassic activity of the North African transform, either along the main transform zone or along some synthetic fault. The interpretation of the broadly parallel Mesorif suture zone lineament is more problematic in the present state of knowledge. One of the gabbroic bodies (Bou Adel) has been dated from the Early/Middle Liassic, but its oceanic-type cover has not been dated yet, nor the low-grade metamorphism observed in these exotic bodies. The association of basaltic breccia, radiolarites, marbles and ophiolitic sands offers a clear Ligurian affinity, and thus can be hypothetically ascribed to the Late Jurassic. At this step of our interpretation, we may propose two alternative hypotheses: (i) the Early–Middle Liassic Bou Adel gabbro and its equivalents would have been exhumed at the northern tip of a NNE-trending Atlantic OCT, next to the distal Continent–Ocean Boundary (COB; see Klingelhofer et al., 2016), then transported along a dextral shear zone during the Middle Jurassic before being overlain by their

sedimentary covers, or (ii) the Mesorif suture gabbros would underline the eastward curvature of the Atlantic margin, and they were exhumed during the Middle Jurassic in the vicinity of the North African transform. The first hypothesis cannot be maintained, as no major dextral movement is documented there during the Middle Jurassic by plate tectonic reconstructions. Therefore, we favor the hypothesis of a curvature of the Atlantic margin that would have surrounded the NW Meseta promontory (Sehoul Block; Michard et al., 2010) to connect eastward with the South Tethyan margin (Fig. 6).

The latter figure is intended to make visible the displacements that the varied tectonic units of the Rif suffered with respect to the Atlas–Meseta foreland. During the Miocene, the movement along the North African transform was sinistral, as this shear zone allowed the westward displacement of the Alboran Domain due to the Gibraltar slab retreat (Jolivet and Faccenna, 2000; van Hinsbergen et al., 2014). The Alboran Domain displacement along the Miocene North African transform is estimated by the latter authors in the range of 400–500 km. The elements belonging to the southern, North African compartment of the Miocene transform were also affected by a westward simple shear deformation (demonstrated in the Temsamane massif of the Mesorif; Frizon de Lamotte, 1985, 1987), but




Fig. 6. New proposal for the Central Atlantic-Alpine Tethys connection involving an eastward curvature of the Atlantic margin around the Moroccan Meseta promontory (this work). This proposal is based on the Liassic age of the Bou Adel gabbro of the Mesorif Suture Zone (MSZ). The curved segment of the Liassic Atlantic margin would have connected due to rift propagation with the Middle Jurassic North African Transform (NAT) whose transported elements form the Mesorif Basalt-Breccia (MBB) lineament. During the Miocene, the Bou Adel massif would have reached its present-day location due to the shortening and SW-ward transport of the External Rif accretionary prism.

they were certainly less displaced than the Alboran Domain itself. At the same time, Africa–Iberia convergence resulted in the building of the External Rif accretionary prism and its tectonic transport toward the foreland. As a conservative hypothesis, we may estimate a southwestward tectonic transport of about 100 km for the Bou Adel-type massifs of the Mesozoic suture. Thus, “unfolding” the External Rif tectonic prism allows us to restore the plausible initial location of the Mesozoic Basalts–Breccias and Suture Zone lineaments along the Jurassic North African transform (Fig. 6). The 30 Ma delay between the emplacement of the Bou Adel gabbro in the MSZ and the MBB fault zone activity would correspond to the propagation of the Central Atlantic rift both eastward (toward the Alpine Tethys) and southward (encroaching the proximal African margin). The Central Atlantic rift would have propagated eastward over about 2000 km in about 20–30 Ma. The rift propagation would have thus occurred at about 10 cm/yr, which is a propagation rate consistent with rates found for other rifts and fault zones worldwide (e.g., Hubert-Ferrari et al., 2003; Manighetti et al., 1997, and references therein).

Two observations support our interpretation. First, the Triassic evaporitic basins that record the earliest rifting of Pangea along the Atlantic margin, culminating with the CAMP magmatism at 201–195 Ma (Leleu et al., 2016; Verati et al., 2007), also surround the NW Moroccan Meseta promontory and are well developed in the Prerif-Mesorif domain itself. Second, the “Slope anomaly” S (i.e. the oldest magnetic anomaly on the African side, also labeled “West Atlantic Margin Anomaly”) adopts an ENE trend west of Tangier, beneath the accretionary prism of the Gulf of Cadiz (Klingelhofer et al., 2016). Martínez-Lorient et al. (2014) also draw a curvature of the Central Atlantic ridge in their reconstruction of the 180 Ma stage (their fig. 10), but their subsequent 155 Ma reconstruction shows a rectilinear Central Atlantic ridge sharply crosscut by the Gibraltar Fault. On the other hand, the presence in the Bou Adel trondhjemite of a Proterozoic (ca. 2 Ga) xenocrystic zircon reveals the existence of an ancient continental crust, possibly inherited from the WAC, beneath the locus of emplacement of the trondhjemite as it is the case for the trondhjemite from the Balagne nappe in Corsica (Rossi et al., 2002) or in the Mauritius trachyte (Ashwal et al., 2017). This is consistent with the MSZ gabbro emplacement in the OCT domain along the northern Meseta. Indeed, the Rabat granite of the Sehoul Block yielded 2-Ga-old inherited zircon grains (Tahiri et al., 2010), and farther to the south, the Meseta Coastal Block displays a shallow, and by place outcropping Paleoproterozoic basement (Letsch et al., 2017; Pereira et al., 2015).

5. Conclusion

At the northwestern corner of the African plate, the ultimate limit of the WAC and WAC-related domains corresponds nowadays to the diffuse Iberia–Africa plate boundary that extends from the Gulf of Cadiz to the Rif–Tell orogen. This plate boundary is inherited from the sinistral Miocene transform zone, active during the westward displacement of the Alboran Domain. The Miocene transform is itself inherited from the Jurassic

transform zone, active during the Pangea breakup and the diachronic opening of the Central Atlantic and Alpine Tethys oceans. The present work suggests that a deformed trace of the Jurassic transform or of some synthetic fault can be found in the Mesorif Basalts–Breccias lineament. Our work also suggests that the arcuate Mesorif Suture Zone, adjacent to the MBB, would represent the ENE-striking continuation of the Atlantic margin.

This preliminary hypothesis is based on the U–Pb zircon dating at 190 ± 2 Ma of one of the gabbro massifs of the Mesorif zone, which appears to be coeval with the early opening of the Central Atlantic. The chaotic formations of the nearby Basalts–Breccias lineament, mainly Kimmeridgian–Tithonian in age, would delineate the border of the North-African margin along the Jurassic transform zone. If correct, this new interpretation can provide the impetus for performing additional field and laboratory studies of the exhumed samples of the up-to-now conceptual Jurassic transform zone.

Acknowledgments

Logistic support for field studies was granted by the Ministry of Energy and Mines, Water and Environment, Rabat. We warmly thank Prof. Marco Chiari for his kind tentative to determine radiolarian remains in four of our samples, collected with the help of our friends Faouziya Haissen and Najib B. Zaghloul. We warmly thank our colleagues Philippe Olivier and Marc-André Gutscher, who achieved thorough and constructive peer reviews of the submitted manuscript. We also thank Isabelle Manighetti who kindly organized the reviewing process and suggested useful improvements to our manuscript.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.crte.2018.05.006>.

References

- Asebriy, L., 1994. *Évolution tectonique et métamorphique du Rif central (Maroc) : définition du domaine subrifain* (Unpublished Thesis, Université Mohamed-V, Rabat-Agdal (283 p.).
- Ashwal, L.D., Wiedenbeck, M., Torsvik, T.H., 2017. Archaean zircons in Miocene oceanic hotspot rocks establish ancient continental crust beneath Mauritius. *Nature*, <http://dx.doi.org/10.1038/ncomms14086>.
- Barnett-Moore, N., Hosseinpour, M., Maus, S., 2016. Assessing discrepancies between previous plate kinematic models of Mesozoic Iberia and their constraints. *Tectonics* 35, <http://dx.doi.org/10.1002/2015TC004019>.
- Bea, F., Montero, P., Haissen, F., Michard, A., Lazar, C., Mouttagi, A., Errami, A., Sadki, O., 2016. First Evidence of Cambrian rift-related magmatism in the West African margin: The Derraman Peralkaline Felsic Complex. *Gondwana Res.* 36, 423–438.
- Ben Yaïch, A., Hervouët, Y., Duée, G., El Hatimi, N., 1989. Âge jurassique et réinterprétation des roches basiques au nord de Ouezzane (Rif externe, Maroc) : signification géodynamique. *C. R. Acad. Sci. Paris, Ser. II* 309, 1197–1202.
- Benzaggagh, M., 2011. Activités volcaniques sous-marines à la limite Jurassique-Créacé dans le Rif externe (Maroc). Âge et relation avec la sédimentation et la paléogéographie du sillon rifain externe. *C. R. Geoscience* 343, 302–311.

- Benzaggagh, M., Mokhtari, A., Rossi, P., Michard, A., El Maz, A., Chalouan, A., Saddiqi, O., Rjimati, E., 2014. Oceanic units in the core of the External Rif (Morocco): intramargin hiatus or South-Tethyan remnants? *J. Geodyn.* 77, 4–21.
- Biari, Y., Klingelhoefer, F., Sahabi, M., Funck, T., Benabdellouahed, M., Schnabel, M., Reichert, C., Gutscher, M.-A., Bronner, A., Austin, J.A., 2017. Opening of the central Atlantic Ocean: implications for geometric rifting and asymmetric initial seafloor spreading after continental breakup. *Tectonics* 36, 1129–1150, <http://dx.doi.org/10.1002/2017TC004596>.
- Bill, M., O'Dogherty, L., Guex, J., Baumgartner, P.O., Masson, H., 2001. Radiolarite ages in Alpine-Mediterranean ophiolites: constraints on the oceanic spreading and the Tethys–Atlantic connection. *Geol. Soc. Amer. Bull.* 113, 129–143.
- Bosch, D., Bruguer, O., Caby, R., Buscaill, F., Hammor, D., 2016. Orogenic development of the Adrar des Iforas (Tuareg Shield, NE Mali): new geochemical and geochronological data and geodynamic implications. *J. Geodyn.* 96, 104–130.
- Bouillin, J.-P., Kornprobst, J., Raoult, J.-F., 1997. Données préliminaires sur le complexe volcanico-sédimentaire de Rekkada-Metletine (ex-Texenna) en Petite Kabylie (Algérie). *Bull. Soc. Geol. Fr.* 19 (7), 805–813.
- Caby, R., 2003. Terrane assembly and geodynamic evolution of central-western Hoggar: a synthesis. *J. Afr. Earth Sci.* 37, 133–159.
- Chalouan, A., El Mrihi, A., El Kadiri, K., Bahmad, A., Salhi, F., Hlila, R., 2006. Mauretanian flysch nappe in the northwestern Rif Cordillera (Morocco): deformation chronology and evidence for a complex nappe emplacement. In: Moratti, G., Chalouan, A. (Eds.), *Tectonics of the western Mediterranean and north Africa*. Geol. Soc. London, Spec. Publ., 262, pp. 161–175.
- Chalouan, A., Michard, A., El Kadiri, Kh., Negro, F., Frizon de Lamotte, D., Soto, J.I., Saddiqi, O., 2008. The Rif belt. In: Michard, A., Saddiqi, O., Chalouan, A., Frizon de Lamotte, D. (Eds.), *Continental Evolution: The Geology of Morocco: Structure, Stratigraphy and Tectonics of the Africa-Atlantic–Mediterranean Triple Junction*. Lect. Notes Earth Sci., 116, pp. 203–302.
- Chalouan, A., Michard, A., El Kadiri, Kh., Saddiqi, O., 2011. Rif central et nord-occidental. In: Michard, A., Saddiqi, O., Chalouan, A., Rjimati, E.C., Mouttaqi, A. (Eds.), *Nouveaux guides géologiques et miniers du Maroc*. Notes Mem. Serv. géol. Maroc, 560, pp. 9–90.
- Chalouan, A., Gil, A.J., Galindo-Zaldívar, J., Ahmamou, M., Ruano, P., de Lacy, M.C., Ruiz-Armenteros, A.M., Bennakhoulf, M., Riguzzi, F., 2014. Active faulting in the frontal Rif Cordillera (Fes region, Morocco): constraints from GPS data. *J. Geodyn.* 77, 110–122.
- Crutchley, G.J., Berndt, C., Klaeschen, D., Masson, D.G., 2011. Insights into active deformation in the Gulf of Cadiz from new 3-D seismic and highresolution bathymetry data. *Geochem. Geophys. Geosyst.* 12, Q07016, <http://dx.doi.org/10.1029/2011GC003576>.
- Davison, I., Daily, P., 2010. Salt tectonics in the Cap Boudjour Area, Aaiun Basin, NW Africa. *Marine Petroleum Geol.* 27, 435–441.
- Ducrot, J., Lancelot, L.R., 1977. Problème de la limite Précambrien-Cambrien : étude radiochronologique par la méthode U-Pb sur zircons du volcan du Jbel Boho (Anti-Atlas F marocain). *Can. Earth J. Sci.* 14, 2771–2777.
- Durand-Delga, M., 1980. La Méditerranée occidentale, étapes de sa genèse et problèmes structuraux liés à celle-ci. *Mem. Soc. Geol. France* 10, 203–224.
- Durand-Delga, M., Rossi, P., Olivier, P., Puglisi, D., 2000. Situation structurale et nature ophiolitique de roches basiques jurassiques associées aux flyschs maghrébins du Rif (Maroc) et de Sicile (Italie). *C. R. Acad. Sci. Paris, Ser. IIa* 331, 29–38.
- Elazzab, D., Galdeano, A., Feinberg, H., Michard, A., 1997. Prolongement en profondeur d'une écale ultrabasique allochtone : traitement des données aéromagnétiques et modélisation 3D des péridotites des Beni Malek (Rif, Maroc). *Bull. Soc. Geol. France* 168, 667–683.
- El Houicha, M., Pereira, M.F., Jouhari, A., Gama, C., Ennih, N., Fekkak, A., Ezzouhairi, H., El Attari, A., Silva, J.B., 2018. Recycling of the Proterozoic crystalline basement in the Coastal Block (Moroccan Meseta): New insights for understanding the geodynamic evolution of the northern peri-Gondwanan realm. *Precambrian Res.* 306, 129–154.
- El Talibi, H., Zaghoul, M.N., Perri, F., Aboumaria, K., Rossi, A., El Moussaoui, S., 2014. Sedimentary evolution of the siliciclastic Aptian-Albian Massylian flysch of the Chouamat Nappe (central Rif, Morocco). *J. Afr. Earth Sci.* 100, 554–568.
- Ennih, N., Liégeois, J.-P., 2001. The Moroccan Anti-Atlas: the West African craton passive margin with limited Pan-African activity. Implications for the northern limit of the craton. *Precambrian Res.* 112, 289–302.
- Ennih, N., Liégeois, J.-P., 2008. The boundaries of the West African craton, with special reference to the basement of the Moroccan metacratonic Anti-Atlas belt. *Geol. Soc. London Spec. Publ.* 297, 1–17.
- Favre, P., 1992. Géologie des massifs calcaires situés au front sud de l'unité de Kétama (Rif, Maroc). *Publ. Départ. Geol. Paleontol. Univ. Genève* 11, (138 p.).
- Favre, P., Stampfli, G., Wildi, W., 1991. Jurassic sedimentary record and tectonic evolution of the northwestern corner of Africa. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 87, 53–73.
- Frizon de Lamotte, D., 1985. La structure du Rif oriental (Maroc), rôle de la tectonique longitudinale et importance des fluides. Thèse d'Etat. Université Paris-6 (85-03, 436 p.).
- Frizon de Lamotte, D., 1987. La structure du Rif externe (Maroc) : mise au point sur le rôle des décrochements, des chevauchements et des glissements gravitaires. *J. Afr. Earth Sci.* 6, 755–766.
- Frizon de Lamotte, D., Saint Bezar, B., Bracène, R., Mercier, E., 2000. The two main steps of the Atlas building and geodynamics of the western Mediterranean. *Tectonics* 19, 740–761, <http://dx.doi.org/10.1029/2000TC900003>.
- Frizon de Lamotte, D., Zizi, M., Missenard, Y., Hafid, M., El Azzouzi, M., Maury, R.C., Charrière, A., Taki, Z., Benammi, M., Michard, A., 2008. The Atlas System. In: Michard, A., Saddiqi, O., Chalouan, A., Frizon de Lamotte, D. (Eds.), *Continental evolution: the Geology of Morocco*. Lect. Notes Earth Sci., 116, pp. 133–202.
- Frizon de Lamotte, D., Raulin, C., Mouchot, N., Wrobel-Daveau, J.C., Blanpied, C., Ringenbach, J.C., 2011. The southernmost margin of the Tethys realm during the Mesozoic and Cenozoic: initial geometry and timing of the inversion processes. *Tectonics* 30, TC3002, <http://dx.doi.org/10.1029/2010TC002691>.
- Frizon de Lamotte, D., Zaghoul, M.N., Haissem, F., Mohn, G., Leprêtre, R., Gimeno-Vives, O., Atouabat, A., El Mourabet, M., Abassi, A., 2017. Rif externe : comment comprendre et expliquer le chaos apparent ? *Géologues Soc. Geol. Fr.* 194, 13–15.
- Galindo-Zaldívar, J., Maldonado, A., Schreider, A.A., 2003. Gorringe Ridge gravity and magnetic anomalies are compatible with thrusting at a crustal scale. *Geophys. J. Int.* 153, 586–594, <http://dx.doi.org/10.1046/j.1365-246X.2003.01922.x>.
- Gärtner, A., Villeneuve, M., Linnemann, U., Gerdes, A., Youbi, N., Guillou, O., Rjimati, E.C., 2014. History of the West African Neoproterozoic Ocean: key to the geotectonic history of circum-Atlantic Peri-Gondwana (Adrar Soutouf Massif, Moroccan Sahara). *Gondwana Res.* 29, 220–233.
- Guerrera, F., Martín-Martín, M., 2014. Geodynamic events reconstructed in the Betic, Maghrebian, and Apennine chains (central-western Tethys). *Bull. Soc. Geol. France* 185, 329–341.
- Gutscher, M.-A., Dominguez, S., Westbrook, G.K., Le Roy, P., 2009. Deep structure, recent deformation and analog modeling of the Gulf of Cadiz accretionary wedge: implications for the 1755 Lisbon earthquake. *Tectonophysics* 475, 85–97.
- Hafid, M., Tari, G., Bouhadioui, B., El Moussaid, I., Eccharfaoui, H., Ait Salem, A., Nahim, M., Dakki, M., 2008. Atlantic Basins. In: Michard, A., Saddiqi, O., Chalouan, A., Frizon de Lamotte, D. (Eds.), *Continental Evolution: The Geology of Morocco*. Lect. Notes Earth Sci., 116, pp. 303–329.
- Hefferan, K., Soulaïmani, A., Samson, S., Admou, H., Inglis, J., Saquaque, A., Heywood, N., 2014. A reconsideration of the Pan-African orogenic cycle in the Anti-Atlas Mountains, Morocco. *J. Afr. Earth Sci.* 98, 34–46.
- Hoepffner, C., Ouanaïmi, H., Michard, A., 2017. La Meseta, un terrain vagabond ou la marge fragmentée de l'Anti-Atlas ? *Géologues Soc. Geol. Fr.* 194, 19–23.
- Hubert-Ferrari, A., King, G., Manighetti, I., Armijo, R., Meyer, B., Tapponnier, P., 2003. Long-term elasticity in the continental lithosphere modelling the Aden Ridge propagation and the Anatolian extrusion process. *Geophys. J. Int.* 153, 111–132.
- Jolivet, L., Faccenna, C., 2000. Mediterranean extension and the Africa-Eurasia collision. *Tectonics* 19, 1094–1106.
- Klingelhoefer, F., Biari, Y., Sahabi, M., Aslanian, D., Schnabel, M., Matias, L., Benabdellouahed, M., Funck, T., Gutscher, M.A., Reict, C., Austin, J.A., 2016. Crustal structure variations along the NW-African continental margin comparison of new: A existing models from wide-angle and reflection seismic data. *Tectonophysics* 674, 227–252.
- Labails, C., Olivet, J.L., Aslanian, D., Roest, W.R., 2010. An alternative early opening scenario for the Central Atlantic Ocean. *Earth Planet. Sci. Lett.* 297 (3), 355–368.
- Lagabrielle, Y., Lemoine, M., 1997. Alpine, Corsican and Apennine ophiolites: the slow-spreading ridge model. *C. R. Acad. Sci. Paris, Ser. II* 325, 909–920.
- Leblanc, D., 1979. Étude géologique du Rif externe oriental au nord de Taza (Maroc). *Notes Mem. Serv. géol. Maroc* 281 (159 p.).
- Le Goff, E., Guerrot, C., Maurin, G., Johan, V., Tegyey, M., Ben Zarga, M., 2001. Découvertes d'éclogites hercyniennes dans la chaîne septentrionale des Mauritanides (Afrique de l'Ouest). *C. R. Acad. Sci. Paris, Ser. IIa* 333, 711–718.

- Leleu, S., Hartley, A.J., van Oosterhout, C., Kennan, L., Ruckwied, K., Gerdes, K., 2016. Structural, stratigraphic and sedimentological characterization of a wide rift system: The Triassic rift system of the Central Atlantic Domain. *Earth Sci. Rev.* 158, 89–124.
- Lemoine, M., Tricart, P., Boillot, G., 1987. Ultramafic and gabbroic ocean floor of the Ligurian Tethys (Alps, Corsica, Apennines): in search of a genetic model. *Geology* 15, 622–625.
- Letsch, D., El Houicha, M., von Quadrt, A., Winkler, W., 2017. A missing link in the peri-Gondwanan terrane collage: the Precambrian basement of the Moroccan Meseta and its lower Paleozoic cover. *Can. J. Earth Sci.*, <http://dx.doi.org/10.1139/cjes-2017-0086>.
- Li, X.H., Faure, M., Lin, W., Manatschal, G., 2013. New isotopic constraints on age and magma genesis of an embryonic oceanic crust: the Chenaillet Ophiolite in the Western Alps. *Lithos* 160–161, 283–291.
- Ludwig, K.R., 2003. ISOPLOT/EX, version 3. A geochronological toolkit for Microsoft Excel. Berkeley Geochronology Center, Spec. Publ. 4. (70 p.).
- Manighetti, I., Tapponnier, P., Courtillot, V., Gruszow, S., Gillot, P.Y., 1997. Propagation of rifting along the Arabia–Somalia plate boundary: The gulfs of Aden and Tadjoura. *J. Geophys. Res. Solid Earth* 102 (B2), 2681–2710.
- Martínez-Loriente, S., Sallarès, V., Gràcia, E., Bartolome, R., Dañobeitia, J.J., Zitellini, N., 2014. Seismic and gravity constraints on the nature of the basement in the Africa–Eurasia plate boundary: New insights for the geodynamic evolution of the SW Iberian margin. *J. Geophys. Res. Solid Earth* 119, 127–149.
- Mascle, J., Mascle, G., 2012. Geological and morpho-tectonic map of the Mediterranean Domain. Publ. CGMW and UNESCO.
- Meghraoui, M., Pondrelli, S., 2012. Active faulting and transpression tectonics along the plate boundary in North Africa. *Annals Geophys.* 55, <http://dx.doi.org/10.4401/ag-4970>.
- Michard, A., Feinberg, H., Elazzab, D., Bouybaouene, M.L., Saddiqi, O., 1992. A serpentinite ridge in a collisional paleomargin setting: the Beni Malek massif, External Rif, Morocco. *Earth Planet. Sci. Lett.* 113, 435–442.
- Michard, A., Frizon de Lamotte, D., Negro, F., 2007. Serpentinite slivers and metamorphism in the External Maghrebides: Arguments for an intra-continental suture in the African paleomargin (Morocco, Algeria). *Rev. Soc. Geol. España* 20, 173–185.
- Michard, A., Soulaïmani, A., Hoepffner, C., Ouainaimi, H., Baïdder, L., Rjimati, E.C., Saddiqi, O., 2010. The south-western branch of the Variscan belt: evidence from Morocco. *Tectonophysics* 492, 1–24.
- Michard, A., Mokhtari, A., Chalouan, A., Saddiqi, O., Rossi, P., Rjimati, E., 2014. New ophiolite slivers in the External Rif Belt, and tentative restoration of a dual Tethyan suture in the Western Maghrebides. *Bull. Soc. Geol. France* 185, 313–328.
- Montero, P., Bea, F., Haissen, F., Molina, J.F., González-Lodeiro, F., Mouttaqi, A., Errami, A., 2017. Dorsale Reguibat et massif des Ouled Dlim, l'avancée des connaissances. *Géologues Soc. Geol. Fr.* 194, 37–46.
- Olivet, J.-L., Bonninn, J., Auzende, J.M., 1984. Cinématique de l'Atlantique nord et central. Centre national pour l'exploration des océans. Rapport scientifiques et techniques de l'Ifremer No. 54. (108 p.).
- Olivier, P., Durand-Delga, M., Manivit, H., Feinberg, H., 1996. Le substratum jurassique des flyschs maurétaniens de l'ouest des Maghrebides : l'unité de Ouareg (région de Targuit, Rif, Maroc). *Bull. Soc. Geol. France* 167, 609–616.
- Ouabid, M., Ouali, H., Garrido, C.J., Acosta-Vigil, A., Román-Alpiste, M.J., Dautria, J.M., Marchesi, C., Hidas, K., 2017. Neoproterozoic granitoids in the basement of the Moroccan Central Meseta: Correlation with the Anti-Atlas at the NW paleo-margin of Gondwana. *Precambrian Res.* 299, 34–57.
- Pereira, M.F., El Houicha, M., Chichorro, M., Armstrong, R., Jouhari, A., El Attari, A., Ennih, N., Silva, J.B., 2015. Evidence of a Paleoproterozoic basement in the Moroccan Variscan Belt (Rehamna Massif, Western Meseta). *Precambrian Res.* 268, 61–73.
- Poujol, A., Ritz, J.F., Vernant, P., Huot, S., Maate, S., Tahayt, A., 2017. Which fault destroyed Fes city (Morocco) in 1755? A new insight from the Holocene deformations observed along the southern border of Gibraltar arc. *Tectonophysics* 712–713, 303–311.
- Rosenbaum, G., Lister, G.S., Duboz, C., 2002. Relative motion of Africa, Iberia and Europe during Alpine orogeny. *Tectonophysics* 339, 117–126.
- Rossi, P., Cocherie, A., Lahondère, D., Fanning, C.M., 2002. La marge européenne de la Téthys jurassique en Corse : datation de trondhjemites de Balagne et indices de croûte continentale sous le domaine balano-ligure. *C. R. Geoscience* 334, 313–322.
- Sallarès, V., Gailler, A., Gutscher, M.A., Graindorge, D., Bartolome, R., Gràcia, E., Díaz, J., Dañobeitia, J.J., Zitellini, N., 2011. Seismic evidence for the presence of Jurassic oceanic crust in the central Gulf of Cadiz (SW Iberia margin). *Earth Planet. Sci. Lett.* 311, 112–123.
- Schettino, A., Turco, E., 2011. Tectonic history of the western Tethys since the Late Triassic. *Geol. Soc. Am. Bull.* 123, 89–105.
- Slama, J., Kosler, J., Condon, D.J., Crowley, J.L., Gerdes, A., Hanchar, J.M., Horstwood, M.S.A., Morris, A.G., Nasdala, L., Norberg, N., Schaltegger, U., Schoene, B., Tubrett, M.N., Withehouse, M.J., 2008. Plésovice zircon—A new natural reference material for U–Pb and Hf isotopic microanalysis. *Chemical Geol.* 249, 1–35.
- Suter, G., 1980a. Carte géologique du Rif, 1/500 000. Notes Mem. Serv. Geol. Maroc. 245a.
- Suter, G., 1980b. Carte structurale du Rif, 1/500 000. Notes Mem. Serv. Geol. Maroc. 245b.
- Tahiri, A., Montero, P., El Hadi, H., Martínez Poyatos, D., Azor, A., Bea, F., Simancas, J.F., González Lodeiro, F., 2010. Geochronological data on the Rabat–Tiflet granitoids: their bearing on the tectonics of the Moroccan Variscides. *J. Afr. Earth Sci.* 57, 1–13.
- van Achterbergh, E., Ryan, C.G., Jackson, S.E., Griffin, W., 2001. Data reduction software for LA-ICP-MS. In: Sylvester, P. (Ed.), *Laser Ablation–ICPMs in the Earth Science*, 29, Mineralogical Association of Canada, St John's, Newfoundland, Canada, pp. 239–243.
- van Hinsbergen, D.J.J., Vissers, R.L.M., Spakman, W., 2014. Origin and consequences of western Mediterranean subduction, rollback, and slab segmentation. *Tectonics* 33, 393–419, <http://dx.doi.org/10.1002/tect.20125>.
- Verati, C., Rapaille, C., Féraud, G., Marzoli, A., Bertrand, H., Youbi, N., 2007. $^{40}\text{Ar}/^{39}\text{Ar}$ ages and duration of the Central Atlantic Magmatic Province volcanism in Morocco and Portugal and its relation to the Triassic–Jurassic boundary. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 244, 308–325.
- Villeneuve, M., Bellon, H., El Archi, A., Sahabi, M., Rehault, J.-P., Olivet, J.-L., 2006. Événements panafricains dans l'Adrar Soutouf (Sahara marocain). *C. R. Geoscience* 338, 359–367.
- Vissers, R., Meijer, P.T., 2012. Mesozoic rotation of Iberia: Subduction in the Pyrenees? *Earth Sci. Rev.* 110, 93–110.
- Wiedenbeck, M., Allé, P., Corfu, F., Griffin, W.L., Meier, M., Oberli, F., von Quadrt, A., Roddick, J.C., Spiegel, W., 1995. Three natural zircon standards for U–Th–Pb, Lu–Hf, trace element and REE analysis. *Geostandards Newslett.* 19, 1–23.
- Zaghoul, M.N., Di Staso, A., Gigliuto, L.G., Maniscalco, R., Puglisi, D., 2005. Stratigraphy and provenance of Lower and Middle Miocene strata within the External Tanger Unit (Intra-Rif sub-Domain, External Domain; Rif, Morocco): first evidence. *Geologica Carpathica* 56, 517–530.
- Zaghoul, M.N., Di Staso, A., de Capoa, P., Perrone, V., 2007. Occurrence of upper Burdigalian silexite beds within the Beni Ider Flysch Fm. in the Ksar-es-Seghir area (Maghrebian Flysch Basin, Northern Rif, Morocco): stratigraphic correlations and geodynamic implications. *Boll. Soc. Geol. It.* 126, 223–239.