

Lithium isotopes in the Loire River Basin (France): hydrogeochemical characterizations at two complementary scales

Romain Millot, Philippe Négrel

► To cite this version:

Romain Millot, Philippe Négrel. Lithium isotopes in the Loire River Basin (France): hydrogeochemical characterizations at two complementary scales. Applied Geochemistry, 2021, 125 (104831), 10.1016/j.apgeochem.2020.104831. hal-03113284

HAL Id: hal-03113284 https://brgm.hal.science/hal-03113284

Submitted on 18 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

- Lithium isotopes in the Loire River Basin (France):
 hydrogeochemical characterizations at two complementary scales
 Romain Millot and Philippe Négrel
 BRGM, F-45060 Orléans, France
- 5 * Corresponding author : r.millot@brgm.fr

6 Abstract

7 We illustrate two different, but complementary, applications of lithium (Li) isotope tracers for 8 river-basin characterization at two different scales within the Loire River basin (LRB) in France. 9 The first example deals with the behaviour of Li and the fractionation of its isotopes during river weathering at the basin scale of the LRB (117,800 km²). The wide δ^{7} Li range (+5.0 to +13.3‰) in 10 11 Loire basin streams, spatialized between the headwaters and the lowlands, is consistent with 12 distinct weathering conditions and the distribution of Li from bedrock in the basin between the 13 rivers and secondary mineral phases during water/rock interaction. Additionally, suspended sediments in the LRB streams are significantly ⁶Li enriched (δ^7 Li from -8.7 to -7.6‰) compared to 14 15 average river waters that range from +5.0 to +13.3%. The second example focuses on the 16 smaller scale Egoutier watershed (13 km²), part of the LRB, which shows that Li isotopes can be 17 useful for distinguishing between natural input and anthropogenic pollution, like effluents from a 18 water treatment plant connected to a hospital. Overall, we confirm that Li isotopes cannot be 19 used as lithological tracers for river waters. However, we did find that Li isotopes can be good 20 tracers of weathering conditions and of anthropogenic sources in an urbanized watershed.

21

Keywords: lithium isotopes, river water, sediment, Loire River Basin, wastewater,
 anthropogenic input, natural spring

- 24
- 25

4872 words (without references or captions)

1- INTRODUCTION

In the Earth Sciences, the field of stable-isotope geochemistry of metals has greatly expanded over the past 20 years (Johnson et al., 2004; Teng et al., 2017), which is mainly due to the recent introduction of a new generation of Multi Collector Inductively Coupled Plasma Mass Spectrometers (MC-ICP-MS) with enhanced sensitivity and better precision. These instrumental innovations have opened new fields of research in metal-isotopes geochemistry (Albarède et al., 2004; Wiederhold, 2015).

34 Among the stable-isotope systematics of metals studied so far, the geochemistry of lithium (Li) has an excellent potential as a tracer of water/rock interactions within low- and high-35 36 temperature systems (Tomascak 2004; Burton and Vigier 2011; Tomascak et al., 2016; Penniston-37 Dorland et al., 2017), whether in the field of surface waters (Huh et al., 1998; 2001; Vigier et al., 38 2009; Wimpenny et al., 2010; Millot et al., 2010a; Dellinger et al., 2014; 2015), groundwaters (Hogan 39 and Blum 2003; Négrel et al., 2010; 2012; Meredith et al., 2013; Pogge von Strandmann et al., 2014; 40 Bagard et al., 2015), geothermal waters (Millot and Négrel, 2007; Millot et al., 2007; Millot et al., 41 2010b; Millot et al., 2011; 2012; Bernal et al., 2014), rainwaters (Millot et al., 2010c) or Li-rich brines 42 (Godfrey et al., 2013; Araoka et al., 2014).

43 Li isotopes are good tracers in geochemistry due to their involvement in water/rock 44 interactions at the Earth's surface. More precisely, Li and its isotopes can provide key information on 45 continental silicate weathering, which is the primary natural drawdown process of atmospheric CO₂ 46 and a major control on climate (Pogge von Strandmann et al., 2020). Li isotopes help our 47 understanding of weathering via globally important processes, such as clay formation and cation 48 retention. Both these processes occur as part of weathering in surface environments, including 49 rivers, soil pore waters, and groundwater, but Li isotopes can also be used for tracking weathering 50 changes across major climate-change events (Misra and Froelich, 2012).

51 Furthermore, Li has a strategic importance for numerous industrial applications, including the 52 production of Li-ion batteries for mobile devices and electric vehicles (World Economic Forum

Report, 2019), or in pharmaceuticals used in the treatment of some mental diseases (Aral and
Vecchio-Sadus 2008; 2011).

Assessing the behaviour of Li and its isotopes during chemical weathering is important for a better definition of water/rock interactions at the Earth's surface and also for refining our understanding of the global Li cycle. Lithium (⁶Li ~ 7.5% and ⁷Li ~ 92.5%) is a fluid, mobile metallic element and, due to the large relative mass difference between its two stable isotopes, it is subject to significant low-temperature mass fractionation that provides key information on weathering processes.

Additionally, the contribution of human activities (industry, agriculture, and domestic inputs) becomes increasingly significant in the chemical composition of dissolved river load (Aral and Vecchio-Sadus 2008; 2011) as well as in that of soil (Négrel et al., 2019). Human factors thus act as an additional key process and, therefore, a mass-balance for the budget of catchments and river basins must also consider anthropogenic disturbance of Li (Choi et al., 2019; Négrel et al., 2020).

66 To date, both the magnitude of Li-isotopic fractionation associated with water/rock 67 interaction processes, and the factors controlling such fractionation, are not totally understood. However, both field- and experimental studies have shown that ⁶Li is preferentially retained by 68 secondary minerals during silicate weathering (Pistiner and Henderson 2003, Kisakürek et al. 2004, 69 70 Pogge von Strandmann et al. 2006, Vigier et al. 2009). Accordingly, the fractionation of Li isotopes 71 depends upon the extent of chemical weathering: strong fractionation seems to occur during 72 incipient and early weathering stages, while little fractionation is observed during more intense or 73 prolonged weathering in a stable environment, such as a low plain with long residence times; Huh et 74 al. 1998, 2001, Pogge von Strandmann et al. 2006, Millot et al., 2010a).

Li-isotope fractionation has been documented in numerous natural environments with experimental and natural data (Tomascak 2004, Burton and Vigier 2011; Tomascak et al., 2016; Penniston-Dorland et al., 2017; Pogge von Strandmann et al., 2020). It was shown that partial dissolution of basalt does not result in Li-isotope fractionation, but that granite dissolution can cause

such fractionation (Pistiner and Henderson 2003; Négrel and Millot, 2019). In addition, adsorption onto mineral surfaces can be another major mechanism of Li-isotopic fractionation in the hydrosphere (Wenshuai and Liu, 2020; Andrews et al., 2020). Moreover, Li is not a nutrient and does not participate in biologically mediated reactions, so no evidence of biological Li-isotope fractionation has been observed to date (Rudnick et al., 2004; Marriott et al., 2004).

Here, we illustrate two applications at complementary scales of Li isotopes used for riverbasin hydrogeochemical surveys. The first example deals with the behaviour of Li and its isotopes at the Loire basin (LRB) scale (117,800 km², France) for river water and sediment transported during river flow. The second example focuses on a smaller scale: the Egoutier watershed (13 km²), part of the LRB. The main objective of our work was to evaluate Li isotopes as 1) tracers of weathering conditions over a large river basin, and 2) tracers of anthropogenic input at the scale of a small watershed.

91

2- ANALYTICAL METHODS

92 2.1. Sampling methods and element concentration measurements

In the LRB (Table 1), river waters were sampled twice, first during low flow and then during high flow (August/September 2012 and April 2013, respectively). Suspended sediments in the Loire at Montjean (the farthest downstream sampling station of this study, 955 km from the Loire spring) were sampled monthly between July 2012 and June 2013. For the Egoutier watershed, we collected river water in April 2015), corresponding to the high-flow stage of the watershed.

In the field, 5 litres of river water were collected in acid-washed containers for major- and trace-element and isotopic measurements (for details of the protocol see Millot et al., 2003).
Samples were filtered in the field, using a Sartorius frontal filtration unit (0.2 µm cellulose acetate filter, 142 mm diameter). After filtration, samples were stored in acid-washed polypropylene bottles.
The river sediments were collected after drying and centrifugation of the river-water filtration retentate. The samples for Li-concentration and Li-isotope analyses were acidified to pH=2 with ultrapure HNO₃. Lithium concentrations in river-water and suspended-sediment samples were

determined by Quadrupole ICP-MS (Thermo X Series II) with indium as an internal standard, with a
 precision of ±5%.

107 **2.2 Lithium isotope measurements**

108 Lithium-isotopic compositions were measured using a Neptune⁺ Multi-Collector ICP-MS at the BRGM (French Geological Survey; Millot et al., 2004). ⁷Li/⁶Li ratios were normalized to the L-SVEC standard 109 110 solution (NIST SRM 8545, Flesch et al., 1973) following the standard-sample bracketing method. Typical in-run precision on the determination of δ^7 Li was about 0.1-0.2‰ ($2\sigma_m$, standard error of the 111 mean). Chemical separation of Li from the matrix before isotope analyses was done with a cationic 112 exchange resin (a single column filled with 3 mL of BioRad AG[®] 50W-X12 resin, 200-400 mesh) and 113 HCl acid (0.2N) rendering 30 ng Li. Blanks for the total chemical extraction were <30 pg Li, which is 114 negligible as this represents a blank/sample ratio of $<10^{-3}$. 115

Successful quantitative measurement of Li-isotopic compositions requires 100% Li recovery during laboratory processing. Therefore, frequent column calibrations were performed and repeated analysis of an L-SVEC standard processed through the columns showed that no isotope fractionation occurred as a result of the purification process.

The accuracy and reproducibility of the entire method (purification procedure + mass analysis) were also tested by repeated measurement of a seawater standard solution (IRMM BCR-403) after separation of Li from the matrix, giving a mean value of δ^7 Li +30.8 ±0.4‰ (2 σ , n=15) over the duration of the analyses. This mean value agrees with our long-term measurements (δ^7 Li +31.0 ±0.5‰, 2 σ , n=30, Millot et al., 2004) and with other values reported in the literature (see for example Carignan et al., 2004 and Tomascak 2004 for a data compilation).

For suspended sediments, a total digestion of 50 mg of crushed sample took place over 4 days at 100 °C in a closed beaker with a mixture of three ultrapure acids: 4 mL HF (23N), 1 mL HNO₃ (14N) and 0.1 mL HClO₄ (12N). The solution obtained was evaporated to dryness and 4 mL HCl (6N) was added and left for a further 4 days at 100 °C. Sample aliquots (30 ng Li) of the residue of the acid dissolution were then dissolved in 0.5 mL HCl (0.2N), before being placed in cation exchange columns for Li separation. Accuracy and reproducibility of the entire procedure for solid samples (dissolution + purification + mass analysis) were tested by repeated measurements of the JB-2 basalt standard (Geological Survey of Japan), which gave a mean value of δ^7 Li = +4.9‰±0.6‰ (2 σ , n=17), in good agreement with published values (see Jeffcoate et al. 2004, Tomascak 2004 and Carignan et al. 2007 for data compilation).

136

3- FIELD SITE DESCRIPTION

137 **3.1 The Loire River Basin**

We undertook a systematic study of the weathering products (both dissolved load and suspended sediments) of the LRB, one of the major river basins in Europe. This area is of particular interest as it has been extensively studied and is scientifically well characterized (Négrel and Grosbois 1999; Grosbois et al., 2001).

142 The Loire River in central France (Fig. 1) is approximately 1010 km long and drains an area of 117,800 km². Initially, the Loire flows north to northwest, originating in the Massif Central and 143 continuing up to the city of Orléans, about 650 km from the source. Beyond Orléans, the river turns 144 145 west to WSW, being one of the main European riverine inputs into the Atlantic Ocean. In the upper 146 basin, the bedrock is Palaeozoic plutonic rock overlain by sub-Recent volcanic rocks. The 147 intermediate basin includes three major tributaries flowing into the Loire from the left bank: the 148 Cher, the Indre and the Vienne rivers. Here, the Loire drains sedimentary rocks of the Paris Basin, 149 mainly carbonate deposits. The lower Loire basin drains Palaeozoic basement rocks of the Armorican 150 Massif and its overlying Mesozoic to Cenozoic sedimentary deposits.

151 Its spring is located at an altitude of 1,404 m (Mont Gerbier de Jonc) and its course is 152 classically divided into three parts from upstream to downstream: (i) the Upper Loire, (ii) the Loire 153 Valley, and (iii) the Lower-Loire, to the estuary. River discharge of the Loire is very irregular: in the 154 Lower Loire it can exceed 7000 m³/s, but at Orléans it is around 350 m³/s and during summer the 155 river can literally dry up, with flows less than 25 m³/s.

BRGM's expert knowledge of the LRB is recognized by the numerous, particularly isotopic, studies, that have been carried out over the past decades (Négrel, 1997; Négrel and Grosbois, 1999; Négrel et al., 2000; Négrel and Petelet-Giraud, 2012; Petelet-Giraud et al., 2018). The Loire has been regularly monitored at Orléans and Tours over several hydrological cycles (Grosbois et al., 2000; 2001). Its physico-chemical parameters, major- and trace elements, and strontium-isotope ratios were determined on the dissolved fraction over time spans ranging from two days to one week, depending on river flow.

163 The relationships between chemical elements and flow as well as between isotopic ratios and 164 flow have shown that the dissolved fraction of the Loire results from a mixture between rainfall input, input from the weathering of silicate- and carbonate bedrocks, and anthropogenic input of 165 166 agricultural and urban origin. Total dissolved-solids flow during a hydrological cycle is estimated at 1300x10³ t/year at Orléans and 1620x10³ t/year at Tours (Grosbois et al., 2001). For the river 167 sediments, work by Négrel and Grosbois (1999) on ⁸⁷Sr/⁸⁶Sr isotope ratios suggests the existence of 168 169 at least two reservoirs of suspended matter transported by the river. One is related to the detrital 170 fraction from silicate erosion, the other to carbonate erosion. The stable isotopes of carbon and 171 oxygen in the fraction extracted by acid leaching from Loire sediments during periods of low water 172 flow (Négrel et al., 2000), confirm the formation of antigenic calcites in isotopic equilibrium with 173 Loire water (Fontes et al., 1973; Dever et al., 1983).

174 **3.2 The Egoutier watershed**

We also investigated anthropogenic Li tracing of wastewater release, using Li isotopes in the small
Egoutier catchment near Orléans (13 km², Fig. 2; in French, 'Egout' means 'Sewer', Desaulty and
Millot 2017).

Located in the Paris Basin on the Beauce plateau just east of Orléans, the Egoutier is a small stream with its spring near the village of Chanteau and a length of 5 km before being channelled and flowing into the Loire at Saint-Jean-de-Braye. For this study, we worked in a small built-up area

181 between its spring in the forest and a point just downstream of the departmental road D2060182 (Fig. 2).

The study area integrates two main potential sources of metals due to human activities: The first one is a psychiatric hospital with about 300 beds and many outpatients. The hospital also houses an active laundry. About a ton of linen from different community sources (fire brigade, schools, etc.) is washed and ironed every day. The second one is an industrial area including a slaughterhouse, factories for processing animal products, and the oil depot of Saint-Jean-de-Braye that supplies fuel oil to the Centre-Val-de-Loire region. This second area is drained by a stream that flows into the Egoutier.

190

4- RESULTS

191 **4.1 The Loire River basin**

In the LRB (Table 1), natural Li concentrations in river water are between 2.0 and 46.5 μ g/L whereas δ^7 Li-isotopic compositions range from +5.0 to +13.3‰. In more detail, there are slight differences depending upon whether the rivers were sampled during low- or high-flow stages.

195 During low flow, Li concentrations are higher (3.4 to 46.5 μ g/L; mean 15.9 μ g/L n=20, Table 1) than 196 during high flow (Li concentrations from 2.0 to 30.0 μ g/L with a mean of 7.4 μ g/L, n=20, Table 1). In 197 parallel, the Li-isotopic compositions are little different for low- and high-flow samples, with δ^7 Li 198 values ranging from +5.0 to +13.3‰ and +5.2 to +11.7‰ respectively. These concentrations are higher than the worldwide riverine average of 1.9 µg/L (Huh et al., 1998; Gaillardet et al., 2014). In 199 addition, Loire basin rivers have lower isotopic compositions than the average δ^7 Li value for modern 200 201 river water of about +23‰ (Huh et al., 1998; Misra and Froelich, 2012). Finally, δ^{7} Li values for the 202 Loire basin rivers agree with previous data reported by Rivé et al. (2013) for the upstream Loire (δ^7 Li 203 from 4.0 to 24.0%). Loire river suspended sediments collected monthly at Montjean/Loire (Table 2) from July 2012 to June 2013 have Li concentrations between 41.3 and 73.0 µg/g (mean 60.2 µg/g, 204 205 n=7), significantly higher than those of the average Upper Continental Crust (UCC) at Li $35 \pm 11 \mu g/g$ 206 $(2\sigma, \text{Teng et al.}, 2004)$. In addition, these sediments have low $\delta^7 \text{Li}$ with strongly negative values

207 ranging from -8.7 to -7.6‰ (mean 8.2‰, n=7) when compared to the UCC value of 0‰ ±2 (Teng et
 208 al., 2004).

209 4.2 The Egoutier watershed

Lithium concentrations in the Egoutier watershed cover a narrow range from 5.3 to 12.7 μ g/L (mean 7.3 μ g/L, n=9, Table 3), whereas the isotopic compositions of Li are comprised between -3.1 and +4.2‰ (mean +0.5‰, n=10, Table 3). The Li concentrations and the δ^7 Li values in the Egoutier watershed are very different from the ones observed at the larger scale of the LRB.

214

5- DISCUSSION

5.1 The Loire River basin: controlling parameters for the distribution of Li and its isotopes

Overall, Li concentrations in Loire River mainstream waters span a wide range from 2.0 to 22.5 μ g/L, whereas δ^7 Li values are between +5.9 and +13.2‰ (Table 1). Figures 3a and 3b show a clear contrast for the main course of the Loire River between headwaters and lowlands. This contrast shows a significant increase in Li concentrations from upstream to downstream (Fig. 3a). This feature is also associated with a strong decrease in the δ^7 Li values as a function of distance from the spring. These variations for Li concentration and δ^7 Li are observed for both high and low flow stages.

In addition, when all δ^7 Li values are plotted for both the Loire and its major tributaries (Cher, 222 223 Indre, Vienne, Maine, Furan, Arroux and Allier rivers, Fig. 1) in a δ^7 Li vs. Na/Li diagram (Fig. 4), the 224 data plot in a triangle formed by three endmembers. The first (Na/Li ~8000 and δ^7 Li ~+12‰) is the 225 upstream Loire at Villerest; the second endmember is close to the Allier river (Na/Li ~500 and δ^{7} Li 226 ~+5‰); and the third could be represented by rainwater. This agrees with the conceptual scheme of the Loire hydrosystem based on δ^{18} O and 87 Sr/ 86 Sr data, suggesting that the Loire River is related to a 227 228 Massif Central surface-water supply for the main Loire and Allier streams, and to water/groundwater 229 interactions in the alluvial plains (Négrel et al., 2003).

For the second endmember, the Allier constitutes a major tributary of the Loire that mostly drains the French Massif Central. As a result, its chemical (Na/Li) and isotopic signatures are influenced by input from thermo-mineralized waters resulting from hydrothermal activity, as 233 demonstrated by Négrel et al. (1997). These thermal springs are well described for the whole Massif 234 Central area and have low δ^7 Li values (Fig. 4, Millot et al., 2007). The third endmember, 235 corresponding to rainwater input, is in a good agreement with our long-term monitoring values of 236 rainwater at Orléans and also near Clermont-Ferrand within the Massif Central, corresponding to the 237 continental signature of rainwaters in France (Millot et al., 2010c).

238 It is well known that rainfall on watersheds can supply an important fraction of dissolved 239 elements in river water (Meybeck, 1983). Atmospheric correction requires knowledge of the 240 chemical composition of rainwater (op. cit.; Négrel et al., 1993). The correction of atmospheric 241 contribution to water, for a given element Z, is estimated by reference to the Cl concentration, called 242 Cl_{ref}, multiplied by the Z/Cl ratio of rainwater. For the LRB and according to Grosbois et al. (2000), we 243 considered the Cl_{ref} at 2.62 µg/L (74 µmol/L). We applied the atmospheric correction to Na, Li and 244 δ^{7} Li values, as earlier done for other studies on Li isotopes (Dellinger et al., 2015; Négrel and Millot, 245 2019; Négrel et al., 2020). Here, we used the characterization of rainwater defined by Négrel and Roy 246 (1998), Négrel et al. (2007) and Millot et al. (2010c), who reported major ions, but also Li 247 concentrations and isotopes (Millot et al., 2010c). The different rainfall stations in France range from near-ocean locations (Brest, Dax) to more continental ones (Orléans). The latter has the lowest Li 248 concentrations (0.37 μ g/L) and δ^7 Li values (+16.1‰). In view of its continental character, this station 249 250 is certainly the one that can best match the rain input over the LRB. Table 1 shows that Li in waters 251 from atmospheric origin ranges from 1.2 to 16.8% and from 1.9 to 28.2%, respectively for low- and 252 high-flow stages.

253 Once atmospheric correction was applied to lithium and its isotopes (Table 1; Fig. 5), the Li-254 isotopic compositions of the Loire River were spatialized along the main course. Figure 5 shows δ^7 Li 255 values plotted against Li concentrations: we see two parallel lines, one for high flow stages and the 256 second for low flow ones. The two lines are not interpreted as a mixing scheme for which the trends 257 may not be linear in such a representation. Nor could this feature be interpreted as a result of a 258 lithological contrast between the upstream and downstream Loire River (plutonic/volcanic rocks *vs.* 259 sedimentary deposits), in agreement with the results of Kısakürek et al. (2005), Vigier et al (2009) and

260

Millot et al. (2010a) that showed that Li isotopes cannot be used as lithological tracers in river basins.

261 This general picture might thus be better explained in terms of water/rock interaction: in the upstream environments of the Loire basin, short water-residence times and rock weathering may 262 263 enhance processes that fractionate Li isotopes, such as adsorption and/or secondary-mineral-phase formation, producing high δ^7 Li in rivers (Négrel and Millot, 2019). By contrast, in the lowlands of the 264 265 Loire basin, longer water-residence time could enhance Li dissolution, with lower isotopic fractionation (low δ^7 Li) approaching that of continental bedrock. Such a Li-isotope distribution has 266 267 already been reported for other large river basins, worldwide (Lemarchand et al., 2010; Millot et al., 2010b; Liu et al., 2013; 2015; Dellinger et al., 2014; 2015; Wang et al., 2015). 268

More specifically, the Li-isotopic composition measured in the Loire basin rivers shows that the dissolved load is significantly enriched in ⁷Li when compared to suspended river load. Lithium concentrations in suspended Loire sediment at Montjean (Table 2) contain 41 to 73 μ g/g and are clearly ⁶Li enriched; δ^7 Li values being negative (-8.7 to -7.6‰), compared to dissolved loads (+5.0 and +13.0‰). This result agrees with the fact that ⁶Li is preferentially incorporated into suspended sediment during weathering as also observed for other river basins worldwide (Huh et al. 1998, 2001, Pogge von Strandmann et al. 2006, Millot et al., 2010b).

Finally, the relationship between δ^7 Li and Al/Li in the suspended sediment (Fig. 6) could also raise the question of the control by different mineral phases (clay minerals *vs.* oxy-hydroxide of Fe or Mn, or carbonates) during Li-isotopic fractionation between water and solids. The variation of the Al/Li ratio could reflect a compositional variation of clay minerals in suspended sediments. It is also likely that the suspended sediments consist of both fine (clay rich) and coarse (carbonate rich) factions.

282 Unfortunately, we do not dispose over specific mineral-characterization data on our samples 283 for further discussion of this point, but it is interesting to observe an inverse correlation between the 284 Li-isotope signatures of suspended sediments and the river discharge (Fig. 7). This trend could

possibly suggest at least two different sources for suspended sediments, one during high flow and
the other during low flow, agreeing with the results from Sr-isotopic ratios of the Loire sediments
(Négrel et al., 2000; Négrel and Roy, 2002).

A final point that might explain such a relationship could be that the δ^7 Li content in solids may reflect weathering intensity (i.e., soil formation/transformation, coarse/fine material), as argued by Dellinger et al. (2014; 2015). Consequently, different pools of river sediments are likely to be transported as a function of the river regime.

292 **5.2** Anthropogenic origin of Li in the Egoutier watershed

Human activities such as industry, agriculture and domestic inputs, generally increase the quantity and modify the quality of chemical compounds in the dissolved and suspended load of rivers (Viers et al., 2009; Royer, 2016; Vanwalleghem et al., 2017). Human factors can also act as another key process for Li in rivers (Choi et al., 2019; Négrel et al., 2020). Therefore, the mass-balance for the budget of catchments and river basins should include anthropogenic disturbance.

Here, we investigate the effect of wastewater release by tracing its impact through using Li isotopes in the small Egoutier watershed (13 km², 5 km long). As a case study, we studied this small watershed with a low housing density in the LRB (Ledieu et al., 2020). Its spring is located in a pristine forested area, but some kilometres downstream it is affected by metal-rich effluents from a hospital water-treatment plant as well as by input from an industrial area (Fig. 2, Desaulty and Millot, 2017).

Within the course of the Egoutier, we clearly see the impact of both human activities. When Li concentrations are plotted as a function of distance from the spring (Fig. 8a), we see two different inputs: one from the waste water treatment plant (WWTP) related to the psychiatric hospital and the second from the stream draining the industrial area.

307 Whereas the natural background for Li in river waters is 5 to 6 μ g/L, the values for the 308 Hospital WWTP release and the industrial area are 12.7 μ g/L and 9.4 μ g/L, respectively. In addition, 309 Li-isotopic compositions are rather homogeneous in water of the Egoutier watershed, with δ^7 Li 310 values of around +0.5‰±1.2 along the main course of the stream (n=7, grey arrow on Fig. 8b), but

the signatures of the hospital and industrial area releases are very different. The WWTP water has a positive δ^7 Li value of +4.2‰, whereas the stream draining the industrial area has a negative value at -3.1‰. The two anthropogenic signatures are quite different for Li isotopes. As mentioned before, the hospital is a psychiatric one and it is likely that the Li released by the WWTP mainly comes from pharmaceutical products, such as Li carbonate used for bipolar disease treatment (Machado-Vieira et al., 2009). The nearby industrial area is mostly impacted by effluents from a slaughterhouse and animal-processing operations (Desaulty and Millot, 2017).

These observations in the Egoutier watershed completely agree with recent work by Choi et al. (2019), who investigated the impact of anthropogenic input on lithium content in river- and tap waters within the metropolitan area of Seoul, South Korea. They showed that Li-enriched wastewater can affect tap waters. They also report Li-isotopic signatures of both therapeutic drugs (Li carbonate) and detergents that are compatible with our findings for the wastewater treatment plant releases in the Egoutier, which combine waste from the laundry and the psychiatric hospital.

This example of the Egoutier watershed shows how Li isotopes can distinguish between natural and anthropic origins at the scale of a small watershed. This ability was recently evoked for Seoul (South Korea; Choi et al. 2019) as well as for the Dommel catchment (Belgium and Netherlands; Négrel et al., 2020). In both cases, lithium and its isotopes are effective tracers of wastewater releases in populated areas, as well as of industrial discharge (smelter effluents) into a river, respectively.

Further investigations are now needed to more precisely determine the Li-isotopic compositions of pharmaceutical formulations, but also the role of wastewater treatment processes in the distribution of metals between liquid and solid phases within a treatment plant. The latter point is particularly relevant, as Choi and al. (2019) showed that there is no significant difference between influent and effluent waters for both Li concentrations and Li isotope compositions, thus making this tracer a perfect conservative tool for tracing Li pollution in the environment.

This study can be considered as pioneering, since Li is found naturally in drinking water (Eyre-Watt et al., 2020; Ewuzie et al., 2020) with various origins. In clinical practice, it is widely used in the treatment of bipolar and of mood disorders. And, very importantly because of the increasing production of lithium-ion batteries worldwide (Naish et al., 2008; Conolly 2010; Christmann et al., 2015), Li pollution in surface- and groundwaters could become a major issue in coming years, as recently suggested by Choi et al. (2019).

342

6- CONCLUDING REMARKS AND FUTURE DIRECTIONS

A large variability of Li-isotopic ratios exists within the Loire river basin (LRB) and δ^7 Li is strongly spatialized between the Loire headwaters and lowlands. In addition, the Allier river, a major tributary in the French Massif Central, is clearly influenced by the contribution of thermo-mineral springs with lower δ^7 Li. Suspended river sediments are ⁶Li enriched (δ^7 Li from -8.7 to -7.6%) compared to river waters (+5.0 to +13.3%), agreeing with the fact that ⁶Li is preferentially incorporated in suspended sediments.

349 Overall, although the weathering mechanisms operating in the LRB must be defined in much 350 more detail, our work confirms that Li isotopes cannot be used as lithological tracers for river waters. 351 However, we also show that Li isotopes are good tracers of river weathering conditions, with an 352 innovative application for tracing anthropogenic Li sources in a small, urbanized watershed. Further 353 work is now required for better characterizing the chemical and isotopic signals of pharmaceutical 354 formulations containing Li as well as those of Li-ion batteries. More specifically, leaks from landfills 355 containing electronic products must be investigated for determining whether curative work is 356 needed to stop the spread of pollutants. To conclude, lithium and δ^7 Li are good proxies for studying 357 weathering and anthropogenic activities.

358

359 Acknowledgements:

This work was supported by the Research Program ISOP2 (Origin and mobility of metals in water and
 sediment of the Loire River Basin), cofounded by BRGM and the AELB Water Agency (Agence de l'Eau

362 Loire Bretagne). Special thanks are due to S. Perret for his help in sample preparation in the clean 363 laboratory and in the field. We acknowledge our fruitful discussions with A.M. Desaulty, X. Bourrain 364 and E. Petelet-Giraud. The BRGM Water, Environment, Process Development and Analysis Division is 365 acknowledged for funding. This study was part of the Service National d'Observation Environnements 366 Urbains: OBSERVIL (French Urban Observatory), accredited by the INSU/CNRS and on site certified by the Zone Atelier Loire: OBSCURE [OBservatoire des Sédiments et de la Cascade sédimentaire en 367 368 milieu URbain : cas de l'Egoutier à Semoy (Loiret)] managed by A. Simonneau. We thank three 369 anonymous reviewers and the Associate Editor for their constructive comments and remarks. 370 H.M. Kluijver edited the final English version of the MS for language and content.

371 RM is particularly grateful to T.D. (Tom) Bullen for helping him in the first stages of Li-isotope

investigations at BRGM, more than 17 years ago; this work is dedicated to his memory.

373

375 References

- Albarède F., Télouk P., Blichert-Toft J., Boyet M., Agranier A., and Nelson B. (2004). Precise and
 accurate isotopic measurements using multiple-collector ICPMS. Geochim. Cosmochim. Acta 68,
 2725-2744.
- 379 Andrews E., Pogge von Strandmann A.E., Fantle M.S. (2020). Exploring the importance of authigenic
- clay formation in the global Li cycle. Geochim. Cosmochim. Acta, 289, 47-68.
- Aral H., Vecchio-Sadus A. (2008). Toxicity of lithium to humans and the environment, A literature
 review. Ecotoxicology and Environmental Safety 70, 349-356.
- Aral H., Vecchio-Sadus A. (2011). Lithium: Environmental Pollution and Health Effects. Encyclopedia
 of Environmental Health. pp 499-508.
- Araoka D., Kawahata H., Takagi T., Watanabe Y., Nishimura K., Nishio Y. (2014). Lithium and
 strontium isotopic systematics in playas in Nevada, USA: constraints on the origin of lithium.
 Mineral. Deposita 49, 371–379.
- Bagard M.L., West A.J., Newman K., Basu A.R. (2015). Lithium isotope fractionation in the Ganges Brahmaputra floodplain and implications for groundwater impact on seawater isotopic
 composition. Earth Planet Sci. Lett. 432, 404–414.
- Bernal N.F., Gleeson S.A., Dean A.S., Liu X.M., Hoskin P. (2014). The source of halogens in geothermal
 fluids from the Taupo Volcanic Zone, North Island, New Zealand. Geochim. Cosmochim. Acta 126,
 265–283.
- Burton K.W., Vigier N. (2011). Lithium isotopes as tracers in marine and terrestrial environments
 Handbook of environmental isotope geochemistry. Advances in Isotope Geochemistry. Ed. Mark
 Baskaran. 2011, vol. 1, pp. 41-60.
- Carignan J., Cardinal D., Eisenhauer A., Galy A., Rehkämper M., Wombacher F., Vigier N. (2004). A
 reflection on Mg, Ca, Cd, Li and Si isotopic measurements and related reference materials.
- 399 Geostandards and Geoanalytical Research 28, 139-148.

- Carignan J., Vigier N., Millot R. (2007). Three secondary reference materials for Li isotopic
 measurements: 7Li-N, 6Li-N and LiCl-N. Geostandards and Geoanalytical Research 31, 7-12.
- 402 Choi H.B., Ryu J.S., Shin W.J., Vigier N. (2019). The impact of anthropogenic inputs on lithium content
- 403 in river and tap water. Nature Communications 10:5371 | https://doi.org/10.1038/s41467-019-
- 404 <u>13376-y</u>
- 405 Conolly D. (2010). A Review of Energy Storage Technologies: for the Integration of Fluctuating
 406 Renewable Energy. Aalborg Universitet. http://vbn.aau.dk/files/
- 407 Christmann P., Gloaguen E., Labbé J.F., Melleton, J., Piantone P. (2015). Global Lithium Resources and
 408 Sustainability Issues. Lithium Process Chemistry. Elsevier, pp. 1–40.
- 409 Dellinger M., Gaillardet J., Bouchez J., Calmels D., Galy V., Hilton R.G., Louvat P., France-Lanord C.
- 410 (2014). Lithium isotopes in large rivers reveal the cannibalistic nature of modern continental
 411 weathering and erosion. Earth Planet Sci. Lett. 401, 359–372.
- 412 Dellinger M., Gaillardet J., Bouchez J., Calmels D., Louvat P., Dosseto A., Gorge C., Alanoca L., Maurice
- 413 L. (2015). Riverine Li isotope fractionation in the Amazon River basin controlled by the weathering
- 414 regimes. Geochim. Cosmochim. Acta 164, 71–93.
- 415 Desaulty A.M., Millot R. (2017). Projet ISOP2 : Origine, mobilité et répartition eaux/sédiments des
- 416 métaux (Pb, Zn, Cu) : exemple de deux sous-bassins versants (Argos et Egoutier) du bassin Loire-
- 417 Bretagne. Rapport final. BRGM/RP-66799 -FR, 127 p.
- 418 Dever L., Durand R., Fontes J.C.H., Vachier P. (1983). Étude pédogénétique et isotopique des
 419 néoformations de calcite dans un sol sur Craie, Caractéristiques et origine. Geochim. Cosmochim.
 420 Acta 47, 2079-2090.
- Ewuzie U., Nnorom I.C., Eze S.O. (2020). Lithium in drinking water sources in rural and urban
 communities in South-eastern Nigeria. Chemosphere 245, 125593.
- 423 Eyre-Watt B, Mahendran E, Suetani S, Firth J, Kisely S, Siskind D. (2020). The association between
- 424 lithium in drinking water and neuropsychiatric outcomes: A systematic review and meta-analysis

425 from across 2678 regions containing 113 million . Australian & New Zealand Journal of Psychiatry.

426 October 2020. doi:10.1177/0004867420963740.

- Flesch G.D., Anderson A.R., Svec H.J. (1973). A secondary isotopic standard for ⁶Li/⁷Li determinations.
 International Journal of Mass Spectrometry and Ion Physics 12, 265-272.
- 429 Fontes J.C.H., Lepvrier C., Melieres F., Pierre C. (1973). Isotopes stables dans les carbonates

évaporitiques du Miocène Supérieur de Mediterranée Occidentale. In: Messinian Events in the

- 431 Mediterranean. Koninklijke Nederlandse Akademie van Wetenschappen, Amsterdam, pp. 91–100.
- Gaillardet J., Viers J., Dupré B., (2014). Trace Elements in River Waters. Treat. Geochem. (second ed.)
 7, 195–235.
- 434 Godfrey L.V., Chan L.H., Alonso R.N., Lowenstein T.K., McDonough W.F., Houston J., Li J., Bobst A.,
- Jordan T.E. (2013). The role of climate in the accumulation of lithium-rich brine in the Central
 Andes. Appl. Geochem. 38, 92–102.
- Grosbois C., Négrel Ph., Fouillac C., Grimaud D. (2000). Dissolved load of the Loire river: Chemical and
 isotopic characterization. Chem. Geol. 170, 179-201.
- Grosbois C., Négrel P., Grimaud D., Fouillac C. (2001) .An overview of dissolved and suspended
 matter fluxes in the Loire River Basin: Natural and anthropogenic inputs. Aqu. Geochem. 7, 81-
- 441 105.

- Hogan J.F., Blum J.D. (2003). Boron and lithium isotopes as groundwater tracers: a study at the Fresh
 Kills Landfill, Staten Island, New York, USA. Appl. Geochem. 18, 615–627.
- 444 Huh Y., Chan L.C., Zhang L., Edmond J.M. (1998). Lithium and its isotopes in major world rivers:
- implications for weathering and the oceanic budget. Geochim. Cosmochim. Acta, 62, 2039-2051.
- 446 Huh Y., Chan L.C., Edmond J.M. (2001). Lithium isotopes as a probe of weathering processes: Orinoco
- 447 River. Earth Planet Sci. Lett. 194, 189-199.
- 448 Jeffcoate A.B., Elliott T., Thomas A., Bouman C. (2004). Precise, small sample size determinations of
- 449 lithium isotopic compositions of Geological Reference Materials and modern seawater by MC-ICP-
- 450 MS. Geostandards and Geoanalytical Research, 28: 161-172.

- Johnson C.M., Beard B.L., Albarède F. (2004). Geochemistry of Non-Traditional Stable Isotopes
 Overview and General Concepts. Reviews in Mineralogy & Geochemistry 55, 1-24.
- 453 Kisakürek B., Widdowson M., James R.H. (2004). Behaviour of Li isotopes during continental 454 weathering: the Bidar laterite profile, India. Chem. Geol. 212, 27-44.
- 455 Kısakürek B., James R.H., Harris N.B.W. (2005). Li and δ^7 Li in Himalayan rivers: Proxies for silicate 456 weathering? Earth Planet. Sci. Lett., 237, 387-401.
- Ledieu L., Simonneau A., Cerdan C., Négrel Ph., Laperche V., Grosbois C., Laggoun F. (2020).
 Geochemical insights into spatial and temporal evolution of sediments at catchment scale. Appl.
 Geochem. 104743.
- Lemarchand E., Chabaux F., Vigier N., Millot R., Pierret M.C. (2010). Lithium isotope systematics in a
 forested granitic catchment (Strengbach, Vosges Mountains, France). Geochim. Cosmochim. Acta
 74, 4612–4628.
- Liu X.M., Rudnick R.L., McDonough W.F., Cummings M.L. (2013). Influence of chemical weathering on
 the composition of the continental crust: Insights from Li and Nd isotopes in bauxite profiles
 developed on Columbia River Basalts. Geochim. Cosmochim. Acta 115, 73–91.
- Liu X.M., Wanner C., Rudnick R.L., McDonough W.F. (2015). Processes controlling delta Li–7 in rivers
 illuminated by study of streams and groundwaters draining basalts. Earth Planet. Sci. Lett. 409,
 212–224.
- Machado-Vieira R., Manji H.K., Zarate C.A. (2009). The role of lithium in the treatment of bipolar
 disorder: convergent evidence for neurotrophic effects as a unifying hypothesis. Bipolar Disord.
 11, 92-109.
- 472 Marriott C.S., Henderson G.M., Crompton R., Staubwasser M., Shaw S. (2004). Effect of mineralogy,
- 473 salinity, and temperature on Li/Ca and Li isotope composition of calcium carbonate. Chem. Geol.
- 474 212, 5-15.

475 Meredith K., Moriguti T., Tomascak P., Hollins S., Nakamura E. (2013). The lithium, boron and
476 strontium isotopic systematics of groundwaters from an arid aquifer system: Implications for
477 recharge and weathering processes. Geochim. Cosmochim. Acta 112, 20–31.

478 Meybeck M. (1983). Atmospheric inputs and river transport of dissolved substances. IAHS Publ. 141,
479 173–192.

480 Millot R., Gaillardet J., Dupré B., Allègre C.J. (2003). Northern latitude chemical weathering rates:
481 clues from the Mackenzie River Basin, Canada. Geochim. Cosmochim. Acta, 67, 1305-1329.

482 Millot R., Guerrot C., Vigier N. (2004). Accurate and high-precision measurement of lithium isotopes

in two reference materials by MC-ICP-MS. Geostandards and Geoanalytical Research 28, 153-159.

484 Millot R., Négrel Ph. (2007). Multi-isotopic tracing (δ^7 Li- δ^{11} B, ⁸⁷Sr/⁸⁶Sr) and chemical

geothermometry: evidence from hydro-geothermal systems in France. Chem. Geol. 244, 664-678.

486 Millot R., Négrel Ph., Petelet-Giraud E. (2007). Multi-isotopic (Li, B, Sr, Nd) approach for geothermal

reservoir characterization in the Limagne Basin (Massif Central, France). Appl. Geochem. 22, 23072325.

Millot R., Vigier N., Gaillardet J. (2010a). Behaviour of lithium and its isotopes during weathering in
the Mackenzie Basin, Canada. Geochim. Cosmochim. Acta, 74, 3897-3912.

Millot R., Scaillet B., Sanjuan B. (2010b). Lithium isotopes in island arc geothermal systems:
Guadeloupe, Martinique (French West Indies) and experimental approach. Geochim. Cosmochim.
Acta 74, 1852-1871.

494 Millot R., Petelet-Giraud E., Guerrot C., Négrel Ph. (2010c). Multi-isotopic composition (δ^7 Li- δ^{11} B- δ D-495 δ^{18} O) of rainwaters in France: origin and spatio-temporal characterization. Appl. Geochem. 25, 496 1510-1524.

Millot R., Guerrot C., Innocent C., Négrel Ph., Sanjuan B. (2011). Chemical, multi-isotopic (Li-B-Sr-U-HO) and thermal characterization of Triassic formation waters from the Paris Basin. Chem. Geol.
283, 226-241.

- 500 Millot R., Hegan A., Négrel Ph. (2012). Geothermal waters from the Taupo Volcanic Zone, New 501 Zealand: Li, B and Sr isotopes characterization. Appl. Geochem. 27, 677-688.
- 502 Misra S., Froelich P.N. (2012). Lithium isotope history of Cenozoic seawater: changes in silicate 503 weathering and reverse weathering. Science 335, 818–823.
- Naish C., McCubbin I., Edberg O., Harfoot M. (2008). Outlook of Energy Storage Technologies. Policy
- 505 Department Economic and Scientific Policy.
- 506 http://www.europarl.europa.eu/document/activities/cont/201109/20110906ATT26009/
- 507 Négrel Ph., Allègre C.J., Dupré B., Lewin E. (1993). Erosion sources determined from inversion of
- 508 major, trace element ratios and strontium isotopic ratio in river water: the Congo Basin case.
- 509 Earth Planet Sci. Lett. 120, 59–76.
- 510 Négrel Ph. (1997). Multi elements chemistry of Loire estuary sediments: Anthropogenic versus
- 511 natural sources. Estuar. Coast. Shelf Sci. 44, 395-411.
- 512 Négrel Ph., Fouillac C, Brach M. (1997). Occurrence of mineral water springs in the stream channel of
- the Allier River (Massif Central, France): chemical and Sr isotope constraints. J. Hydrol. 203, 143153.
- Négrel Ph., Roy S. (1998). Rain chemistry in the Massif Central (France). A strontium isotopic and
 major elements study. Appl. Geochem. 13, 941–952.
- 517 Négrel Ph., Grosbois C. (1999). Changes in chemical and ⁸⁷Sr/⁸⁶Sr signature distribution patterns of
- 518 suspended matter and bed sediments in the upper Loire river basin (France). Chem. Geol. 156,
- 519 231-249.
- 520 Négrel Ph., Grosbois C., Kloppmann W. (2000). The labile fraction of suspended matter in the Loire
- river (France): multi-element chemistry and isotopic (Rb-Sr and C-O) systematics. Chem. Geol.
 166, 271-285.
- 523 Négrel Ph., Roy S. (2002). Investigating the sources of the labile fraction in sediments from silicate-
- drained rocks using trace elements, and strontium and lead isotopes. Sci. Tot. Env. 298, 163-181.

- 525 Négrel Ph., Petelet-Giraud E., Barbier J., Gautier E. (2003). Surface water–groundwater interactions in
- 526 an alluvial plain: chemical and isotopic systematics. J. Hydrol. 277(3-4), 248-267.
- 527 Négrel Ph., Guerrot C., Millot R. (2007). Chemical and strontium isotope characterization of rainwater
- in France: influence of sources and hydrogeochemical implications. Isotope Environ. Health Stud.
 43, 179–196.
- Négrel Ph., Millot R., Brenot A., Bertin C. (2010). Lithium isotopes as tracers of groundwater
 circulation in a peat land. Chem. Geol. 276, 119–127.
- Négrel Ph., Millot R., Guerrot C., Petelet-Giraud E., Brenot A., Malcuit E. (2012). Heterogeneities and
 interconnections in groundwaters: Coupled B, Li and stable-isotope variations in a large aquifer

534 system (Eocene Sand aquifer, Southwestern France). Chem. Geol. 296, 83–95.

- 535 Négrel Ph., Petelet-Giraud E. (2012). Isotopic evidence of lead sources in Loire river sediment. App.
 536 Geochem. 27, 2019-2030.
- Négrel Ph., Millot R. (2019). Behaviour of Li isotopes during regolith development on a granitic
 bedrock (Massif Central, France): controls on the dissolved load of waters, saprolites, soils and
 sediments. Chem. Geol. 523, 121-132.

540 Négrel Ph., Ladenberger A., Reimann C., Birke M., Demetriades A., Sadeghi M., & the GEMAS Project

541 Team. (2019). GEMAS: Geochemical background and mineral potential of emerging tech-critical

- elements in Europe revealed from low-sampling density geochemical mapping. App. Geochem.
- 543 111, 104425.
- 544 Négrel Ph., Millot R., Petelet-Giraud E., Klaver G. (2020). Li and δ^7 Li as proxies for weathering and 545 anthropogenic activities: application to the Dommel River (Meuse basin). App. Geochem. 104674.
- Penniston-Dorland S., Liu X.M., Rudnick R.L. (2017). Lithium Isotope Geochemistry Reviews in
 Mineralogy and Geochemistry. 82, 165-217.
- 548 Petelet-Giraud E., Négrel Ph., Casanova J. (2018). Tracing surface water mixings and groundwater

549 inputs using δ^{18} O- δ^{2} H and 87 Sr/ 86 Sr isotope fingerprints at basin scale: the Loire River (France).

550 App. Geochem. 97, 279-290.

- Pistiner J.S., Henderson G.M. (2003). Lithium isotope fractionation during continental weathering
 processes. Earth Planet Sci. Lett. 214, 327-339.
- 553 Pogge von Strandmann P.A.E., Burton K.W., James R.H., van Calsteren P., Gislason S.R., Mokadem F.
- (2006). Riverine behaviour of uranium and lithium isotopes in an actively glaciated basaltic
 terrain. Earth Planet Sci. Lett. 251, 134-147.
- 556 Pogge von Strandmann P.A.E., Porcelli D., James R.H., van Calsteren P., Schaefer B., Cartwright I.,
- 557 Reynolds B.C., Burton K.W. (2014). Chemical weathering processes in the Great Artesian Basin:
- 558 Evidence from lithium and silicon isotopes. Earth Planet. Sci. Lett. 406, 24–36.
- 559 Pogge von Strandmann P.A.E., Kasemann S.A., Wimpenny J.B. (2020). Lithium and lithium isotopes in
- 560 Earth's surface cycles. Elements, 16, 253-258.
- 561 Rivé K., Rad S., Assayag N. (2013). Carbon sources and water-rock interactions in the Allier River,
 562 France. Procedia Earth Planet. 7, 713-716.
- Royer T.V. (2016). Human-Dominated Rivers and River Management in the Anthropocene. Book
 Chapter: Stream Ecosystems in a Changing Environment, 491-524.
- Rudnick R.L., Tomascak P.B., Njo H.B., Gardner L.R. (2004). Extreme lithium isotopic fractionation
 during continental weathering revealed in saprolites from South Carolina. Chemical Geology, 212:
- 567 45-57.
- 568 Teng F.Z., McDonough W.F., Rudnick R.L., Dalpé C., Tomascak P.B., Chappell B.W., Gao S. (2004).
- 569 Lithium isotopic composition and concentration of the upper continental crust. Geochim.
- 570 Cosmochim. Acta, 68, 4167-4178.
- 571 Teng F.Z., Dauphas N., Watkins J.M. (2017). Non-traditional stable isotopes: Retrospective and 572 Prospective Reviews in Mineralogy and Geochemistry 82, 1-26.
- 573 Tomascak P.B. (2004). Developments in the understanding and application of lithium isotopes in the 574 earth and planetary sciences. Reviews in Mineralogy & Geochemistry 55, 153-195.
- 575 Tomascak P.B., Magna T, Dohmen R. (2016). Advances in Lithium Isotope Geochemistry. ISSN: 2364-
- 576 5105 Springer.

- Vanwalleghem T., Gómez J.A., Infante Amate J., González de Molina M., Giráldez J.V. (2017). Impact
 of historical land use and soil management change on soil erosion and agricultural sustainability
 during the Anthropocene. Anthropocene, 17, 13-29.
- 580 Viers J. Dupré B., Gaillardet J. (2009). Chemical composition of suspended sediments in World Rivers:
 581 New insights from a new database. Science of the Total Environment, 407, 853-868.
- 582 Vigier N., Gislason S.R., Burton K.W., Millot R., Mokadem F. (2009). The relationship between riverine
- 583 lithium isotope composition and silicate weathering rates in Iceland. Earth Planet Sci. Lett. 287,
 584 434-441.
- 585 Wang Q.L., Chetelat B., Zhao Z.Q., Ding H., Li SL, Wang B.L., Li J., Liu X.L. (2015). Behavior of lithium
- isotopes in the Changjiang River system: Sources effects and response to weathering and erosion.
- 587 Geochim. Cosmochim. Acta 151, 117–132.
- Wenshuai L., Liu X.M. (2020). Experimental investigation of lithium isotope fractionation during
 kaolinite adsorption: Implications for chemical weathering. Geochimica et Cosmochimica Acta,
 284: 156-172.
- 591 Wiederhold J.G. (2015). Metal stable isotope signatures as tracers in environmental geochemistry.
 592 Environ. Sci. Technol. 49, 2606–2624.
- 593 Wimpenny J., James R.H., Burton K.W., Gannoun A., Mokadem F., Gíslason S.R. (2010). Glacial effects
- 594 on weathering processes: New insights from the elemental and lithium isotopic composition of
- 595 West Greenland rivers. Earth Planet Sci. Lett. 290, 427-437.
- World Economic Forum (2019). A Vision for a Sustainable Battery Value Chain in 2030. Unlocking the
 Full Potential to Power Sustainable Development and Climate Change Mitigation.
 https://www.weforum.org/reports/a-vision-for-a-sustainable-battery-value-chain-in-2030
- 599

601 Figure and Table Captions

- Figure 1: Map showing the Loire River Basin within France (right). The geological map shows thedifferent sampling points for the Loire River mainstream and the different tributaries.
- 604 Figure 2: Map of the Egoutier watershed. 2a: General location of the sampling points, 2b: Land cover
- 605 map, 2c: Geological map. Modified from Desaulty and Millot (2017). The locations of the hospital and
- 606 the industrial area are shown on 2b.
- **Figure 3**: : Li concentrations (μ g/L, Fig. 3a) and Li isotopic compositions (δ^7 Li, ‰, Fig. 3b) in the Loire mainstream for low- and high-flow stages plotted as a function of distance from the source (km).
- 609 **Figure 4:** δ^7 Li (‰) plotted as a function of Na/Li mass ratio for the Loire River mainstream. Na
- 610 concentrations are from Desaulty and Millot (2017).
- **Figure 5:** δ^7 Li rainwater corrected (‰) plotted as a function of Li concentrations after rainwater correction (µg/L) for both low- and high-flow water samples. See text for explanation.
- **Figure 6:** δ^7 Li (‰) plotted as a function of the Al/Li mass ratio for suspended Loire sediments of the
- Loire at Montjean. Al data are from Desaulty and Millot (2017). (Determination coefficient R² = 0.60,
- 615 Correlation Coefficient R = 0.78).
- **Figure 7:** Li isotopic compositions (δ^7 Li, ∞) in the Loire River sediments at Montjean as a function of
- 617 river discharge (m^3/s) (Coefficient of determination $R^2 = 0.48$, Correlation coefficient R = 0.69).
- **Figure 8**: Li concentrations (μ g/L, Fig. 8a) and Li isotopic compositions (δ^7 Li, ‰, Fig. 8b) in Egoutier waters plotted as a function of distance from the source (km).
- 620

621 Table 1:

- 622 Major elements (Na, Cl), trace (Li) concentrations and Li-isotopic compositions (δ' Li, ∞) of Loire 623 waters and its main tributaries, at both high- and low-flow stages. The * means after atmospheric
- 624 correction (see text for explanations). River discharge data are also reported here.
- 625 Table 2:

- 626 Aluminium and Lithium concentrations and Li-isotopic compositions (δ^7 Li, ∞) of suspended river
- 627 sediments sampled at Montjean/Loire. River discharge data are also reported here.
- 628 Table 3:
- 629 Li concentrations and Li isotopic compositions (δ^7 Li, ∞) of Egoutier catchment waters.
- 630





635

636

.















Table 1

ŝ	CI (low)	Li (low)	Li* (low)	8 ⁷ Li (low)	8 ⁷ Li* (low) d	ischarge (Iow)	_
_	mg/L	hg/L	hg/L	%	%o	m³/s	
	16.1	3.4	2.8	12.6	11.9	10.7	
	22.7	11.3	10.8	13.2	13.1	41.6	
	21.8	10.8	10.3	13.3	13.1	27.6	
	20.9	13.0	12.4	12.1	11.9	67.0	
	22.2	22.5	21.9	8.7	8.5	89.9	
	23.2	21.5	20.9	7.5	7.3	68.6	
_	19.3	21.9	21.3	7.8	7.6	60.4	
	20.2	18.4	17.8	7.2	6.9	60.4	
_	21.4	19.3	18.7	7.0	6.7	78.7	
	21.1	19.4	18.8	6.8	6.5	78.7	
	21.9	19.6	19.0	7.4	7.1	77.1	
	22.9	13.6	13.0	7.3	6.9	127.0	
	25.7	12.6	12.0	7.8	7.4	144.0	
	24.4	5.7	5.1	10.3	9.7	9.7	
	30.3	4.8	4.2	11.2	10.5	3.3	
	22.5	4.3	3.7	13.2	12.7	31.4	
	38.4	5.1	4.5	8.9	8.0		
	73.3	29.9	29.3	11.3	11.2	1.7	
	20.7	14.3	13.7	13.1	13.0	7.7	
_	25.1	46.5	45.9	5.0	4.9	37.1	

667 Table 2

River	Sampling location	Sampling date	AI	Li	δ ⁷ Li	discharge
			µg/g	µg/g	‰	m³/s
Loire	Montjean/Loire	July 2012	45779	56	-8.7	384
Loire	Montjean/Loire	August 2012	46525	51	-8.3	181
Loire	Montjean/Loire	September 2012	50735	59	-8.6	153
Loire	Montjean/Loire	April 2013	75021	68	-7.6	62.75
Loire	Montjean/Loire	May 2013	74611	73	-8.2	64.69
Loire	Montjean/Loire	June 2013	67308	73	-7.8	69.78
Loire	Montjean/Loire	July 2013	38580	41	-8.3	78.47

670 Table 3

River	Sampling location	Li	δ ⁷ Li	
		µg/L	‰	
Egoutier	Source	5.3	1.1	
Egoutier	point 2	8.4	2.9	
Egoutier	point 2 bis	12.7	4.2	
Egoutier	point 3	6.9	0.0	
Egoutier	point 4	5.7	0.3	
Egoutier	point 5	9.4	-3.1	
Egoutier	point 6	5.4	0.7	
Egoutier	point 7	5.8	-0.2	
Egoutier	end	5.9	-1.2	