

Influence of nutrients on bioleaching efficiency of polymetallic sulfidic mining residues

Agathe Hubau, Catherine Joulian, Douglas Pino Herrera, Anne-Gwenaelle

Guezennec

▶ To cite this version:

Agathe Hubau, Catherine Joulian, Douglas Pino Herrera, Anne-Gwenaelle Guezennec. Influence of nutrients on bioleaching efficiency of polymetallic sulfidic mining residues. Biomining'21, Jun 2021, online, United Kingdom. 10.1016/j.hydromet.2020.105484 . hal-03227569

HAL Id: hal-03227569 https://brgm.hal.science/hal-03227569

Submitted on 17 May 2021 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. Event: Biomining'21 conference

Title: Influence of nutrients on bioleaching efficiency of polymetallic sulfidic mining residues.

Authors: Agathe HUBAU, Catherine JOULIAN, Anne-Gwénaëlle GUEZENNEC BRGM - Water, Environment, Processes and Analyses Division - 3, Avenue Claude Guillemin – BP36009 – 45060 Orléans CEDEX 2 - FRANCE

Our study aimed at assessing the influence of nutrient medium on the bioleaching of a polymetallic and polymineral mining residue. Bioleaching experiments in 2 L-stirred tank reactors maintained at 55°C with 20% solid concentration, in the presence of variable concentrations of ammonium and potassium, were compared. Increasing nutrient concentration enhanced the dissolution yields of Co and sulfides, suggesting a decrease of the microbial activity with nutrients-poor media. Ni, Zn, and Cu yields remained the same, which confirms results from a previous study showing their dissolution is mainly chemical [1]. The composition of the nutrient medium also highly affected the precipitation phenomena, thus modifying the solid-liquid separation and the acid demand. Changes in biomass concentration were correlated to nutrients consumption and showed lower ammonium concentration was detrimental to biomass growth, while lowering potassium concentration did not limit the growth but rather the maintenance of the cells during the stationary phase.

[1] Hubau, A., Guezennec, A.-G., Joulian, C., Falagán, C., Dew, D., Hudson-Edwards, K.A., 2020.
Bioleaching to reprocess sulfidic polymetallic primary mining residues: Determination of metal leaching mechanisms. Hydrometallurgy 197, 105484. <u>https://doi.org/10.1016/j.hydro-met.2020.105484</u>