



HAL
open science

Potential safe geological storage of CO₂ as mixed gas hydrates along the European continental margins

André Burnol, Ricardo Leon

► **To cite this version:**

André Burnol, Ricardo Leon. Potential safe geological storage of CO₂ as mixed gas hydrates along the European continental margins. ECGH 2022 – European Conference on Gas Hydrate, Jun 2022, Lyon, France. hal-03660225

HAL Id: hal-03660225

<https://brgm.hal.science/hal-03660225v1>

Submitted on 5 May 2022


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Potential safe geological storage of CO₂ as mixed gas hydrates along the European continental margins

André Burnol¹, Ricardo León Buendía²
¹ BRGM, Orléans, a.burnol@brgm.fr
² IGME, Madrid, r.leon@igme.es

A pan-European GIS focused on gas hydrates has been developed in the GARAH project (<https://geoera.eu/projects/garah4/>) to assess for the first time the gas hydrate related risk along the European continental margins. Several factors and variables have been taken into account to assess the abundance of sediment-hosted gas hydrates and to estimate the safe storage volume of CO₂ as mixed gas hydrates in deep-sea sediments. The baseline scenario is that gas hydrate occurrence is only possible in seafloor areas where pressure (bathymetry) and seafloor temperature conditions are inside the theoretical gas hydrate stability zone (GHSZ) and that the injected CO₂(l) is expected to percolate downwards or upwards to the neutral buoyancy level (NBZ) depending on the CO₂ quality and on the seafloor depth. In the Bay of Biscay, the safe CO₂ deep offshore storage capacity is estimated in both cases (pure CO₂ or 96%ml CO₂). The French EEZ storage volume estimate of 3,422 km³ is of the same order of magnitude as the total Spanish EEZ storage volume estimate of 3,700 km³. There is, however, a big difference resulting from the seafloor depth: in the French zone, almost all the stored volume is occupied by gas hydrates, whereas in the Spanish zone, about the half (1,728 km³) is occupied by CO₂ in hydrate phase and the other half (1,972 km³) by liquid CO₂. This GIS-database has been conceived as a first step or base-line for future gas hydrate related research.



Acknowledgment GARAH project. GeoERA - GeoE.171.00

- Candidature to the best PhD poster presentation
- Candidature to the best Young Researcher oral presentation (PhD student or young researcher – PhD defended < 5 years)
- I want to deposit (end of June, I will be contacted by reviewers) a full document for reviewing and publication