



HAL
open science

Abandoned Mine Lands Reclamation by Plant Remediation Technologies

Jesús D Peco, Pablo Higuera, Juan A Campos, José M Esbrí, Marta M
Moreno, Fabienne Battaglia-Brunet, Luisa M Sandalio

► **To cite this version:**

Jesús D Peco, Pablo Higuera, Juan A Campos, José M Esbrí, Marta M Moreno, et al.. Abandoned Mine Lands Reclamation by Plant Remediation Technologies. Sustainability, 2021, 13 (12), pp.6555. 10.3390/su13126555 . hal-03745740

HAL Id: hal-03745740

<https://brgm.hal.science/hal-03745740>




Submitted on 4 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Review

Abandoned Mine Lands Reclamation by Plant Remediation Technologies

Jesús D. Peco ^{1,2,*} , Pablo Higuera ² , Juan A. Campos ¹, José M. Esbrí ² , Marta M. Moreno ¹, Fabienne Battaglia-Brunet ³ and Luisa M. Sandalio ^{4,*}

- ¹ Escuela Técnica Superior de Ingenieros Agrónomos, Universidad de Castilla-La Mancha (UCLM), 13071 Ciudad Real, Spain; juanantonio.campos@uclm.es (J.A.C.); martamaria.moreno@uclm.es (M.M.M.)
- ² Instituto de Geología Aplicada, Universidad de Castilla-La Mancha (UCLM), 13400 Ciudad Real, Spain; pablo.higuera@uclm.es (P.H.); josemaria.esbri@uclm.es (J.M.E.)
- ³ Bureau de Recherches Géologiques et Minières (BRGM), 3 Avenue Claude Guillemin, 45060 Orléans, France; f.battaglia@brgm.fr
- ⁴ Departamento de Bioquímica, Biología Celular y Molecular de Plantas, Estación Experimental del Zaidín, (CSIC), 18008 Granada, Spain
- * Correspondence: jesusdaniel.peco@uclm.es (J.D.P.); luisamaria.sandalio@eez.csic.es (L.M.S.)

Abstract: Abandoned mine lands (AMLs), which are considered some of the most dangerous anthropogenic activities in the world, are a source of hazards relating to potentially toxic elements (PTEs). Traditional reclamation techniques, which are expensive, time-consuming and not well accepted by the general public, cannot be used on a large scale. However, plant-based techniques have gained acceptance as an environmentally friendly alternative over the last 20 years. Plants can be used in AMLs for PTE phytoextraction, phytostabilization, and phytovolatilization. We reviewed these phytoremediation techniques, paying particular attention to the selection of appropriate plants in each case. In order to assess the suitability of plants for phytoremediation purposes, the accumulation capacity and tolerance mechanisms of PTEs was described. We also compiled a collection of interesting actual examples of AML phytoremediation. On-site studies have shown positive results in terms of soil quality improvement, reduced PTE bioavailability, and increased biodiversity. However, phytoremediation strategies need to better characterize potential plant candidates in order to improve PTE extraction and to reduce the negative impact on AMLs.

Keywords: phytoremediation; phytostabilization; phytovolatilization; phytoextraction; abandoned mine lands; heavy metals; reclamation; oxidative stress; accumulation



Citation: Peco, J.D.; Higuera, P.; Campos, J.A.; Esbrí, J.M.; Moreno, M.M.; Battaglia-Brunet, F.; Sandalio, L.M. Abandoned Mine Lands Reclamation by Plant Remediation Technologies. *Sustainability* **2021**, *13*, 6555. <https://doi.org/10.3390/su13126555>

Academic Editors: Iyyakkannu Sivanesan and Muthu Thiruvengadam

Received: 13 April 2021

Accepted: 6 June 2021

Published: 8 June 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In recent decades, many countries have realized that abandoned metal mining operations greatly contribute to environmental degradation [1]. Metal mining radically transforms the natural environment, causing modifications to the land, as well as increased accumulation of potentially toxic elements (PTEs) in the ecosystem, which seriously undermines soil health and the viability of all living organisms [2]. The principal problem is caused by mining activity carried out in the past using inefficient technology in the absence of environmental protection regulations, leading to the accumulation of mining waste which is now a source of PTEs [3]. Regarding to their role in biological systems, PTEs can be divided into essential and non-essential categories. Despite being required by living organisms for physiological and biochemical functions, at high concentrations, essential PTEs, such as copper (Cu), nickel (Ni), iron (Fe), and zinc (Zn), can have a toxic environmental impact. On the other hand, living organisms do not require non-essential PTEs such as lead (Pb), mercury (Hg), cadmium (Cd), arsenic (As), chromium (Cr) and antimony (Sb) [4]. Large concentrations of PTEs present in abandoned mine waste can migrate to nearby lands and cause severe agricultural soil and water pollution and undermine food safety

and human health [5]. The 2030 Agenda for Sustainable Development, which includes 17 Sustainable Development Goals (SDGs), also refers to this problem. SDG target 15.3, in particular, aims, *by 2030, to combat desertification, restore degraded land and soil, including land affected by desertification, drought and floods, and to strive to achieve a land degradation-neutral world* [6]. The achievement of these objectives in the near future would also necessitate the identification, implementation, and optimization of AML reclamation strategies.

Over the last ten years, the following remediation strategies have been developed to reclaim abandoned mine lands (AMLs): (I) surface stabilization to prevent eolian and water erosion; (II) prevention of toxic pollutant dispersion to the surrounding environment; and (III) restoration of the degraded natural landscape [7]. AMLs can be treated in situ using the following physico-chemical techniques: soil isolation and containment, solidification and stabilization, vitrification, soil flushing, and electrokinetic reclamation [8]. However, these methods, which are expensive, time-consuming and unacceptable to the general public, cannot be applied on a large scale [9]. On the other hand, many organic and inorganic materials, which can be used as in-situ alternatives to expensive and disruptive physico-chemical techniques, can potentially be used as soil amendments [10]. These organic amendments include wood ash, manure, sawdust, compost, sewage sludge, biosolids, and wood chips. Commonly used inorganic amendments include phosphate-derived products, liming materials, alkaline sludge, and sugar foam [11]. However, despite their productive results, these techniques do not eliminate PTEs from the soil.

Plant-based techniques can be used to restore degraded soil in a cost-effective and environmentally friendly manner compared with physico-chemical processes [12]. While physico-chemical techniques irreversibly alter soil properties, phytoremediation generally improves the physical, chemical, and biological quality of mining lands [13]. In addition, according to Wan et al. [14], the total cost of a two-year project devoted to the phytoremediation of soil contaminated by As, Cd, and Pb was estimated to be US\$ 75,375.2/hm² or US\$ 37.7/m³, which is significantly lower than that for most other techniques studied. Phytoremediation is defined as the use of living plant species to clean up soil, water and air that have been polluted with hazardous contaminants [15]. The following phytoremediation strategies are principally used to restore AMLs: in situ phytostabilization as well as phytoextraction and phytovolatilization clean-up techniques (Figure 1) [16]. These strategies involve selecting suitable plant species that can be used to decontaminate polluted soils. A knowledge of the mechanisms involved in natural plant tolerance to PTEs is therefore crucial in order to design new phytoremediation strategies [17–19]. The accumulation of and tolerance to PTEs involve many different genes, whose selection and transfer to candidate plants constitute a potential genetic engineering strategy to obtain new organisms with characteristics more suited to environmental remediation [20,21]. It is important to note that, apart from the need to remove human-made structures and PTEs in AML reclamation projects, it is also necessary to restore the biodiversity [3]. Thus, simultaneously or following phytoremediation, revegetation with native flora suited to the specific ecosystem is required in order to return AMLs to their pre-mining state. However, for natural colonization by native plant species to occur in most AMLs, a prior reclamation process is essential [6].

In recent years, although considerable progress has been made in the application of phytoremediation techniques, significant additional research needs to be carried out. In this review, we discussed the current state of research into AML phytoremediation techniques, analyzed some actual examples, and determined what areas require further research in the future. Particular attention should be paid to the search for appropriate wild plants for use in phytoremediation practices.




Figure 1. Principal mechanisms involved in phytoremediation.

2. Occurrence of AMLs and Their Ecological Impact Worldwide

Throughout history, mining activity has produced an enormous number of impacted sites, many of which have undergone remediation, restoration, and reclamation, with differing levels of success, aimed at reducing its environmental impact. However, too often, ancient mining sites, such as those in Italy, Spain, and Latin America, were worked intensely and abandoned without any restorative measures being carried out. More recently, during the first half of the 20th century, very large French and British mining operations such as the Sao Domingos mine in Portugal [22], the Tharsis mine in Spain [23], and the Salafossa mine in Italy [24] left a legacy of abandonment. However, a comprehensive list of mining activities around the world, with different degrees of disruption, is too long for inclusion in this review. Most of these sites contain wastes accumulated in tailing dumps, which are

major sources of pollution for the surrounding environment [25] though aerial dispersion, percolation, water leakage, and acid mine drainage [26].

AML abandonment has had a diverse range of consequences for the local and regional environment, some of which have had catastrophic effects, particularly in the form of waste accumulation, direct mine residue dumps, and processed waste dams, which have passed through various mineral processing stages, including froth flotation, calcination, and fusion. Each stage produces modifications in the characteristics of the original accumulated materials. Some of these changes further mobilize chemical pollutants and hence maximize the release of chemicals into the human food chain. An important related issue is the degradation of soil health [27], which includes a reduction in certain soil enzyme activities, mainly dehydrogenase, as well as changes in bacterial communities [28].

3. Plant Remediation Technologies for AML Restoration

Over the last thirty years, the use of phytoremediation for AML regeneration has been gaining in importance. Plant-based remediation, which is eco-friendly, innovative, cost-effective, practical, accepted by the general public, and carried out in-situ, can increasingly be used for PTE-contaminated soils [29]. The use of plants on AMLs enables PTEs to be extracted and immobilized without damaging the soil structure while regenerating soil fertility and quality [16]. The potential negative effects of metal-tolerant and hyperaccumulator plants are offset by their efficient uptake and accumulation of toxic elements. This, together with their adaptation to most environmental conditions, makes these plants a versatile remediation tool [30]. However, the use of plants for AML remediation has certain limitations; with their low organic content, plant nutrient levels and pH, as well as high toxic element levels, mine tailings usually lack the appropriate edaphic characteristics to establish vegetation [31]. Consequently, phytoremediation, which involves the selection of suitable plants and their establishment in AMLs, could be slower or even a long-term process as compared with physico-chemical processes [32]. In addition, implementation initially requires AML evaluation, mineralogical and physico-chemical analysis, monitoring of pH extremes, salinity and erodibility, as well as plant nutrient and microbial community analyses (Figure 1) [28,33,34]. Another important factor to be taken into account is that phytoremediation, which most often involves AML revegetation, can be considered a reconstruction and reestablishment of the original ecosystem [35].

3.1. Clean-Up Techniques

Despite being thought of as the most effective and practical plant-based PTE removal technique [30,36], phytoextraction is the most difficult to carry out given the need to find good plant candidates [16]. This technique involves selecting plants that can grow in contaminated soils and of taking up large amounts of PTEs from the soil for final accumulation in the aerial part of the plant [30]. At the end of this process, plants are harvested and incinerated after drying, which generates concentrations of ash pollutants, which need to be properly disposed of, processed to recover the metals, or simply used for landfill, with their incineration potentially producing an alternative source of energy [37]. While the removal of PTE contamination from a two-acre site to a depth of 18 inches normally produces 5000 tons of contaminated soil, the ash produced by phytoremediation techniques would only amount to between 25 and 30 tons [38]. On the other hand, PTE-contaminated biomass combustion can be hazardous due to metal dispersion through flying ashes. However, this can be counteracted using scrubbers, filters, and electrostatic precipitators which remove most PTEs and flying ash from incinerator air emissions to comply with regulatory thresholds [39]. In addition to cleaning up toxic metal-contaminated areas, at the current world prices of some PTEs, phytoextraction could be an economically viable phytomining alternative [40]. Phytomining is a phytoextraction-based technology that uses hyperaccumulator plant capacity to extract valuable elements from metal-contaminated soils; this process produces bio-ore from the harvested biomass, which facilitates metal recovery through smelting [41]. Thus, mine tailings could be used as secondary sources of valuable

metals through the incineration of the harvested biomass to obtain bio-ore in much higher concentrations. Li et al. [42] reported that the 20,000 $\mu\text{g g}^{-1}$ Ni contained in *Alyssum murale* biomass became 32 wt.% in ash. Recently, the establishment of the first tropical 'metal farm', using the 'metal crop' species *Phyllanthus rufuschaneyi*, was reported in Malaysia [43]. Metal extraction from ash could, in turn, be carried out using different processes. For example, an efficient method to recover valuable elements from the ash of hyperaccumulating *Alyssum murale* plants by leaching the ashes using a sulphuric acid solution has been reported [44]. Other similar processes that could be used to recover metals from incineration ashes from solid municipal waste, including magnetic separation for ferrous compounds, Eddy current separation for non-ferrous metals, and X-ray transmission sorting for light/heavy metal separation [45], could also be applied to plant ashes. The benefits and limitations of phytoextraction were elegantly reviewed by Corzo et al. [40].

The most critical step in the implementation of phytoextraction is the selection of hyperaccumulator plants capable of accumulating at least 10 g kg^{-1} (1%) of Mn and Zn, 1 g kg^{-1} (0.1%) of As, Co, Cr, Cu, Ni, Pb, Sb, Se, and Tl, and 0.1 g kg^{-1} (0.01%) of Cd in the aerial parts of the plant without affecting its survival [16]. Research carried out on plant populations collected from soils severely contaminated with PTEs has identified many plant species with high PTE tolerance and accumulation capacities, although this is not the only requirement. According to Alkorta et al. [36], the appropriate plants for phytoextraction should have the following features: high tolerance to PTE exposure, PTE accumulation in the aerial parts of the plant, rapid growth rates, high biomass production, a proper root system, easy crop management, genetic stability, and non-edibility for animals. Table 1 provides a summary of potentially good hyperaccumulator candidates for use in phytoextraction strategies. However, these plant populations cannot easily compensate for all the contaminated AMLs, where hyperaccumulator plants growing on this land are highly sensitive to changes in prevailing soil and climate conditions. In addition, as the tolerance and accumulation mechanisms of these plants are usually PTE-specific, the presence of other contaminants could negatively affect plant growth [7,16]. Some studies have found that a plant's phytoextraction capacity is linearly correlated with the amount of PTEs in the soil, meaning that, after successive crops, phytoextraction ceases to be effective, with a certain amount of contaminants remaining in the soil [46].

As phytoextraction can be a very slow process as compared to other physico-chemical techniques [16], plant communities and high biomass generation capacity need to be established for one or more seasons, depending on whether herbaceous, shrub, or arboreal plants are used. The need for long periods of time to remediate polluted soils sometimes renders phytoextraction nonviable and impractical [16]. In order to accelerate the extraction process, the use of soil amendments, such as compost and shredded tree bark, with their higher seed germination rate and plant growth potential due to their enhanced physico-chemical soil properties and organic matter supply [31], as well as the use of chelators, which increase the plant metal absorption, have been proposed [47]. For example, the use of ethylenediamine disuccinic acid (EDDS) and vermicompost in *Helianthus annuus* L. exposed to Cd and Pb has notably been found to increase the extraction of the metals mentioned above [48]. Similarly, marble waste, when added to the soil, increased leaf Pb accumulation 2.5-fold and that of Cd by 55% in *Atriplex halimus* as compared with control plants [49]. On the other hand, despite the reported induced phytoextraction potential of ethylenediaminetetraacetic acid (EDTA) and diethylenetriaminepentaacetic acid (DTPA), these chelators exhibit poor biodegradability, which may trigger groundwater contamination [50,51]. This was demonstrated by Grčman et al. [47] in soil from a former Pb and Zn industrial smelter site. Despite its enhancement of PTE accumulation in the green parts of *Brassica rapa* L. var. *pekinensis*, they showed that the addition of EDTA also caused Pb, Zn, and Cd leaching through the soil and increased the toxic effects on test plants and soil microorganisms. Many studies have investigated the possibility of using other less toxic compounds for the environment than EDTA and DTPA. Meers et al. [52] evaluated whether the biodegradable chelator EDDS could be used to enhance

phytoextractive processes. EDDS, which initially sparked an intense mobilization of PTEs, followed by a sharp decrease due to its rapid degradation over a 3.8–7.5 day period, was judged as unsuitable for phytoremedial use. Similarly, nitrilotriacetic acid (NTA) and other organic acids were found to require continuous supplementation in large quantities due to their rapid biodegradation also [53]. Although the addition of citric acid, NH₄-citrate/citric acid, oxalic acid, EDDS, and NTA to *Brassica juncea* and *Lolium perenne* sharply increased PTE phytoextraction, their use was found to be impractical due to the time required to achieve acceptable levels of reduction in soil contaminant concentrations [54]. The goal of many studies to obtain compounds to facilitate absorption by the plant without any adverse effects [53,54] has yet to be achieved. Thus, improvements in phytoextraction techniques are currently based on appropriate plant selection and soil enhancement.

Phytovolatilization clean-up involves plant uptake of contaminants and their release into the atmosphere in a less toxic form [32]. This technique, which enables gaseous volatilized PTEs to be redeposited in surrounding lands, lakes, and oceans, can be precisely analyzed [15]. In particular, the most common PTEs, Hg, As, and Se, can be removed by their conversion into volatile forms [29]. Some interesting examples of plants with a phytovolatilization capability are shown in Table 2 and many more exist in the literature, some of which are described below. *Pteris vittata* has been shown to efficiently volatilize approximately 90% of total As uptake from contaminated soils in greenhouse experiments [55]. A bio-phytoremediation technique was tested for As remediation potential by adding *Stenotrophomonas maltophilia* and *Agrobacterium* sp. to the roots of *Arundo donax*. By the end of the experiment, ≈25% of As remained in the sand, ≈0.15% was accumulated in the plant, while the remaining 75% was volatilized by transpiration [56]. Another experiment with *Zea mays* exposed to Hg showed that plants inoculated with *Serratia marcescens* and *Pseudomonas* sp. increased mercury volatilization by 47.16% and 62.42%, respectively [57]. Although phytovolatilization could be used for effective AML remediation, prior evaluations of possible environmental risks linked to certain volatilized PTEs are required.

3.2. On-Site Stabilization Techniques

Phytostabilization, a remediation technique currently attracting increasing interest, involves the use of plant roots to limit PTE mobility and bioavailability in the soil [6]. Plants used for AML phytostabilization need to: (I) decrease the amount of water in the soil matrix, which prevents metals from leaching into aquifer systems; (II) prevent soil erosion and reduce PTE dispersion to other areas; (III) act as a barrier to reduce direct contact between animals and contaminated soils; (IV) sequester large amounts of PTEs through plant roots, thus reducing their bioavailability and leaching; and (V) to promote PTE immobilization in the rhizosphere through complexation/precipitation associated with the production of certain root exudates [16,58]. As with the other techniques described above involving PTE-tolerant plant selection, in the case of phytostabilization, the plant species selected also need a high root biomass production capacity, as well as root systems with PTE accumulation and uptake capabilities [59]. Most phytostabilization research has focused on the initial stage of selecting appropriate candidate plant species, some of which are listed in Table 3. These species accumulate large pollutant concentrations in the roots rather than in the aerial part of the plant, thus avoiding the drawbacks associated with phytoextraction, such as the final harvesting and the possible ingestion of the green parts and fruits by animals [7,36]. However, as these species have other drawbacks such as their inability to reduce the total amount of PTEs in the soil, plants with a long lifespan are preferable to ensure long-term phytostabilization [60].

In order to effectively establish appropriate plant communities to implement phytostabilization programs, the use of soil amendments has been proposed to improve nutritional status and soil structure [59]. The treatment of Pb/Zn mine tailings with Fe-rich compounds, furnace slag and red mud, combined with *Miscanthus sinensis* and *Pteridium aquilinum*, led to a sharp decrease in Pb, Cd, and Zn bioavailability following phytostabilization. Treatment with various amendments and the use of *M. sinensis* decreased the

mobility of Cd, Pb, and Zn to 72–92%, 83–90%, and 79–88% of control, respectively, while, with *P. aquilinum*, the decrease in Cd, Pb, and Zn mobility was 79–96%, 77–91%, and 77–96% of control, respectively [61]. Another study found that a combination of compost, cyclonic ash, and steel shots had a beneficial effect on the plant growth and colonization of an industrial soil highly contaminated with Zn (1620 mg kg⁻¹), Cd (60 mg kg⁻¹), Cu (319 mg kg⁻¹), and Pb (756 mg kg⁻¹), suggesting that these amendments could also be used to enhance phytostabilization [62]. Similarly, the addition of manure was shown to be effective for PTE stabilization by enhancing the phytostabilization capacity of *Brassica juncea* [63]. In summary, soil amendments can be combined with phytostabilization to overcome physico-chemical limitations and to establish green cover.

It is important to note that many microorganisms in the rhizosphere play an important role in soil structure maintenance and plant nutrient recycling as a barrier against pathogens, as well as in the detoxification of some PTEs [64]. These microorganisms can modify soil metal speciation and PTE bioavailability and enhance phytostabilization in some cases [53]. However, the ecological effect of foreign bacteria on indigenous microbial communities and their possible impact on human health need to be taken into account. It has been reported that the addition of *Pseudomonas fluorescens* and *Microbacterium* sp. to *Brassica napus* stimulates plant growth and Pb accumulation [65] and that the addition of *Sanguibacter* sp. and *Pseudomonas* sp. increases *Nicotiana tabacum* growth, as well as Cd, Fe, and Zn uptake [66]. It is important to point out that changes in the plant rhizosphere can also enhance tolerance to certain PTEs. Thus, Quiñones et al. [67] found that *Lupinus albus* plants inoculated with a *Bradyrhizobium canariense* bacterium became more Hg-tolerant as compared with control plants. In summary, increases or modifications in the rhizosphere enhance phytoextraction, phytostabilization, and adaptation to AMLs and also increase PTE uptake and immobilization by the plant.

4. Selection of Suitable Plants for AML Restoration

Plant selection is critical for the development of an effective phytoremediation strategy. It is vital to carry out a preliminary study of the AMLs to be restored and to set clear objectives [17]. The major plant selection criteria for phytoremediation include a good level of tolerance to the PTEs present in the AMLs, a sufficiently high potential level of PTE accumulation, translocation, and uptake, a high growth rate, and biomass yield and extreme weather tolerance. Soil physico-chemical properties, the plant's terrestrial, aquatic, or semi-aquatic habitat, as well as the characteristics and depth of roots, also need to be considered [18]. However, exotic plants, even those with good phytoremediation, rapid growth, and excellent colonization capacity, which could lead to unwanted invasions of whole ecosystems, should not be used [68]. In general, the marketing of plants, which have highly specific PTE accumulation and tolerance capacities, as well as specific climate and soil requirements, may not be economically viable [16]. Plants also have certain maintenance requirements such as the addition of fertilizers, periodic replanting and harvesting, as well as the implementation of feasibility studies [29]. In summary, for the purposes of AML remediation, it is necessary to select plants with sufficient growth capacity in the degraded areas and with specific PTE detoxification characteristics [19,69,70]. Researchers have also tried to identify natural metallophyte species that could be useful for AML reclamation. In recent years, molecular genetic techniques, which can be used to improve plant phytoremediation capacity, have been playing an increasingly important role.

4.1. Wild Plant Species

Great strides have been made in the search for and characterization of PTE hyper-accumulating and hypertolerant plants to PTEs. All these data have been collected in databases such as PHYTOREM in Canada and the metal-accumulating plants database METALS funded by the Environmental Consultancy of the University of Sheffield (ECUS) in the UK [71]. However, their use by phytoremediation researchers has been limited. Unfortunately, PTE accumulation and tolerance are influenced by many variables such

as AML physico-chemical characteristics and climatic conditions, in addition to plant physiology. Thus, given the extreme difficulty of using non-native plants for remediation of specific AML areas, the first step in any phytoremediation strategy remains the selection and characterization of AML native plants.

Hyperaccumulator and hyper-tolerant plants often appear to be restricted to metalliferous soils, where their tolerance and accumulation mechanisms operate against PTEs [69,72]. The collection and spread of hyperaccumulators may therefore present challenges due to the difficulty of harvesting, separating, and cleaning natural seeds, as well as to the germination and cultivation requirements of the places where hyperaccumulators naturally grow [73]. The characteristics of hyperaccumulator plant populations often depend on whether they come from metalliferous or nonmetalliferous soils [74]. In research carried out by Posic et al. [72], thallium tolerance and accumulation were compared in one non-metallicolous and three metallicolous populations of TI hyperaccumulator *Biscutella laevigata*. Their results show that TI hyperaccumulation and hyper-tolerance are population-specific traits of *B. laevigata*. In research by Ke et al. [75], a *Rumex japonicus* plant population from a copper mine and another from an uncontaminated site were exposed to Cu. Their results showed that metallicolous plant populations had higher tolerance to Cu and lower nutrient availability than populations from the uncontaminated site. This indicated that the tolerance and accumulation of certain PTEs are population-specific.

Once the plants to be used have been selected, a preliminary study of their PTE accumulation capacity and tolerance mechanisms is recommended (Figure 1) [17,18] in order to assess the suitability of these plants for AML phytoremediation.

4.1.1. Metal Uptake and Accumulation Mechanisms

Several mechanisms enable accumulator plants to absorb PTEs into the roots which are then translocated via various systems to the plant's aerial parts for eventual storage [76]. PTE accumulation goes through four main stages: (1) PTE phytoavailability and uptake from the soil through the roots; (2) accumulation in the roots; (3) translocation from roots to shoots through xylem bundles; and (4) accumulation in the aerial parts of the plant through various mechanisms [33]. PTE phytoavailability depends on soil characteristics such as pH [63,77,78], salinity [79], humidity [80], soil organic matter content [63,77,79], speciation [81], inter-metal competition [78], and cation exchange capacity [77]. These characteristics enable plants to accumulate PTEs in different forms such as free ions or complexes containing organic and inorganic matter [82]. The synergism and antagonism among PTEs and other elements can also have a major impact on their absorption, translocation, and accumulation. In research carried out by Dong [83], competition between Cd and Mn absorption and translocation was observed in *Lycopersicon esculentum* exposed to Cd. Similarly, the addition of Ca reduced Cd absorption by soybean and wheat roots, while the addition of K only diminished Cd absorption by wheat roots [84]. On the other hand, plant roots produce chelating agents and compounds which promote changes in soil pH, as well as redox reactions. Thus, plants are able to solubilize and absorb PTEs present in the substrate [85]. *Triticum aestivum* and *Sorghum bicolor*, which produce phytosiderophores under Fe deficiency stress, facilitate the uptake of other PTEs [86]. Similarly, the use of root exudates from *Belamcanda chinensis* in *Echinochloa crus-galli* exposed to Pb, Cd, and Cu increases plant metal uptake levels [87]. An opposite pattern is observed with respect to *Medicago polymorpha*, whose root organic acid production reduces PTE phytoavailability [88]. The ability of this plant to produce certain compounds that facilitate PTE uptake may be of particular interest with regard to enhancing phytoremediation efficiency.

Once phytoavailable, PTEs can enter roots via apoplastic and symplastic pathways. In the apoplastic pathway, PTEs enter the cells through intracellular spaces, while in the symplastic pathway they are absorbed by the roots through protein transporters located in the membrane [30]. These transporters specialize in the translocation of nutrients within the root, although some PTEs, whose chemical characteristics are very similar to those of certain nutrients, can use the same entry channels [17]. Since PTE uptake depends on the

number of membrane proteins present, PTE absorption is saturable according to Michaelis-Menten kinetics [89]. Each species contains different types and numbers of transporters in each cell, thus making the absorption of a certain element characteristic of the species [76]. Numerous studies of these transporters have been carried out, with the most important Cd transporter families being ZIP, YSL, and NRAMP [90–92], while those for Cu uptake are ZIP, COPT, Cu-ATPases, ATX, CCS, and YSL [93–96]. Thus, PTE absorption by plants, which is influenced by many factors, is a highly complex process. After entering the root, PTEs are translocated to the aerial parts of the plant through xylem bundles, which are involved in the transport of water and dissolved salts [97]. This energy-consuming process only permits the entry of metals through the symplastic pathway, as the Casparian strip in the endodermis closes off access during transport to the xylem [30]. Many natural chelates are synthesized in order to transport the metal to the xylem, which is then transported to the aerial part of the plant. Glutathione, phytochelatins (PCs), organic acids, histidine, and nicotinamide have been reported to facilitate this process [98–102].

After their translocation from the roots to the aerial part of the plant, a large number of chelator compounds play an important role in the accumulation of PTEs, which bind to thiol-rich peptides and phytochelatins (PCs) to facilitate their accumulation in specific locations in the roots and aerial parts of plants [19,69,103–105]. PCs are enzymatically synthesized from glutathione or its homologs through a reaction catalyzed by the PC synthase (PCS), a PTE-activated enzyme [105,106]. These peptides are made up of three amino acids with a typical (γ -GluCys)n-Gly (n = 2–11) sequence. In some plants, the C-terminal glycine is replaced by serine, glutamine, glutamate, and alanine with (γ -Glu-Cys)n- β -Ala, (γ -Glu-Cys)n-Ser and (γ -Glu-Cys)n-Glu sequences [107–111], whose mechanisms of action have been elucidated in *Nicotiana tabacum* in response to Cd. After crossing the plasma membrane, Cd forms complexes with PCs in the cytosol, which are then actively transported into the vacuole [112]. This mechanism facilitates the sequestering of Cd, which is kept away from cellular machinery to prevent any toxic effects on the plant [15]. PCs are induced in response to many PTE ions such as Ag^+ , H_2AsO_3^- , H_2AsO_4^- , Cd^{2+} , Cu^{2+} , Ca^{3+} , Hg^{2+} , Ln^{3+} , Ni^{2+} , Pb^{2+} , SeO_4^{2-} , and Zn^{2+} [107].

Hyperaccumulating plants are usually characterized by limited compartmentalization of PTEs in root vacuoles, which are mainly translocated to the aerial parts of the plant. This was demonstrated by Yang et al. [113], who compared two populations of *Sedum alfredii*. Their results showed that the non-accumulating plant population retained 2.7 times more Zn in root vacuoles than the hyperaccumulating population. The bioaccumulation factor (BF) and translocation factor (TF) are used to evaluate the phytoextraction potential of a given species. The BF indicates a plant's PTE accumulation efficiency in leaves as compared to PTE soil concentrations, while the TF indicates a plant's efficiency in transporting PTEs from roots to leaves [114]. Thus, the BF, which was found to be less than one in metal excluder species, was greater than one in metal hyperaccumulator species [64]. The TF, which indicates where most of the metal is accumulated in roots and shoots, could be an accurate indicator of whether a species is a good phytoextraction or phytostabilization candidate. A TF of less than one indicates that the metal is mainly accumulated in roots, while a TF of greater than one indicates that most of the metal is accumulated in leaves [19,69,70].

4.1.2. Tolerance Mechanisms

Plant growth and development are directly dependent on optimal environmental and nutritional factors, any changes in which produce stress [115]. PTE-related stress can lead to multiple toxic effects of differing intensities depending on plant species, concentration, and phytoavailability, as well as the duration of exposure [16]. PTE toxicity triggers morphological and physiological imbalances in plants [19,48,69,70,116,117]. This is mainly due to the inhibition of most cellular processes at different metabolic levels caused by the appearance of reactive oxygen species (ROS), which produces a state of oxidative stress in the plant [19,118]. From a physiological perspective, it is important to highlight the

decrease in photosynthetic pigment content and in leaf water potential and stomatal closure due to oxidative stress and reduction in photosynthesis [119]. At the biochemical level, the similarity of some PTEs to certain essential elements can result in the replacement of elements in active enzyme sites with PTEs. This results in a reduction or inhibition in enzyme activity which undermines the correct functioning of the entire system [120]. Many PTEs also affect the uptake of essential nutrients which compete for the same transporters, thus causing deficiencies in nutritional status [121].

PTE-related stress is directly correlated with an increase in the accumulation of ROS, which are a group of free radicals produced by sequential reductions in molecular oxygen (O_2). The most common ROS include singlet oxygen (1O_2), hydrogen peroxide (H_2O_2), as well as superoxide (O_2^-) and hydroxyl ($\cdot OH$) radicals [122], which play important roles in the plant's normal metabolism by maintaining a balance between their production and elimination by antioxidant systems. However, exposure to PTEs can cause ROS to exceed the threshold levels required for plant metabolism and trigger damage to proteins, lipids, and nucleic acids, which are the main cellular macromolecules [123]. Tolerance to PTEs in plants is dependent on their ability to activate certain tolerance systems that enable ROS levels to be maintained within normal limits [124]. Plants have complex enzymatic and non-enzymatic antioxidant defenses, which are their main protection against PTEs and enable them to maintain ROS levels compatible with those of regularly functioning cells [124]. The principal enzymatic antioxidative defenses are catalase (CAT), superoxide dismutase (SOD), peroxidase (POD), NADP-dependent dehydrogenases, glutathione S-transferase (GST), and the Foyer-Halliwell-Asada cycle enzymes ascorbate peroxidase (APX), monodehydroascorbate reductase (MDHAR), dehydroascorbate reductase (DHAR), and glutathione reductase (GR) [122]. SOD, which removes O_2^- , is one of the primary antioxidative defenses against PTE toxicity [125]. An increase in SOD activity against a large number of PTEs, such as Pb [69], Cd [126], Cu [127], Hg [128], As [129], Sb [130], Cr [131], and Zn [132], has been reported. To deal with an increase in H_2O_2 , plants activate different enzymes, particularly CAT, POD, and the Foyer-Halliwell-Asada cycle enzymes. Many studies have shown that, by using these enzymes to increase the rate of H_2O_2 removal, certain plants can better tolerate PTEs such as Cd [19], Cu [70], As [129], Sb [133], Cr [131], and Hg [134]. The correct functioning of the Foyer-Halliwell-Asada cycle is promoted by an increase in the NADP-dehydrogenase activities NADP-G6PDH, NADP-IDH, and NADP-MS, which supply the reduction power required by the AsA-GSH cycle [135]. An induction of certain NADP dehydrogenases against Cd [19], Cu [70], and As has been reported [136]. GST enzymes are a diverse group of enzymes which catalyse the conjugation of electrophilic xenobiotic substrates with GSH, thus removing cytotoxic and genotoxic compounds that could damage the cells [137]. This group of enzymes plays an important role in defenses against As [129,137], Cd [19], Cu [70], Zn [138], and Cr [139]. Non-enzymatic components of the antioxidative defense system include ascorbate (AsA), glutathione (GSH), tocopherol, carotenoids, and phenolic compounds [118]. GSH, which is the most abundant low-molecular weight thiol-bearing molecule in cells, constitutes an essential component of PTE resistance due to its role as a substrate in enzymatic antioxidant systems and as a precursor of the PTE chelating molecules phytochelatin and metallothioneins [140]. More recent studies have reported that nitric oxide, (NO), which can prevent oxidative damage by enhancing antioxidant defenses, could be a key factor in tolerance to PTEs [124,141].

4.2. Genetically Modified Plants

Although plants use their natural metabolic mechanisms to take up PTEs from the soil, these mechanisms are often insufficient for the purposes of phytoremediation, which is sometimes ruled out due to the difficulty of finding plants with all the characteristics required [30]. The use of genetic engineering to modify plant characteristics has emerged as a promising solution to overcome the common drawbacks of conventional plants used in phytoremediation strategies [142]. Scientists have managed to develop genetically mod-

ified (GM) plants to overcome the limitations of natural plants, including low biomass production, slow growth, and the lack of tolerance and accumulation mechanisms [143]. However, certain safety and regulatory requirements can limit the commercial use of GM plants [29]. The main problem associated with these types of plants, which show greater survival potential than wild plants, is their possible invasiveness [144]. Consequently, phytoremediation techniques involving GM plants are still at the initial stage of seeking genes that improve the efficiency and development of new transgenic plants which are then tested under laboratory conditions. Several studies, including one carried out by Sanz-Fernández et al. [20], have assembled a “molecular toolbox” of genes for phytoremediation purposes which could be used by other scientists to obtain GM plants. Research carried out on these types of plants shows that genetic engineering could become an important remediation tool. After exposure to PTEs, *Beta vulgaris* L. plants, which were transformed by gamma-glutamylcysteine synthetase (GGCS)-glutathione synthetase (GS) from *Streptococcus thermophilus*, accumulated more Cd, Zn and Cu than wild-type plants and showed higher GSH and PC levels [145]. PpMT2, a metallothionein-encoding gene from *Physcomitrella patens*, was used to improve the tolerance of *Arabidopsis thaliana* plants to high concentrations of Cu and Cd [146]. Similarly, *Arabidopsis thaliana* transgenic plants, expressing the natural resistance-associated macrophage protein 6 (Nramp6) gene from *Sedum alfredii*, exhibited higher Cd accumulation levels as compared with wild-type plants [91]. Although GM plants have yet to be tested on natural AMLs, the ability to genetically engineer plants with ideal characteristics, unlike other remediation techniques, could be of great significance in the future.

5. Real Examples of Using Plants for AML Restoration

Although often misunderstood as simply the removal of human-created structures and waste, AML restoration involves the regeneration of all previously existing biodiversity in an area prior to the commencement of mining operations [3]. While the removal of PTEs accumulated over decades in the environment is insufficient, it is essential to study the whole range of organisms that previously inhabited the location to be returned to its initial state [147]. AML restoration projects initially involve field experiments to regenerate the essential soil fertility conditions through the removal and immobilization of PTEs and through soil restoration. Once this initial stage is completed, the next step will be the restoration of the entire ecosystem.

Given the extent of the progress made, one of the most interesting examples of using plants for AML restoration is the area affected by a mining accident in Aznalcóllar in southwestern Spain in 1998. It was caused by a dam collapse due to an excessive volume of mining waste deposits. The spill affected approximately 4286 ha of alluvial soil, which were flooded by slurry containing high PTE concentrations, including 0.27 mg L^{-1} of As, 0.85 mg L^{-1} of Cd, 3.6 mg L^{-1} of Pb, and 463 mg L^{-1} of Zn, a few hours after the accident [148]. Following the accident and clean-up operations, a large number of studies were carried out on the defensive behavior of herbaceous plants against PTEs accumulated in nearby areas for at least 20 years [149–152]. Later studies focused on testing the potential of using different plant species for phytoremediation purposes. For example, *Helianthus annuus* plants, which were grown in the spill-affected soil, were evaluated for their PTE removal capacity. Unfortunately, the results showed that plant growth in contaminated soils was negatively affected and that amount of PTEs absorbed was very low. Though ruled out for use in phytoextraction, some studies favored this plant species for use in soil conservation [153]. Similarly, two *Brassica juncea* crops containing different organic soil amendments were analyzed on the site affected by the toxic spillage. However, as in the previous case, total PTE uptake was relatively low, rendering its use in the phytoextraction of pluri-contaminated sites ill-advised [154]. Likewise, two cereal crops, barley and triticale, and two *Brassica* sp. crops were grown, whose absorption rates were excessively slow, which cast doubt on the viability of using these crops for decontamination [155]. On the other hand, despite their slow PTE uptake rates, *Lupinus albus* plants showed an increase

in acid soil pH, a decrease in soluble As and Cd fractions, as well as high As and Cd accumulation levels in roots. According to the results obtained, it would be possible to extract approximately $0.54 \text{ g Cd ha}^{-1} \text{ year}^{-1}$ and $5.26 \text{ g As ha}^{-1} \text{ year}^{-1}$ from the contaminated soil by harvesting and removing the lupin shoots. Given these positive results, this plant species was recommended for use in the phytostabilization and revegetation of the affected area [156]. Large samplings were carried out to determine which plants could be used in phytoremediation programs. It is important to highlight the research by Del Rio et al. [149], who carried out periodic field surveys to identify spontaneously growing metal-tolerant species in the area affected by the Aznalcóllar accident. Their results show that, of the 99 plants studied, *Anchusa azurea*, *Beta vulgaris*, *Chamaemelum fuscum*, *Convolvulus arvensis*, *Cynodon dactylon*, *Diplotaxis virgata*, *Erodium aethiopicum*, *Lavatera cretica*, *Malva nicaeensis*, *Silybum marianum* and, particularly, *Amaranthus blitoides* could be used in the remediation of the contaminated area. A phytoremediation experiment was therefore performed to assess the potential use of native shrub species for soil restoration in Aznalcóllar. The results showed that *Retama sphaerocarpa* was the most effective plant for use in phytostabilization programs [157]. Later on, an afforestation soil remediation program was carried out involving the planting of Mediterranean riparian woodland species, such as *Populus alba*, *Tamarix africana*, *Fraxinus angustifolia*, and *Salix atrocinerea*, and the usual drier upland trees such as *Quercus ilex* subsp. *ballota*, *Olea europaea* var. *sylvestris*, *Phillyrea angustifolia*, *Pistacia lentiscus*, *Rosmarinus officinalis*, and *Retama sphaerocarpa* [158]. Domínguez et al. [159] monitored survival and growth patterns in these plantations over a 3-year period. Plant survival rates reached nearly 100% in riparian sites, while some species in the upland terrace sites showed the lowest survival rates. They did not find any relationship between plant efficiency and soil pollution, with PTE accumulation rates found to be very low, except for the Salicaceae species, which showed high Cd and Zn accumulation rates (1 and 200 mg kg^{-1} , respectively). Another study carried out in the area found that nurse shrubs, such as *Retama sphaerocarpa*, were highly effective in reducing *Quercus ilex* mortality, and thus are a promising tool for the revegetation of polluted areas [160]. *Retama sphaerocarpa*, which recorded the highest survival rates and strong tolerance levels, was found to be the most effective plant species for the restoration of these soils [161]. In addition to the revegetation mentioned above, plant growth phytostabilization had a considerable impact on PTE bioavailability [148,157]. In general, the amount of PTEs accumulated in the aerial parts of trees and shrubs was usually observed to be at the very low normal levels for higher plants [158,159], while different levels of accumulated PTEs were recorded in roots [157,162]. For example, *Quercus ilex* was capable of accumulating up to 7 g kg^{-1} of Cd in its fine roots [162] and *Myrtus communis*, *Arbutus unedo*, and *Retama sphaerocarpa* accumulated As in their roots [163]. The area affected by the Aznalcóllar mining accident has become an exceptional location for studying the phytoremediation potential of numerous species in a realistic context and for providing input for future phytoremediation projects.

Given Spain's long history of a highly dynamic mining industry, large areas of the country have been affected by abandoned mining waste [164]. Some studies have collected the results of different plant-based rehabilitation projects in these locations [10,49,165,166]. For example, a reclamation project involving the plantation of *Pinus* sp. or *Eucalyptus* sp. trees and waste material amendments (sewage sludge and paper mill residues) in a Cu mining area located in Touro (Galicia, Spain) [165]. An evaluation of the different reclamation practices showed that soil planted with pines and eucalyptus trees increased Ni, Pb, and Zn retention in non-mobile soil fractions, while the amendments also attenuated Cr- and Cu-related pollution. The results obtained showed that vegetation provided by these tree species, together with the amendments, decreased PTE pollution in the area affected [165]. A field experiment involving the agricultural crop plants *Hordeum vulgare*, *Triticum aestivum*, and *Lupinus luteus* was also performed in an area called Almadén (Ciudad Real, Spain) containing the most unusual large soil concentrations of Hg in the world [167]. The wheat crop was found to extract concentrations of Hg of up to $0.479 \text{ } \mu\text{g g}^{-1}$. However,

the most interesting plant subjected to phytoextraction was barley, which accumulated up to 719 mg ha⁻¹ of Hg [168]. Later studies carried out in this area showed that most Hg uptake by plants was through leaves, with root uptake rates being rather slow [169]. The Cartagena-La Union mining district, located on the Mediterranean coast in the southeast of Spain, is also affected by high levels of PTE accumulation (\approx 1100, 4700 and 5000 mg kg⁻¹ of As, Pb, and Zn, respectively) [49,166]. A remediation experiment combining red mud, compost, and hydrated lime soil amendments with the halophyte plant species *Atriplex helium* and *Zygophyllum fabago* was performed. On the whole, soil amendments increased tailing pH and reduced PTE solubility and extractability, while the halophytes tested, which managed to grow only in amended tailings, accumulated low levels of Al, As, Cd, and Zn. This research showed that the combined use amendments and halophyte plant species is an effective AML phytostabilization strategy [166]. A large-scale field experiment, also conducted in this area, involved the reclamation of a tailing's disposal pond through the spontaneous colonization of native species mediated by tilling, marble waste, and pig slurry. Five years later, *Atriplex halimus*, showing high metal concentrations, mainly in leaves, had become the predominant species. Accumulation levels were increased differentially using different amendments, with pig slurry increasing Cu concentrations and marble waste increasing Pb and Cd accumulation levels. In summary, *Atriplex halimus* could be a good candidate for AML phytoremediation [49]. Another study carried out in the Sarzedas mining area (Castelo Branco, Portugal), which was greatly affected by As (11.1–651.1 mg kg⁻¹), Sb (30.5–5986.4 mg kg⁻¹), and W (0.8–684.0 mg kg⁻¹) contamination, analyzed spontaneous vegetation growth. The results showed that the following plants could be used to reduce PTE bioavailability: *Pinus pinaster* for As and W, *Cistus ladanifer* and *Cistus vulgaris* for Sb and W, *Digitalis purpurea* and *Erica umbellata* for Sb, and *Quercus ilex*, *Chamaespartium tridentatum*, and *Eucalyptus globulus* for W [170]. Elsewhere in Europe, with its large number of AMLs, a study of a tin mining area in Abbaretz (Loire-Atlantique, Pays de la Loire, France) highlighted the importance of improving soil conditions to establish vegetation correctly. In this research, the effect of biochar amendments combined with iron sulphate supplements on the growth and uptake of three Salicaceae species was evaluated. The results showed that these amendments improved soil health and raised PTE concentrations in the roots of *Salix purpurea*, which could be a good candidate for PTE phytostabilization [171]. Similarly, a field experiment was conducted in the Campo Pisano mining area (Sardinia, Italy), with 3250 mg kg⁻¹ of Pb and 11,949 mg kg⁻¹ of Zn, to study the feasibility of using *Pistacia lentiscus* and *Scrophularia bicolor* combined with soil amendments for phytoremediation purposes. The results showed that the plant species *Pistacia lentiscus* could be used effectively for PTE phytostabilization due to its capacity to accumulate PTEs in roots, as well as its high resistance and biomass production [172]. Other studies have investigated the establishment of plant cover to improve soil quality. For example, in two AMLs in northwestern Transylvania (Romania), a study was carried out to monitor the effect of different plant covers on soil health after 40 years of vegetation growth. The results revealed that the chemical, physical, and biological properties of the soil, which improved significantly during the period of the study, depended on the plant cover [35].

In the United States, over 550,000 abandoned mining sites have generated 45 billion tons of mining waste, including waste rock and tailing material, containing worrying levels of PTE concentrations [3]. While some researchers have been evaluating emerging phytoremediation technologies, most researchers have focused on the use of revegetation techniques to improve soil health. A revegetation study of three mining sites in Ohio, using different mixes of standard reclamation and native plants, compared the growth of each type of revegetation and its impact on soil microorganisms over a 2-year period. The results obtained highlighted the importance of using native plants which can improve soil health more effectively than non-native plants [173]. An Iron King Mine and Humboldt Smelter Superfund (IKMHSS) field study carried out by Gil-Loaiza, which found over 2000 mg kg⁻¹ of Pb, As, and Zn, also demonstrated the importance of incorporating good

native plant cover in AMLs [174]. Six different treatments containing different amounts of compost and a mix of native plants, which were pre-tested in the laboratory, were studied. It is interesting to note that plant growth in tailing soils was found to stabilize pH levels through rhizosphere alkalisation and to prevent smelter tailing acidification. In summary, both studies showed that the establishment of vegetation cover improved soil quality through increased microbial activity and PTE phytostabilization [173,174].

In China, about 2.88 Mha of land, which is increasing at an alarming rate of 46,700 ha per year, has been degraded by mining operations [175]. The many studies of phytoremediation techniques used in China, outlining the most important advances made in this field, are reviewed below. The remediation of the Dabaoshan polymetallic mining area in the Guangdong province, with 681, 343 and 339 mg kg⁻¹ of Pb, Cu and Zn, respectively, combining lime, organic fertilizer, and biochar soil amendments with *Hibiscus cannabinus* and *Boehmeria nivea* plant species, was studied. The results showed that amendments and plant growth improved soil health by reducing PTE bioavailability, solubilization, and migration in mine tailings [176]. The phytoextraction capacity of heavy metal-accumulating plants and chemical chelators has also been evaluated in Guangdong paddy fields that had been continuously irrigated with mining wastewater for 55 years (868.5, 433.6, 58.6, and 2.2 mg kg⁻¹ of Pb, Zn, Cu, and Cd, respectively). The results suggested that *Amaranthus hypochondriacus* was highly effective for in situ PTE phytoremediation and that the application of ethylenediaminetetraacetic acid (EDTA) accelerated the phytoextraction effect [177]. A field experiment involving plants, microorganisms, and amendments was also carried out on the Anshan Mining Group Corporation's metal-contaminated mine tailings in the Liaoning province (51.7, 151.5, 43.6, and 1960.0 mg kg⁻¹ of Cu, Zn, Pb, and Mn, respectively). The experiment, which combined *Glycine max* and *Mucor circinelloides* fungi, as well as the application of rice husk, biochar, and ceramsite soil amendments in 17 phytoremediation tree plots, was carried out over a period of 105 days. The combination of these techniques was found to improve PTE phytoextraction capacity, reduce PTE soil concentrations, and to increase microbial soil activity [178]. Similar results were reported by Yu et al. [179] using *Celosia argentea* and ammonium chloride, *Bacillus megaterium*, and citric acid soil amendments for 10 months in mine tailing-contaminated soil in southern China (3.68 mg kg⁻¹ of Cd). Soil Cd concentrations were decreased after three harvesting events, with overall reductions in Cd of 40.5%, 46.1%, and 105% following applications of ammonium chloride, *B. megaterium*, and citric acid, respectively. The study showed that the use of *C. argentea* combined with soil amendments could greatly improve soil enzyme activity and the quality of Cd-contaminated soils. Another experiment in Chenzhou (Hunan province), one of the largest ferrous metal mines in China, assessed the restoration and accumulation potential of selected hybrid sweet sorghum species in contaminated areas (288–1270, 97.7–1250 and 2–5.7 mg kg⁻¹ of Zn, Pb, and Cd, respectively). The results showed that hybrid sweet sorghum species had greater accumulation and tolerance capacity than local species. The study concluded that it would be economically viable to use sweet sorghum, which densifies biofuel production, for PTE remediation in this mining area [180]. Similarly, an evaluation of the *Manihot esculenta* bioethanol plantation in a severely PTE-contaminated mining area found that *Manihot esculenta* could produce 3680–4160 L of bioethanol per hectare (L ha⁻¹) on contaminated soils. *Manihot esculenta* also improved soil fertility by reducing mobile and bioavailable metal fractions. The cultivation of *Manihot esculenta* could therefore be used for AML remediation and also generate economic benefits [181]. Other AML phytoremediation projects have evaluated the relationship between revegetation and improved chemical, physical, and microbiological soil properties. Research conducted in the AnTaibao, Anjialing, and East opencast coal mining areas analyzed the effects of revegetation on physical, chemical, and microbiological soil properties in nine plots for 1–13 years. The results obtained showed that the plant species significantly enhanced soil health by improving soil field capacity, organic content, aggregation, microbiological composition, and total microorganisms excluding other factors such as soil texture in these areas [175]. Revegetation projects were carried out in the Lechand

Pb/Zn mine in Guangdong province, which is severely contaminated by Pb (29.9 mg kg^{-1}), Zn (50.7 mg kg^{-1}) Cu (14.7 mg kg^{-1}), and Cd (0.52 mg kg^{-1}). A combination of domestic waste and complex fertilizer amendments and plants was evaluated to assess their suitability for revegetation purposes. The results showed that *Vertiveria zizanioides* was the best plant species for tailings revegetation, followed by *Paspalum notatum*, *Cynodon dactylon*, and *Imperata cylindrica* var. major [182]. Similarly, research carried out in the Jiangxi province, which is affected by ionic rare earth element mine tailings, showed that a combination of organic amendments and plants could regenerate bacterial community structures and diversity, improve soil quality, and possibly reduce PTE bioavailability [183]. Similar results were reported by Mi et al. [184] in the Kee Open Pit Mine in Shanxi province, where the use of mine dumps for crop cultivation improved soil health as measured by soil properties and microbial communities. On the other hand, different periods of revegetation were evaluated in metal mining dump areas (up to 306.5, 1863.9, 494.5, 353, and 133.3 mg kg^{-1} of Cu, Mn, Cr, Ni, and Zn, respectively) in the Sichuan province to determine the long-term impact of revegetation. The results showed that plant species with low vegetation diversity had a strong impact on PTE remediation, but that the effect of soil fertility restoration was less than that of natural vegetation, while spontaneous vegetation succession reduced the rate of PTE remediation. The study suggested that revegetation techniques need to be monitored to ensure that succession does not disrupt soil remediation [185]. Finally, it is important to highlight the study carried out by Wan et al. [14], in which the costs and benefits of a phytoremediation project involving soil contaminated with As, Cd, and Pb are calculated. They concluded that phytoremediation costs for this project were lower than those reported for other remediation projects. Moreover, given the cost of environmental pollution, the benefits of phytoremediation are expected to offset project costs in less than seven years.

The Asia-Pacific region, one the largest producers of coal, accounts for over 67% of total world production. Land degradation due to coal mining operations is reported to affect 1400 additional hectares each year [186]. Several studies have described phytoremediation projects, in which different soil restoration strategies involving herbaceous plants and trees in coal mining areas have been used [186–189]. The use of *Miscanthus x giganteus* as an effective plant to restore soil in a lignite mine has been analyzed by Jeżowski et al. [188]. This 3-year field study of lignite mining waste heaps showed positive results in terms of soil regeneration. A phytoremediation study of the removal of toxic iron and sulphur aqueous waste spilled on soils around coal mines was also carried out in the Jaintia hills (Meghalaya, northeastern India). The use of agricultural lime and organic manure soil amendments, together with plants, reduced sulphate concentrations from 22 to 5 mg kg^{-1} and soluble iron from 476 to 109 mg kg^{-1} . Moreover, this reclamation strategy improved soil fertility by increasing plant nutrients, pH (up to normal levels) and microbial biomass [186]. The phytoextraction capacity of *Eucalyptus* sp. was evaluated in the Jharia coal mine (Dhanbad, India), the results of which were compared with PTE content measured in soil and trees in 1995. The study concluded that plant-based reclamation reduced Pb, Zn, and Cu concentrations to their levels in 1995, while PTEs mainly accumulated in the stem bark (Pb), leaves (Zn and Mn), and roots (Cu and Co) of trees [189].

The results of AML restoration projects described above showed that the establishment of vegetation cover in these areas effectively reduced PTE bioavailability and increased soil quality, which constituted a further step towards a total restoration of the ecosystem.

Table 1. List of hyperaccumulator plants that could be used for phytoextraction. Asterisks indicate approximate values (*).

Species	Maximum Data of PTEs Concentration in Leaves (mg kg^{-1} DW)		References
<i>Alyssum</i> sp.	Ni	20,100	[190]

Table 1. Cont.

Species	Maximum Data of PTEs Concentration in Leaves (mg kg ⁻¹ DW)		References
<i>Arabidopsis halleri</i>	Cd	5722	[191]
	Zn	9491	
<i>Arabis gemmifera</i>	Cd	1810	[192]
	Zn	20,300	
<i>Atriplex halimus</i>	Cd	850 *	[193]
<i>Berkheya coddii</i>	Ni	31,300	[194]
<i>Bidens pilosa</i>	Cd	144.1	[195]
<i>Biscutella laevigata</i>	Tl	66.7	[196]
<i>Brassica pekinensis</i>	Pb	2670	[197]
<i>Carpobrotus rossi</i>	Cd	172	[198]
<i>Lantana câmara</i>	Cd	423.06	[46]
<i>Leersia hexandra</i>	Cr	5005	[199]
<i>Lonicera japonica</i>	Cd	57.22	[200]
<i>Phytolacca acinosa</i>	Mn	19,300	[201]
<i>Polygonum lapathifolium</i>	Mn	18,841.7	[202]
<i>Pteris vittata</i>	As	1530	[203]
<i>Schima superba</i>	Mn	9975.61	[204,205]
<i>Sedum alfredii</i>	Cd	9000	[206]
	Zn	4807	[207]
<i>Sesbania drummondii</i>	Pb	5000 *	[208]
<i>Solanum</i> sp.	Cd	142.7	[209]
<i>Thlaspi</i> sp.	Cd	164	[210]
	Ni	16,600	[190]
	Zn	19,070	[210]

Table 2. List of plants capable of volatilizing PTEs.

Species	PTEs	References
<i>Pteris vittata</i>	As	[211]
<i>Arundo donax</i>		[56]
Transgenic <i>Arabidopsis thaliana</i> (fungal gene WaarsM)		[212]
<i>Dittrichia viscosa</i>		[213]
<i>Oryza sativa</i>		[214]
<i>Zea mays</i>		[57]
Transgenic <i>Nicotiana tabacum</i> (bacterial gene merA and merB)	Hg	[215]
<i>Halimione portulacoides</i>		[216]
<i>Arabidopsis thaliana</i>		[217]

Table 2. Cont.

Species	PTEs	References
<i>Brassica juncea</i>		[218]
<i>Oryza sativa</i> <i>Brassica oleracea</i> <i>Daucus carota</i> <i>Hordeum vulgare</i> <i>Medicago sativa</i> <i>Lycopersicon esculentum</i> <i>Cucumis sativus</i> <i>Gossypium hirsutum</i> <i>Solanum melongena</i> <i>Zea mays</i>	Se	[219]
<i>Populus tremula x alba</i>		[220]
<i>Stanleya pinnata</i>		[221]
<i>Agrostis tenuis</i> <i>Hordeum vulgare</i>		[222]

Table 3. List of suitable plants for phytostabilization purposes.

Species	PTEs	References
<i>Acanthus ilicifolius</i>	Cd	[223]
<i>Agrostis capillaris</i>	Pb	[224]
<i>Arundo donax</i>	Ni, Pb and Hg	[225]
<i>Atriplex halimus</i>	Pb and Cd	[49,94]
<i>Biscutella auriculata</i>	Pb, Cu and Cd	[19,69,70]
<i>Brassica juncea</i>	Cu and Zn	[63]
<i>Brassica juncea</i> <i>Dactylis glomerata</i>	Cd, Pb and Zn	[226]
<i>Casdaminiopsis arenosa</i>	Zn, Cd and Pb	[227]
<i>Conocarpus erectus</i> <i>Populus deltoides</i>	As	[228]
<i>Dahlia pinnata</i>	As	[229]
<i>Imperata condensata</i>	Cu	[230]
<i>Iris lactea</i>	Cd	[126]
<i>Jatropha curcas</i>	Al, Zn and Cu	[231]
<i>Lolium perenne</i>	Cu	[232]
<i>Lupinus albus</i>	Cd, As and Pb	[156,233]
<i>Miscanthus sacchariflorus</i>	Cd	[234]
<i>Miscanthus sinensis x giganteus</i>	Zn, Cd and Pb	[235]
<i>Pistacia lentiscus</i>	Pb	[236]
<i>Pteridium aquilinum</i>	Cd, Cu, Pb and Zn	[236]
<i>Quercus ilex</i>	Cd	[162]
<i>Ricinus communis</i>	Pb	[237]
<i>Salix purpurea</i>	As	[171]
<i>Typha latifolia</i>	Co, Cd, Ni, Mn, Cr and As	[238]

6. Conclusions

Soil restoration in accordance with ecologically sustainable development norms is essential once mining operations are closed. The goal of land reclamation is to return degraded AMLs to their original state and to ensure that the ecosystem's normal interactions between biotic and abiotic components take place. Plant-based techniques, which include phytoextraction, phytostabilization, and phytovolatilization, are environmentally-friendly and cost-effective alternatives to traditional remediation procedures which are usually more destructive to soils. However, the principal drawback of this remediation strategy is the need to select plants that can meet the objectives set out in the strategy. Their PTE tolerance and accumulation systems must enable the plants selected to play an effective role in AML reclamation. Implementation of a phytoremediation strategy will therefore include the following steps: (a) soil physical-chemical characterization; (b) description of climatic conditions; (c) identification of plants capable of growing in the area; (d) assessment of phytoremediation capacities; (e) characterization of PTE tolerance and accumulation mechanisms; and, finally, (f) in situ experiments to assess the success of the strategy. In many cases, little is known about the underlying tolerance and accumulation mechanisms. A better understanding of these mechanisms is crucial to decipher and thus to enhance a plant's capacity to be used for phytoextraction, phytostabilization, and phytovolatilization. This will also facilitate the development of transgenic plants, which may be of great potential significance in the future. In situ phytoremediation studies of AMLs demonstrated that plant-based techniques regenerate soil fertility, reduce PTE bioavailability, and increase biodiversity on-site. New market opportunities in mine land phytoremediation, which uses plants and bio-ores to obtain, in particular, critical precious metals, should be explored. However, research in this field, which is still in its infancy, needs to better identify plants suitable for these purposes in order to enhance their capacity to extract, reduce, and stabilize PTE concentrations in AMLs, as well as to accumulate metals in shoots to facilitate precious metal recovery.

Author Contributions: Conceptualization, J.D.P.; resources and investigation, J.D.P., P.H., J.A.C., J.M.E., L.M.S., M.M.M. and F.B.-B.; writing—original draft preparation, J.D.P. and P.H.; writing—review and editing, P.H., J.A.C., J.M.E., L.M.S., M.M.M. and F.B.-B.; supervision, L.M.S., P.H., and J.A.C.; project administration, P.H. and L.M.S.; funding acquisition, L.M.S., P.H. and M.M.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Spanish Ministry of Science, Innovation and Universities (CGL2015-67644-R and PGC2018-098372-B-100), the Castilla-La Mancha Regional Government (SBPLY/17/180501/000273) and the University of Castilla La-Mancha RDi Internal Research Plan (predoctoral contract 2016/14100).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We wish to thank Michael O'Shea for proofreading the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Elmayel, I.; Esbrí, J.M.; García-Ordiales, E.; Elouaer, Z.; Garcia-Noguero, E.M.; Bouzid, J.; Campos, J.A.; Higuera, P.L. Biogeochemical assessment of the impact of Zn mining activity in the area of the Jebel Trozza mine, Central Tunisia. *Environ. Geochem. Health* **2020**, *42*, 3529–3542. [[CrossRef](#)] [[PubMed](#)]
2. Jain, M.K.; Das, A. Impact of Mine Waste Leachates on Aquatic Environment: A Review. *Curr. Pollut. Rep.* **2017**, *3*, 31–37. [[CrossRef](#)]
3. Venkateswarlu, K.; Nirola, R.; Kuppasamy, S.; Thavamani, P.; Naidu, R.; Megharaj, M. Abandoned metalliferous mines: Ecological impacts and potential approaches for reclamation. *Rev. Environ. Sci. Biotechnol.* **2016**, *15*, 327–354. [[CrossRef](#)]
4. Tchounwou, P.B.; Yedjou, C.G.; Patlolla, A.K.; Sutton, D.J. Heavy metal toxicity and the environment. *Mol. Clin. Environ. Toxicol.* **2012**, *101*, 133–164.

5. Sengupta, M. *Environmental Impacts of Mining: Monitoring, Restoration, and Control*; Routledge: London, UK, 2018; ISBN 9781351450539.
6. United Nations. Transforming Our World: The 2030 Agenda for Sustainable Development. Available online: <https://sustainabledevelopment.un.org/post2015/transformingourworld> (accessed on 27 May 2021).
7. Xie, L.; van Zyl, D. Distinguishing reclamation, revegetation and phytoremediation, and the importance of geochemical processes in the reclamation of sulfidic mine tailings: A review. *Chemosphere* **2020**, *252*, 126446. [CrossRef] [PubMed]
8. Dada, E.; Njoku, K.; Osuntoki, A.; Akinola, M. A review of current techniques of in situ Physico-chemical and biological remediation of heavy metals polluted soil. *Ethiop. J. Environ. Stud. Manag.* **2015**, *8*, 606. [CrossRef]
9. Mahar, A.; Wang, P.; Ali, A.; Awasthi, M.K.; Lahori, A.H.; Wang, Q.; Li, R.; Zhang, Z. Challenges and opportunities in the phytoremediation of heavy metals contaminated soils: A review. *Ecotoxicol. Environ. Saf.* **2016**, *126*, 111–121. [CrossRef]
10. Rodríguez, L.; Gómez, R.; Sánchez, V.; Villaseñor, J.; Alonso-Azcárate, J. Performance of waste-based amendments to reduce metal release from mine tailings: One-year leaching behaviour. *J. Environ. Manag.* **2018**, *209*, 1–8. [CrossRef]
11. Rodríguez, L.; Gómez, R.; Sánchez, V.; Alonso-Azcárate, J. Chemical and plant tests to assess the viability of amendments to reduce metal availability in mine soils and tailings. *Environ. Sci. Pollut. Res.* **2016**, *23*, 6046–6054. [CrossRef]
12. Ruiz, E.; Rodríguez, L.; Alonso-Azcárate, J.; Rincón, J. Phytoextraction of metal polluted soils around a Pb-Zn mine by crop plants. *Int. J. Phytoremediat.* **2009**, *11*, 360–384. [CrossRef]
13. Yan, A.; Wang, Y.; Tan, S.N.; Mohd Yusof, M.L.; Ghosh, S.; Chen, Z. Phytoremediation: A Promising Approach for Revegetation of Heavy Metal-Polluted Land. *Front. Plant Sci.* **2020**, *11*, 1–15. [CrossRef]
14. Wan, X.; Lei, M.; Chen, T. Cost-benefit calculation of phytoremediation technology for heavy-metal-contaminated soil. *Sci. Total Environ.* **2016**, *563–564*, 796–802. [CrossRef]
15. Ali, H.; Khan, E.; Sajad, M.A. Phytoremediation of heavy metals—Concepts and applications. *Chemosphere* **2013**, *91*, 869–881. [CrossRef]
16. Suman, J.; Uhlík, O.; Viktorova, J.; Macek, T. Phytoextraction of heavy metals: A promising tool for clean-up of polluted environment? *Front. Plant Sci.* **2018**, *9*, 1476. [CrossRef]
17. Dickinson, N. Phytoremediation. *Encycl. Appl. Plant Sci.* **2016**, *3*, 327–331.
18. Sarma, H. Metal hyperaccumulation in plants: A review focusing on phytoremediation technology. *J. Environ. Sci. Technol.* **2011**, *4*, 118–138. [CrossRef]
19. Peco, J.D.; Campos, J.A.; Romero-Puertas, M.C.; Olmedilla, A.; Higuera, P.; Sandalio, L.M. Characterization of mechanisms involved in tolerance and accumulation of Cd in *Biscutella auriculata* L. *Ecotoxicol. Environ. Saf.* **2020**, *201*, 110784. [CrossRef]
20. Sanz-Fernández, M.; Rodríguez-Serrano, M.; Sevilla-Perea, A.; Pena, L.; Mingorance, M.D.; Sandalio, L.M.; Romero-Puertas, M.C. Screening *Arabidopsis* mutants in genes useful for phytoremediation. *J. Hazard. Mater.* **2017**, *335*, 143–151. [CrossRef]
21. Macek, T.; Kotrba, P.; Svatos, A.; Novakova, M.; Demnerova, K.; Mackova, M. Novel roles for genetically modified plants in environmental protection. *Trends Biotechnol.* **2008**, *26*, 146–152. [CrossRef]
22. Ferreira, R.A.; Pereira, M.F.; Magalhães, J.P.; Maurício, A.M.; Caçador, I.; Martins-Dias, S. Assessing local acid mine drainage impacts on natural regeneration-revegetation of São Domingos mine (Portugal) using a mineralogical, biochemical and textural approach. *Sci. Total Environ.* **2021**, *755*, 142825. [CrossRef]
23. Moreno González, R.; Cánovas, C.R.; Olías, M.; Macías, F. Seasonal variability of extremely metal rich acid mine drainages from the Tharsis mines (SW Spain). *Environ. Pollut.* **2020**, *259*, 113829. [CrossRef]
24. Pavoni, E.; Covelli, S.; Adami, G.; Baracchini, E.; Cattelan, R.; Crosera, M.; Higuera, P.; Lenaz, D.; Petranich, E. Mobility and fate of Thallium and other potentially harmful elements in drainage waters from a decommissioned Zn-Pb mine (North-Eastern Italian Alps). *J. Geochem. Explor.* **2018**, *188*, 1–10. [CrossRef]
25. Sánchez-Donoso, R.; Martín-Duque, J.F.; Crespo, E.; Higuera, P.L. Tailing’s geomorphology of the San Quintín mining site (Spain): Landform catalogue, aeolian erosion and environmental implications. *Environ. Earth Sci.* **2019**, *78*, 166. [CrossRef]
26. Akcil, A.; Koldas, S. Acid Mine Drainage (AMD): Causes, treatment and case studies. *J. Clean. Prod.* **2006**, *14*, 1139–1145. [CrossRef]
27. United States Department of Agriculture of Soil Health. NRCS Soils. Available online: <https://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/> (accessed on 2 March 2021).
28. Gallego, S.; Esbrí, J.M.; Campos, J.A.; Peco, J.D.; Martín-Laurent, F.; Higuera, P. Microbial diversity and activity assessment in a 100-year-old lead mine. *J. Hazard. Mater.* **2020**, *420*, 124618. [CrossRef]
29. Muthusaravanan, S.; Sivarajasekar, N.; Vivek, J.S.; Paramasivan, T.; Naushad, M.; Prakashmaran, J.; Gayathri, V.; Al-Duaij, O.K. Phytoremediation of heavy metals: Mechanisms, methods and enhancements. *Environ. Chem. Lett.* **2018**, *16*, 1339–1359. [CrossRef]
30. Ghori, Z.; Iftikhar, H.; Bhatti, M.F.; Nasar-um-Minullah; Sharma, I.; Kazi, A.G.; Ahmad, P. *Phytoextraction*; Elsevier Inc.: Amsterdam, The Netherlands, 2015; ISBN 9780128031582.
31. Grobelak, A. Organic soil amendments in the phytoremediation process. In *Phytoremediation: Management of Environmental Contaminants*; Springer International Publishing: Cham, Switzerland, 2016; Volume 4, pp. 21–39; ISBN 9783319418117.
32. Jadia, C.D.; Fulekar, M.H. Phytoremediation of heavy metals: Recent techniques. *Afr. J. Biotechnol.* **2009**, *8*, 921–928.
33. Zhao, X.; Sun, Y.; Huang, J.; Wang, H.; Tang, D. Effects of soil heavy metal pollution on microbial activities and community diversity in different land use types in mining areas. *Environ. Sci. Pollut. Res.* **2020**, *27*, 20215–20226. [CrossRef]

34. Sheoran, V.; Sheoran, A.S.; Poonia, P. Soil Reclamation of Abandoned Mine Land by Revegetation: A Review. *Int. J. Soil Sediments Water* **2010**, *3*, 13.
35. Buta, M.; Blaga, G.; Paulette, L.; Păcurar, I.; Roșca, S.; Borsai, O.; Grecu, F.; Sînziana, P.E.; Negrușier, C. Soil Reclamation of Abandoned Mine Lands by Revegetation in Northwestern Part of Transylvania: A 40-Year Retrospective Study. *Sustainability* **2019**, *11*, 3393. [[CrossRef](#)]
36. Alkorta, I.; Hernández-Allica, J.; Becerril, J.M.; Amezaga, I.; Albizu, I.; Garbisu, C. Recent findings on the phytoremediation of soils contaminated with environmentally toxic heavy metals and metalloids such as zinc, cadmium, lead, and arsenic. *Rev. Environ. Sci. Biotechnol.* **2004**, *3*, 71–90. [[CrossRef](#)]
37. Liu, L.; Li, W.; Song, W.; Guo, M. Remediation techniques for heavy metal-contaminated soils: Principles and applicability. *Sci. Total Environ.* **2018**, *633*, 206–219. [[CrossRef](#)] [[PubMed](#)]
38. Black, H. Absorbing possibilities: Phytoremediation. *Environ. Health Perspect.* **1995**, *103*, 1106–1108. [[CrossRef](#)] [[PubMed](#)]
39. Chalot, M.; Blaudez, D.; Rogaume, Y.; Provent, A.S.; Pascual, C. Fate of trace elements during the combustion of phytoremediation wood. *Environ. Sci. Technol.* **2012**, *46*, 13361–13369. [[CrossRef](#)]
40. Corzo Remigio, A.; Chaney, R.L.; Baker, A.J.M.; Edraki, M.; Erskine, P.D.; Echevarria, G.; van der Ent, A. Phytoextraction of high value elements and contaminants from mining and mineral wastes: Opportunities and limitations. *Plant Soil.* **2020**, *449*, 11–37. [[CrossRef](#)]
41. Novo, L.A.B.; Castro, P.M.L.; Alvarenga, P.; da Silva, E.F. Phytomining of rare and valuable metals. In *Phytoremediation: Management of Environmental Contaminants*; Springer International Publishing: Cham, Switzerland, 2017; Volume 5, pp. 469–486. ISBN 9783319523811.
42. Li, Y.M.; Chaney, R.L.; Brewer, E.P.; Angle, J.S.; Nelkin, J. Phytoextraction of nickel and cobalt by hyperaccumulator *Alyssum* species grown on nickel-contaminated soils. *Environ. Sci. Technol.* **2003**, *37*, 1463–1468. [[CrossRef](#)]
43. Nkrumah, P.N.; Echevarria, G.; Erskine, P.D.; Chaney, R.L.; Sumail, S.; van der Ent, A. Growth effects in tropical nickel-agromining “metal crops” in response to nutrient dosing. *J. Plant Nutr. Soil Sci.* **2019**, *182*, 715–728. [[CrossRef](#)]
44. Barbaroux, R.; Plasari, E.; Mercier, G.; Simonnot, M.O.; Morel, J.L.; Blais, J.F. A new process for nickel ammonium disulfate production from ash of the hyperaccumulating plant *Alyssum murale*. *Sci. Total Environ.* **2012**, *423*, 111–119. [[CrossRef](#)]
45. Gisbertz, K.; Hilgendorf, S.; Friedrich, B.; Heinrichs, S.; Rießmann, D.; Pretz, T. Maximising metal recovery from incineration ashes. In Proceedings of the European Metallurgical Conference, EMC 2013, Weimar, Germany, 23–26 June 2013; pp. 1127–1132.
46. Liu, S.; Ali, S.; Yang, R.; Tao, J.; Ren, B. A newly discovered Cd-hyperaccumulator *Lantana camara* L. *J. Hazard. Mater.* **2019**, *371*, 233–242. [[CrossRef](#)]
47. Grčman, H.; Velikonja-Bolta, Š.; Vodnik, D.; Kos, B.; Leštan, D. EDTA enhanced heavy metal phytoextraction: Metal accumulation, leaching and toxicity. *Plant Soil* **2001**, *235*, 105–114. [[CrossRef](#)]
48. Moslehi, A.; Feizian, M.; Higuera, P.; Eisvand, H.R. Assessment of EDDS and vermicompost for the phytoextraction of Cd and Pb by sunflower (*Helianthus annuus* L.). *Int. J. Phytoremediat.* **2019**, *21*, 191–199. [[CrossRef](#)]
49. Acosta, J.A.; Abbaspour, A.; Martínez, G.R.; Martínez-Martínez, S.; Zornoza, R.; Gabarrón, M.; Faz, A. Phytoremediation of mine tailings with *Atriplex halimus* and organic/inorganic amendments: A five-year field case study. *Chemosphere* **2018**, *204*, 71–78. [[CrossRef](#)]
50. Nörtemann, B. Biodegradation of Chelating Agents: EDTA, DTPA, PDTA, NTA, and EDDS. *Biogeochem. Chelating Agents* **2005**, 150–170. [[CrossRef](#)]
51. Sun, B.; Zhao, F.J.; Lombi, E.; McGrath, S.P. Leaching of heavy metals from contaminated soils using EDTA. *Environ. Pollut.* **2001**, *113*, 111–120. [[CrossRef](#)]
52. Meers, E.; Ruttens, A.; Hopgood, M.J.; Samson, D.; Tack, F.M.G. Comparison of EDTA and EDDS as potential soil amendments for enhanced phytoextraction of heavy metals. *Chemosphere* **2005**, *58*, 1011–1022. [[CrossRef](#)]
53. Meers, E.; Hopgood, M.; Lesage, E.; Vervaeke, P.; Tack, F.M.G.; Verloo, M.G. Enhanced phytoextraction: In search of EDTA alternatives. *Int. J. Phytoremediat.* **2004**, *6*, 95–109. [[CrossRef](#)]
54. Duquène, L.; Vandenhove, H.; Tack, F.; Meers, E.; Baeten, J.; Wannijn, J. Enhanced phytoextraction of uranium and selected heavy metals by Indian mustard and ryegrass using biodegradable soil amendments. *Sci. Total Environ.* **2009**, *407*, 1496–1505. [[CrossRef](#)]
55. Sakakibara, M.; Watanabe, A.; Sano, S.; Inoue, M.; Kaise, T. Phytoextraction and phytovolatilization of arsenic from as-contaminated soils by *Pteris vittata*. In Proceedings of the Association for Environmental Health and Sciences—22nd Annual International Conference on Contaminated Soils, Sediments and Water 2006, Amherst, MA, USA, 16–19 October 2006; Volume 12, pp. 258–263.
56. Guarino, F.; Miranda, A.; Castiglione, S.; Ciatelli, A. Arsenic phytovolatilization and epigenetic modifications in *Arundo donax* L. assisted by a PGPR consortium. *Chemosphere* **2020**, *251*, 126310. [[CrossRef](#)]
57. Mello, I.S.; Targanski, S.; Pietro-Souza, W.; Frutuoso Stachack, F.F.; Terezo, A.J.; Soares, M.A. Endophytic bacteria stimulate mercury phytoremediation by modulating its bioaccumulation and volatilization. *Ecotoxicol. Environ. Saf.* **2020**, *202*, 110818. [[CrossRef](#)]
58. Raskin, I.; Ensley, B.D. *Phytoremediation of Toxic Metals-Using Plants to Clean Up the Environment*; Wiley Inter-Science Publication: New York, NY, USA, 2001.
59. Shackira, A.M.; Puthur, J.T. Phytostabilization of Heavy Metals: Understanding of Principles and Practices. In *Plant-Metal Interactions*; Springer International Publishing: Cham, Switzerland, 2019; pp. 263–282.

60. Marques, A.P.G.C.; Rangel, A.O.S.S.; Castro, P.M.L. Remediation of heavy metal contaminated soils: Phytoremediation as a potentially promising clean-Up technology. *Crit. Rev. Environ. Sci. Technol.* **2009**, *39*, 622–654. [[CrossRef](#)]
61. Lee, S.H.; Park, H.; Koo, N.; Hyun, S.; Hwang, A. Evaluation of the effectiveness of various amendments on trace metals stabilization by chemical and biological methods. *J. Hazard. Mater.* **2011**, *188*, 44–51. [[CrossRef](#)]
62. Ruttens, A.; Mench, M.; Colpaert, J.V.; Boisson, J.; Carleer, R.; Vangronsveld, J. Phytostabilization of a metal contaminated sandy soil. I: Influence of compost and/or inorganic metal immobilizing soil amendments on phytotoxicity and plant availability of metals. *Environ. Pollut.* **2006**, *144*, 524–532. [[CrossRef](#)]
63. Pérez-Esteban, J.; Escolástico, C.; Moliner, A.; Masaguer, A.; Ruiz-Fernández, J. Phytostabilization of metals in mine soils using *Brassica juncea* in combination with organic amendments. *Plant Soil* **2014**, *377*, 97–109. [[CrossRef](#)]
64. McGrath, S.P.; Zhao, F.J.; Lombi, E. Plant and rhizosphere processes involved in phytoremediation of metal-contaminated soils. *Plant Soil* **2001**, *232*, 207–214. [[CrossRef](#)]
65. Sheng, X.F.; Xia, J.J.; Jiang, C.Y.; He, L.Y.; Qian, M. Characterization of heavy metal-resistant endophytic bacteria from rape (*Brassica napus*) roots and their potential in promoting the growth and lead accumulation of rape. *Environ. Pollut.* **2008**, *156*, 1164–1170. [[CrossRef](#)]
66. Mastretta, C.; Taghavi, S.; Van Der Lelie, D.; Mengoni, A.; Galardi, F.; Gonnelli, C.; Barac, T.; Boulet, J.; Weyens, N.; Vangronsveld, J. Endophytic bacteria from seeds of *Nicotiana tabacum* can reduce cadmium phytotoxicity. *Int. J. Phytoremediat.* **2009**, *11*, 251–267. [[CrossRef](#)]
67. Quiñones, M.A.; Ruiz-Díez, B.; Fajardo, S.; López-Berdonces, M.A.; Higuera, P.L.; Fernández-Pascual, M. *Lupinus albus* plants acquire mercury tolerance when inoculated with an Hg-resistant *Bradyrhizobium* strain. *Plant Physiol. Biochem.* **2013**, *73*, 168–175. [[CrossRef](#)]
68. Rock, S.; Pivetz, B.; Madalinski, K.; Adams, N.; Wilson, T. *Introduction to Phytoremediation*; U.S. Environmental Protection Agency: Washington, DC, USA, 2019.
69. Peco, J.D.; Higuera, P.; Campos, J.A.; Olmedilla, A.; Romero-Puertas, M.C.; Sandalio, L.M. Deciphering lead tolerance mechanisms in a population of the plant species *Biscutella auriculata* L. from a mining area: Accumulation strategies and antioxidant defenses. *Chemosphere* **2020**, *261*, 127721. [[CrossRef](#)]
70. Peco, J.D.; Sandalio, L.M.; Higuera, P.; Olmedilla, A.; Campos, J.A. Characterization of the biochemical basis for copper homeostasis and tolerance in *Biscutella auriculata* L. *Physiol. Plant.* **2020**. [[CrossRef](#)]
71. Reeves, R.D.; Baker, A.J.M.; Jaffré, T.; Erskine, P.D.; Echevarria, G.; van der Ent, A. A global database for plants that hyperaccumulate metal and metalloids trace elements. *New Phytol.* **2018**, *218*, 407–411. [[CrossRef](#)]
72. Pošćić, F.; Marchiol, L.; Schat, H. Hyperaccumulation of thallium is population-specific and uncorrelated with caesium accumulation in the thallium hyperaccumulator, *Biscutella laevigata*. *Plant Soil* **2013**, *365*, 81–91. [[CrossRef](#)]
73. McIntyre, T. Phytoremediation of heavy metals from soils. *Adv. Biochem. Eng. Biotechnol.* **2003**, *78*, 97–123. [[PubMed](#)]
74. Pollard, A.J.; Powell, K.D.; Harper, F.A.; Smith, J.A.C. The genetic basis of metal hyperaccumulation in plants. *Crit. Rev. Plant Sci.* **2002**, *21*, 539–566. [[CrossRef](#)]
75. Ke, W.; Xiong, Z.T.; Chen, S.; Chen, J. Effects of copper and mineral nutrition on growth, copper accumulation and mineral element uptake in two *Rumex japonicus* populations from a copper mine and an uncontaminated field sites. *Environ. Exp. Bot.* **2007**, *59*, 59–67. [[CrossRef](#)]
76. Kabata-Pendias, A. *Trace Elements in Soils and Plants*; CRC press: Boca Raton, FL, USA, 2001; ISBN 0849315751.
77. Wang, X.P.; Shan, X.Q.; Zhang, S.Z.; Wen, B. A model for evaluation of the phytoavailability of trace elements to vegetables under the field conditions. *Chemosphere* **2004**, *55*, 811–822. [[CrossRef](#)]
78. Kim, K.-R.; Owens, G.; Naidu, R. Heavy metal distribution, bioaccessibility, and phytoavailability in long-term contaminated soils from Lake Macquarie, Australia. *Soil Res.* **2009**, *47*, 166. [[CrossRef](#)]
79. Filipović, L.; Romić, M.; Romić, D.; Filipović, V.; Ondrašek, G. Organic matter and salinity modify cadmium soil (phyto) availability. *Ecotoxicol. Environ. Saf.* **2018**, *147*, 824–831. [[CrossRef](#)]
80. Stafford, A.; Jeyakumar, P.; Hedley, M.; Anderson, C. Influence of Soil Moisture Status on Soil Cadmium Phytoavailability and Accumulation in Plantain (*Plantago lanceolata*). *Soil Syst.* **2018**, *2*, 9. [[CrossRef](#)]
81. Ren, J.H.; Ma, L.Q.; Sun, H.J.; Cai, F.; Luo, J. Antimony uptake, translocation and speciation in rice plants exposed to antimonite and antimonate. *Sci. Total Environ.* **2014**, *475*, 83–89. [[CrossRef](#)]
82. Kim, R.Y.; Yoon, J.K.; Kim, T.S.; Yang, J.E.; Owens, G.; Kim, K.R. Bioavailability of heavy metals in soils: Definitions and practical implementation—a critical review. *Environ. Geochem. Health* **2015**, *37*, 1041–1061. [[CrossRef](#)]
83. Dong, J.; Wu, F.; Zhang, G. Influence of cadmium on antioxidant capacity and four microelement concentrations in tomato seedlings (*Lycopersicon esculentum*). *Chemosphere* **2006**, *64*, 1659–1666. [[CrossRef](#)]
84. Yang, C.M.; Juang, K.W. Alleviation effects of calcium and potassium on cadmium rhizotoxicity and absorption by soybean and wheat roots. *J. Plant Nutr. Soil Sci.* **2015**, *178*, 748–754. [[CrossRef](#)]
85. Tangahu, B.V.; Sheikh Abdullah, S.R.; Basri, H.; Idris, M.; Anuar, N.; Mukhlisin, M.; Sheikh Abdullah, S.R.; Idris, M.; Tangahu, B.V.; Basri, H. A Review on Heavy Metals (As, Pb, and Hg) Uptake by Plants through Phytoremediation. *Int. J. Chem. Eng.* **2011**, *2011*, 939161. [[CrossRef](#)]
86. Römheld, V.; Awad, F. Significance of root exudates in acquisition of heavy metals from a contaminated calcareous soil by graminaceous species. *J. Plant Nutr.* **2000**, *23*, 1857–1866. [[CrossRef](#)]

87. Kim, S.; Lim, H.; Lee, I. Enhanced heavy metal phytoextraction by *Echinochloa crus-galli* using root exudates. *J. Biosci. Bioeng.* **2010**, *109*, 47–50. [[CrossRef](#)]
88. Montiel-Rozas, M.M.; Madejón, E.; Madejón, P. Effect of heavy metals and organic matter on root exudates (low molecular weight organic acids) of herbaceous species: An assessment in sand and soil conditions under different levels of contamination. *Environ. Pollut.* **2016**, *216*, 273–281. [[CrossRef](#)]
89. Martin, M.H.; Marschner, H. The Mineral Nutrition of Higher Plants. *J. Ecol.* **1988**, *76*, 1250. [[CrossRef](#)]
90. Lombi, E.; Tearall, K.L.; Howarth, J.R.; Zhao, F.J.; Hawkesford, M.J.; McGrath, S.P. Influence of iron status on cadmium and zinc uptake by different ecotypes of the hyperaccumulator *Thlaspi caerulescens*. *Plant Physiol.* **2002**, *128*, 1359–1367. [[CrossRef](#)]
91. Chen, S.; Han, X.; Fang, J.; Lu, Z.; Qiu, W.; Liu, M.; Sang, J.; Jiang, J.; Zhuo, R. *Sedum alfredii* SaNramp6 Metal Transporter Contributes to Cadmium Accumulation in Transgenic *Arabidopsis thaliana*. *Sci. Rep.* **2017**, *7*, 13318. [[CrossRef](#)]
92. Feng, S.; Tan, J.; Zhang, Y.; Liang, S.; Xiang, S.; Wang, H.; Chai, T. Isolation and characterization of a novel cadmium-regulated Yellow Stripe-Like transporter (SnYSL3) in *Solanum nigrum*. *Plant Cell Rep.* **2017**, *36*, 281–296. [[CrossRef](#)]
93. Yruela, I. Copper in plants: Acquisition, transport and interactions. *Funct. Plant Biol.* **2009**, *36*, 409–430. [[CrossRef](#)]
94. Puig, S. Function and Regulation of the Plant COPT Family of High-Affinity Copper Transport Proteins. *Adv. Bot.* **2014**, *2014*, 476917. [[CrossRef](#)]
95. Andrés-Colás, N.; Sancenón, V.; Rodríguez-Navarro, S.; Mayo, S.; Thiele, D.J.; Ecker, J.R.; Puig, S.; Peñarrubia, L. The Arabidopsis heavy metal P-type ATPase HMA5 interacts with metallochaperones and functions in copper detoxification of roots. *Plant J.* **2006**, *45*, 225–236. [[CrossRef](#)]
96. Dai, J.; Wang, N.; Xiong, H.; Qiu, W.; Nakanishi, H.; Kobayashi, T.; Nishizawa, N.K.; Zuo, Y. The yellow stripe-like (YSL) gene functions in internal copper transport in peanut. *Genes* **2018**, *9*, 635. [[CrossRef](#)]
97. Uruguchi, S.; Mori, S.; Kuramata, M.; Kawasaki, A.; Arao, T.; Ishikawa, S. Root-to-shoot Cd translocation via the xylem is the major process determining shoot and grain cadmium accumulation in rice. *J. Exp. Bot.* **2009**, *60*, 2677–2688. [[CrossRef](#)]
98. Li, Q.; Guo, J.; Zhang, X.; Yu, H.; Huang, F.; Zhang, L.; Zhang, M.; Li, T. Changes of non-protein thiols in root and organic acids in xylem sap involved in cadmium translocation of cadmium-safe rice line (*Oryza sativa* L.). *Plant Soil* **2019**, *439*, 475–486. [[CrossRef](#)]
99. Kerkeb, L.; Krämer, U. The role of free histidine in xylem loading of nickel in *Alyssum lesbiacum* and *Brassica juncea*. *Plant Physiol.* **2003**, *131*, 716–724. [[CrossRef](#)]
100. Centofanti, T.; Sayers, Z.; Cabello-Conejo, M.I.; Kidd, P.; Nishizawa, N.K.; Kakei, Y.; Davis, A.P.; Sicher, R.C.; Chaney, R.L. Xylem exudate composition and root-to-shoot nickel translocation in *Alyssum* species. *Plant Soil* **2013**, *373*, 59–75. [[CrossRef](#)]
101. Saathoff, A.J.; Ahner, B.; Spanswick, R.M.; Walker, L.P. Detection of Phytochelatin in the Xylem Sap of *Brassica napus*. *Environ. Eng. Sci.* **2011**, *28*, 103–111. [[CrossRef](#)]
102. Wei, Z.G.; Wong, J.W.C.; Zhao, H.Y.; Zhang, H.J.; Li, H.X.; Hu, F. Separation and determination of heavy metals associated with low molecular weight chelators in xylem saps of Indian mustard (*Brassica juncea*) by size exclusion chromatography and atomic absorption spectrometry. *Biol. Trace Elem. Res.* **2007**, *118*, 146–158. [[CrossRef](#)]
103. Yadav, S.K.K. Heavy metals toxicity in plants: An overview on the role of glutathione and phytochelatins in heavy metal stress tolerance of plants. *S. Afr. J. Bot.* **2010**, *76*, 167–179. [[CrossRef](#)]
104. Dago, À.; González, I.; Ariño, C.; Martínez-Coronado, A.; Higuera, P.; Díaz-Cruz, J.M.; Esteban, M. Evaluation of mercury stress in plants from the Almadén mining district by analysis of phytochelatin and their Hg complexes. *Environ. Sci. Technol.* **2014**, *48*, 6256–6263. [[CrossRef](#)] [[PubMed](#)]
105. Dago, A.; Ariño, C.; Díaz-Cruz, J.M.; Esteban, M. Analysis of phytochelatin and Hg-phytochelatin complexes in *Hordeum vulgare* plants stressed with Hg and Cd: HPLC study with amperometric detection. *Int. J. Environ. Anal. Chem.* **2014**, *94*, 668–678. [[CrossRef](#)]
106. Heiss, S.; Wachter, A.; Bogs, J.; Cobbett, C.; Rausch, T. Phytochelatin synthase (PCS) protein is induced in *Brassica juncea* leaves after prolonged Cd exposure. *J. Exp. Bot.* **2003**, *54*, 1833–1839. [[CrossRef](#)]
107. Liu, W.; Zhang, X.; Liang, L.; Chen, C.; Wei, S.; Zhou, Q. Phytochelatin and oxidative stress under heavy metal stress tolerance in plants. In *Reactive Oxygen Species and Oxidative Damage in Plants under Stress*; Springer International Publishing: Cham, Switzerland, 2015; pp. 191–217. ISBN 9783319204215.
108. Grill, E.; Zenk, M.H.; Winnacker, E.-L. Induction of heavy metal-sequestering phytochelatin by cadmium in cell cultures of *Rauvolfia serpentina*. *Naturwissenschaften* **1985**, *72*, 432–433. [[CrossRef](#)]
109. Klapheck, S.; Chrost, B.; Starke, J.; Zimmermann, H. γ -Glutamylcysteinylserine—A New Homologue of Glutathione in Plants of the Family Poaceae. *Bot. Acta* **1992**, *105*, 174–179. [[CrossRef](#)]
110. Meuwly, P.; Thibault, P.; Rauser, W.E. γ -Glutamylcysteinylglutamic acid—a new homologue of glutathione in maize seedlings exposed to cadmium. *FEBS Lett.* **1993**, *336*, 472–476. [[CrossRef](#)]
111. Kubota, H.; Sato, K.; Yamada, T.; Maitani, T. Phytochelatin homologs induced in hairy roots of horseradish. *Phytochemistry* **2000**, *53*, 239–245. [[CrossRef](#)]
112. Sylwia, W.; Anna, R.; Ewa, B.; Stephan, C.; Danuta Maria, A. The role of subcellular distribution of cadmium and phytochelatin in the generation of distinct phenotypes of AtPCS1- and CePCS3-expressing tobacco. *J. Plant Physiol.* **2010**, *167*, 981–988. [[CrossRef](#)]
113. Yang, X.E.; Li, T.Q.; Long, X.X.; Xiong, Y.H.; He, Z.L.; Stoffella, P.J. Dynamics of zinc uptake and accumulation in the hyperaccumulating and non-hyperaccumulating ecotypes of *Sedum alfredii* Hance. *Plant Soil* **2006**, *284*, 109–119. [[CrossRef](#)]

114. Melo, E.E.C.; Costa, E.T.S.; Guilherme, L.R.G.; Faquin, V.; Nascimento, C.W.A. Accumulation of arsenic and nutrients by castor bean plants grown on an As-enriched nutrient solution. *J. Hazard. Mater.* **2009**, *168*, 479–483. [[CrossRef](#)]
115. Pandey, S.; Fartyal, D.; Agarwal, A.; Shukla, T.; James, D.; Kaul, T.; Negi, Y.K.; Arora, S.; Reddy, M.K. Abiotic Stress Tolerance in Plants: Myriad Roles of Ascorbate Peroxidase. *Front. Plant Sci.* **2017**, *8*, 581. [[CrossRef](#)]
116. Fuentes, A. Phytotoxicity and heavy metals speciation of stabilised sewage sludges. *J. Hazard. Mater.* **2004**, *108*, 161–169. [[CrossRef](#)]
117. Sandalio, L.M.; Dalurzo, H.C.; Gómez, M.; Romero-Puertas, M.C.; del Río, L.A. Cadmium-induced changes in the growth and oxidative metabolism of pea plants. *J. Exp. Bot.* **2001**, *52*, 2115–2126. [[CrossRef](#)]
118. Sharma, P.; Jha, A.B.; Dubey, R.S.; Pessarakli, M. Reactive Oxygen Species, Oxidative Damage, and Antioxidative Defense Mechanism in Plants under Stressful Conditions. *J. Bot.* **2012**, *2012*, 1–26. [[CrossRef](#)]
119. Kalaivanan, D.; Ganeshamurthy, A.N. Mechanisms of Heavy Metal Toxicity in Plants. In *Abiotic Stress Physiology of Horticultural Crops*; Springer: New Delhi, India, 2016; pp. 85–102.
120. Nagajyoti, P.C.; Lee, K.D.; Sreekanth, T.V.M. Heavy metals, occurrence and toxicity for plants: A review. *Environ. Chem. Lett.* **2010**, *8*, 199–216. [[CrossRef](#)]
121. Manzoor, J.; Sharma, M.; Wani, K.A. Heavy metals in vegetables and their impact on the nutrient quality of vegetables: A review. *J. Plant Nutr.* **2018**, *41*, 1744–1763. [[CrossRef](#)]
122. Sandalio, L.M.; Rodríguez-Serrano, M.; Gupta, D.K.; Archilla, A.; Romero-Puertas, M.C.; Del Río, L.A. Reactive Oxygen species and Nitric Oxide in plants under cadmium stress: From toxicity to signaling. In *Environmental Adaptations and Stress Tolerance of Plants in the Era of Climate Change*; Springer: New York, NY, USA, 2012; pp. 199–215. ISBN 9781461408154.
123. Molina-Moya, E.; Terrón-Camero, L.C.; Pescador-Azofra, L.; Sandalio, L.M.; Romero-Puertas, M.C. Reactive Oxygen Species and Nitric Oxide Production, Regulation and Function During Defense Response. In *Reactive Oxygen, Nitrogen and Sulfur Species in Plants*; Wiley: Hoboken, NJ, USA, 2019; pp. 573–590.
124. Romero-Puertas, M.C.; Terrón-Camero, L.C.; Peláez-Vico, M.Á.; Olmedilla, A.; Sandalio, L.M. Reactive oxygen and nitrogen species as key indicators of plant responses to Cd stress. *Environ. Exp. Bot.* **2019**, *161*, 107–119. [[CrossRef](#)]
125. Gupta, A.S.; Webb, R.P.; Holaday, A.S.; Allen, R.D. Overexpression of superoxide dismutase protects plants from oxidative stress. Induction of ascorbate peroxidase in superoxide dismutase-overexpressing plants. *Plant Physiol.* **1993**, *103*, 1067–1073. [[CrossRef](#)]
126. Guo, Q.; Meng, L.; Zhang, Y.-N.N.; Mao, P.-C.C.; Tian, X.-X.X.; Li, S.-S.S.; Zhang, L. Antioxidative systems, metal ion homeostasis and cadmium distribution in *Iris lactea* exposed to cadmium stress. *Ecotoxicol. Environ. Saf.* **2017**, *139*, 50–55. [[CrossRef](#)]
127. Gong, Q.; Wang, L.; Dai, T.; Zhou, J.; Kang, Q.; Chen, H.; Li, K.; Li, Z. Effects of copper on the growth, antioxidant enzymes and photosynthesis of spinach seedlings. *Ecotoxicol. Environ. Saf.* **2019**, *171*, 771–780. [[CrossRef](#)]
128. Xun, Y.; Feng, L.; Li, Y.; Dong, H. Mercury accumulation plant *Cyrtomium macrophyllum* and its potential for phytoremediation of mercury polluted sites. *Chemosphere* **2017**, *189*, 161–170. [[CrossRef](#)]
129. Singh, S.; Sounderajan, S.; Kumar, K.; Fulzele, D.P. Investigation of arsenic accumulation and biochemical response of in vitro developed *Vetiveria zizanoides* plants. *Ecotoxicol. Environ. Saf.* **2017**, *145*, 50–56. [[CrossRef](#)]
130. Ortega, A.; Garrido, I.; Casimiro, I.; Espinosa, F. Effects of antimony on redox activities and antioxidant defence systems in sunflower (*Helianthus annuus* L.) plants. *PLoS ONE* **2017**, *12*, e0183991. [[CrossRef](#)]
131. Zaheer, I.E.; Ali, S.; Saleem, M.H.; Imran, M.; Alnusairi, G.S.H.; Alharbi, B.M.; Riaz, M.; Abbas, Z.; Rizwan, M.; Soliman, M.H. Role of iron-lysine on morpho-physiological traits and combating chromium toxicity in rapeseed (*Brassica napus* L.) plants irrigated with different levels of tannery wastewater. *Plant Physiol. Biochem.* **2020**, *155*, 70–84. [[CrossRef](#)]
132. Singh, P.; Shukla, A.K.; Behera, S.K.; Tiwari, P.K. Zinc Application Enhances Superoxide Dismutase and Carbonic Anhydrase Activities in Zinc-Efficient and Zinc-Inefficient Wheat Genotypes. *J. Soil Sci. Plant Nutr.* **2019**, *19*, 477–487. [[CrossRef](#)]
133. Zhu, Y.M.; Wu, Q.H.; Lv, H.Q.; Chen, W.X.; Wang, L.Z.; Shi, S.J.; Yang, J.G.; Zhao, P.P.; Li, Y.P.; Christopher, R.; et al. Toxicity of different forms of antimony to rice plants: Effects on reactive oxidative species production, antioxidative systems, and uptake of essential elements. *Environ. Pollut.* **2020**, *263*, 114544. [[CrossRef](#)]
134. Pirzadah, T.B.; Malik, B.; Tahir, I.; Irfan, Q.M.; Rehman, R.U. Characterization of mercury-induced stress biomarkers in *Fagopyrum tataricum* plants. *Int. J. Phytoremediat.* **2018**, *20*, 225–236. [[CrossRef](#)]
135. Corpas, F.J.; Barroso, J.B.; Sandalio, L.M.; Distefano, S.; Palma, J.M.; Lupiáñez, J.A.; Del Río, L.A. A dehydrogenase-mediated recycling system of NADPH in plant peroxisomes. *Biochem. J.* **1998**, *330 Pt 2*, 777–784. [[CrossRef](#)]
136. Rodríguez-Ruiz, M.; Aparicio-Chacón, M.V.; Palma, J.M.; Corpas, F.J. Arsenate disrupts ion balance, sulfur and nitric oxide metabolisms in roots and leaves of pea (*Pisum sativum* L.) plants. *Environ. Exp. Bot.* **2019**, *161*, 143–156. [[CrossRef](#)]
137. Kumar, S.; Trivedi, P.K. Glutathione S-transferases: Role in combating abiotic stresses including arsenic detoxification in plants. *Front. Plant Sci.* **2018**, *9*, 751. [[CrossRef](#)] [[PubMed](#)]
138. Namdjoyan, S.; Kermanian, H.; Abolhasani Soorki, A.; Modarres Tabatabaei, S.; Elyasi, N. Interactive effects of Salicylic acid and nitric oxide in alleviating zinc toxicity of Safflower (*Carthamus tinctorius* L.). *Ecotoxicology* **2017**, *26*, 752–761. [[CrossRef](#)] [[PubMed](#)]
139. Handa, N.; Kohli, S.K.; Sharma, A.; Thukral, A.K.; Bhardwaj, R.; Abd_Allah, E.F.; Alqarawi, A.A.; Ahmad, P. Selenium modulates dynamics of antioxidative defence expression, photosynthetic attributes and secondary metabolites to mitigate chromium toxicity in *Brassica juncea* L. plants. *Environ. Exp. Bot.* **2019**, *161*, 180–192. [[CrossRef](#)]
140. Noctor, G.; Mhamdi, A.; Chaouch, S.; Han, Y.; Neukermans, J.; Marquez-Garcia, B.; Queval, G.; Foyer, C.H. Glutathione in plants: An integrated overview. *Plant. Cell Environ.* **2012**, *35*, 454–484. [[CrossRef](#)]

141. Terrón-Camero, L.C.; Peláez-Vico, M.Á.; Del-Val, C.; Sandalio, L.M.; Romero-Puertas, M.C.; Brouquisse, R. Role of nitric oxide in plant responses to heavy metal stress: Exogenous application versus endogenous production. *J. Exp. Bot.* **2019**, *70*, 4477–4488. [[CrossRef](#)]
142. Maestri, E.; Marmioli, N. Transgenic plants for phytoremediation. *Int. J. Phytoremediat.* **2011**, *13*, 264–279. [[CrossRef](#)]
143. Cherian, S.; Oliveira, M.M. Transgenic plants in phytoremediation: Recent advances and new possibilities. *Environ. Sci. Technol.* **2005**, *39*, 9377–9390. [[CrossRef](#)]
144. Gunarathne, V.; Mayakaduwa, S.; Ashiq, A.; Weerakoon, S.R.; Biswas, J.K.; Vithanage, M. Transgenic Plants: Benefits, Applications, and Potential Risks in Phytoremediation. Benefits, Applications, and Potential Risks in Phytoremediation. In *Transgenic Plant Technology for Remediation of Toxic Metals and Metalloids*; Elsevier: Amsterdam, The Netherlands, 2018; pp. 89–102; ISBN 9780128143902.
145. Liu, D.; An, Z.; Mao, Z.; Ma, L.; Lu, Z. Enhanced Heavy Metal Tolerance and Accumulation by Transgenic Sugar Beets Expressing *Streptococcus thermophilus* StGCS-GS in the Presence of Cd, Zn and Cu Alone or in Combination. *PLoS ONE* **2015**, *10*, e0128824. [[CrossRef](#)]
146. Liu, Y.; Kang, T.; Cheng, J.S.; Yi, Y.J.; Han, J.J.; Cheng, H.L.; Li, Q.; Tang, N.; Liang, M.X. Heterologous expression of the metallothionein PpMT2 gene from *Physcomitrella patens* confers enhanced tolerance to heavy metal stress on transgenic *Arabidopsis* plants. *Plant Growth Regul.* **2020**, *90*, 63–72. [[CrossRef](#)]
147. Evans, D.M.; Zipper, C.E.; Burger, J.A.; Strahm, B.D.; Villamagna, A.M. Reforestation practice for enhancement of ecosystem services on a compacted surface mine: Path toward ecosystem recovery. *Ecol. Eng.* **2013**, *51*, 16–23. [[CrossRef](#)]
148. Madejón, P.; Domínguez, M.T.; Madejón, E.; Cabrera, F.; Marañón, T.; Murillo, J.M. Soil-plant relationships and contamination by trace elements: A review of twenty years of experimentation and monitoring after the Aznalcóllar (SW Spain) mine accident. *Sci. Total Environ.* **2018**, *625*, 50–63. [[CrossRef](#)]
149. Del Río, M.; Font, R.; Almela, C.; Vélez, D.; Montoro, R.; De Haro Bailón, A. Heavy metals and arsenic uptake by wild vegetation in the Guadiamar river area after the toxic spill of the Aznalcóllar mine. In Proceedings of the Journal of Biotechnology. *J. Biotechnol.* **2002**, *98*, 125–137. [[CrossRef](#)]
150. Madejón, P.; Murillo, J.M.; Marañón, T.; Cabrera, F. Bioaccumulation of trace elements in a wild grass three years after the Aznalcóllar mine spill (South Spain). *Environ. Monit. Assess.* **2006**, *114*, 169–189. [[CrossRef](#)]
151. Madejón, P.; Murillo, J.M.; Marañón, T.; Lepp, N.W. Factors affecting accumulation of thallium and other trace elements in two wild *Brassicaceae* spontaneously growing on soils contaminated by tailings dam waste. *Chemosphere* **2007**, *67*, 20–28. [[CrossRef](#)]
152. Murillo, J.M.; Marañón, T.; Cabrera, F.; López, R. Accumulation of heavy metals in sunflower and sorghum plants affected by the Guadiamar spill. *Sci. Total Environ.* **1999**, *242*, 281–292. [[CrossRef](#)]
153. Madejón, P.; Murillo, J.M.; Marañón, T.; Cabrera, F.; Soriano, M.A. Trace element and nutrient accumulation in sunflower plants two years after the Aznalcóllar mine spill. *Sci. Total Environ.* **2003**, *307*, 239–257. [[CrossRef](#)]
154. Clemente, R.; Walker, D.J.; Bernal, M.P. Uptake of heavy metals and As by *Brassica juncea* grown in a contaminated soil in Aznalcóllar (Spain): The effect of soil amendments. *Environ. Pollut.* **2005**, *138*, 46–58. [[CrossRef](#)]
155. Soriano, M.A.; Fereres, E. Use of crops for in situ phytoremediation of polluted soils following a toxic flood from a mine spill. *Plant Soil* **2003**, *256*, 253–264. [[CrossRef](#)]
156. Vázquez, S.; Agha, R.; Granada, A.; Sarro, M.J.; Esteban, E.; Peñalosa, J.M.; Carpena, R.O. Use of white lupin plant for phytostabilization of Cd and as polluted acid soil. *Water Air Soil Pollut.* **2006**, *177*, 349–365. [[CrossRef](#)]
157. Moreno-Jiménez, E.; Vázquez, S.; Carpena-Ruiz, R.O.; Esteban, E.; Peñalosa, J.M. Using Mediterranean shrubs for the phytoremediation of a soil impacted by pyritic wastes in Southern Spain: A field experiment. *J. Environ. Manag.* **2011**, *92*, 1584–1590. [[CrossRef](#)]
158. Domínguez, M.T.; Marañón, T.; Murillo, J.M.; Schulin, R.; Robinson, B.H. Trace element accumulation in woody plants of the Guadiamar Valley, SW Spain: A large-scale phytomanagement case study. *Environ. Pollut.* **2008**, *152*, 50–59. [[CrossRef](#)]
159. Domínguez, M.T.; Madejón, P.; Marañón, T.; Murillo, J.M. Avorestation of a trace-element polluted area in SW Spain: Woody plant performance and trace element accumulation. *Eur. J. For. Res.* **2010**, *129*, 47–59. [[CrossRef](#)]
160. Domínguez, M.T.; Pérez-Ramos, I.M.; Murillo, J.M.; Marañón, T. Facilitating the afforestation of Mediterranean polluted soils by nurse shrubs. *J. Environ. Manag.* **2015**, *161*, 276–286. [[CrossRef](#)]
161. De la Fuente, C.; Pardo, T.; Alburquerque, J.A.; Martínez-Alcalá, I.; Bernal, M.P.; Clemente, R. Assessment of native shrubs for stabilisation of a trace elements-polluted soil as the final phase of a restoration process. *Agric. Ecosyst. Environ.* **2014**, *196*, 103–111. [[CrossRef](#)]
162. Domínguez, M.T.; Madrid, F.; Marañón, T.; Murillo, J.M. Cadmium availability in soil and retention in oak roots: Potential for phytostabilization. *Chemosphere* **2009**, *76*, 480–486. [[CrossRef](#)]
163. Moreno-Jiménez, E.; Peñalosa, J.M.; Carpena-Ruiz, R.O.; Esteban, E. Comparison of arsenic resistance in Mediterranean woody shrubs used in restoration activities. *Chemosphere* **2008**, *71*, 466–473. [[CrossRef](#)]
164. Martín-Crespo, T.; Gómez-Ortiz, D.; Martín-Velázquez, S.; Esbrí, J.M.; de Ignacio-San José, C.; Sánchez-García, M.J.; Montoya-Montes, I.; Martín-González, F. Abandoned mine tailings in cultural itineraries: Don Quixote Route (Spain). *Eng. Geol.* **2015**, *197*, 82–93. [[CrossRef](#)]
165. Asensio, V.; Vega, F.A.; Singh, B.R.; Covelo, E.F. Effects of tree vegetation and waste amendments on the fractionation of Cr, Cu, Ni, Pb and Zn in polluted mine soils. *Sci. Total Environ.* **2013**, *443*, 446–453. [[CrossRef](#)] [[PubMed](#)]

166. Pardo, T.; Bernal, M.P.; Clemente, R. Phytostabilisation of severely contaminated mine tailings using halophytes and field addition of organic and inorganic amendments. *Chemosphere* **2017**, *178*, 556–564. [[CrossRef](#)]
167. Hernández, A.; Jébrak, M.; Higuera, P.; Oyarzun, R.; Morata, D.; Munhá, J. The Almaden mercury mining district, Spain. *Miner. Depos.* **1999**, *34*, 539–548. [[CrossRef](#)]
168. Rodríguez, L.; Lopez-Bellido, F.J.; Carnicer, A.; Recreo, F.; Tallos, A.; Monteagudo, J.M. Mercury recovery from soils by phytoremediation. In *Environmental Chemistry: Green Chemistry and Pollutants in Ecosystems*; Springer: Berlin/Heidelberg, Germany, 2005; pp. 197–204; ISBN 3540228608.
169. Naharro, R.; Esbrí, J.M.; Amorós, J.Á.; García-Navarro, F.J.; Higuera, P. Assessment of mercury uptake routes at the soil-plant-atmosphere interface. *Geochem. Explor. Environ. Anal.* **2018**, *19*, 146–154. [[CrossRef](#)]
170. Pratas, J.; Prasad, M.N.V.; Freitas, H.; Conde, L. Plants growing in abandoned mines of Portugal are useful for biogeochemical exploration of arsenic, antimony, tungsten and mine reclamation. *J. Geochem. Explor.* **2005**, *85*, 99–107. [[CrossRef](#)]
171. Simiele, M.; Lebrun, M.; Miard, F.; Trupiano, D.; Poupart, P.; Forestier, O.; Scippa, G.S.; Bourgerie, S.; Morabito, D. Assisted phytoremediation of a former mine soil using biochar and iron sulphate: Effects on As soil immobilization and accumulation in three Salicaceae species. *Sci. Total Environ.* **2020**, *710*, 136203. [[CrossRef](#)] [[PubMed](#)]
172. Bacchetta, G.; Cao, A.; Cappai, G.; Carucci, A.; Casti, M.; Fercia, M.L.; Lonis, R.; Mola, F. A field experiment on the use of *Pistacia lentiscus* L. and *Scrophularia canina* L. subsp. bicolor (Sibth. et Sm.) Greater for the phytoremediation of abandoned mining areas. *Plant Biosyst.* **2012**, *146*, 1054–1063. [[CrossRef](#)]
173. Swab, R.M.; Lorenz, N.; Byrd, S.; Dick, R. Native vegetation in reclamation: Improving habitat and ecosystem function through using prairie species in mine land reclamation. *Ecol. Eng.* **2017**, *108*, 525–536. [[CrossRef](#)]
174. Gil-Loaiza, J.; White, S.A.; Root, R.A.; Solís-Dominguez, F.A.; Hammond, C.M.; Chorover, J.; Maier, R.M. Phytostabilization of mine tailings using compost-assisted direct planting: Translating greenhouse results to the field. *Sci. Total Environ.* **2016**, *565*, 451–461. [[CrossRef](#)]
175. Zhao, Z.; Shahrour, I.; Bai, Z.; Fan, W.; Feng, L.; Li, H. Soils development in opencast coal mine spoils reclaimed for 1–13 years in the West-Northern Loess Plateau of China. *Eur. J. Soil Biol.* **2013**, *55*, 40–46. [[CrossRef](#)]
176. Zhang, P.; Yang, F.; Lan, M.; Liu, W.; Yang, W.; Tang, Y.; Qiu, R. Phytostabilization with tolerant plants and soil amendments of the tailings of the Dabaoshan polymetallic mine in Guangdong Province. *Huanjing Kexue Xuebao/Acta Sci. Circumstantiae* **2019**, *39*, 545–552.
177. Tai, Y.P.; Yang, Y.F.; Li, Z.A.; Yang, Y.; Wang, J.X.; Zhuang, P.; Zou, B. Phytoextraction of 55-year-old wastewater-irrigated soil in a Zn-Pb mine district: Effect of plant species and chelators. *Environ. Technol.* **2018**, *39*, 2138–2150. [[CrossRef](#)]
178. Li, X.; Wang, X.; Chen, Y.; Yang, X.; Cui, Z. Optimization of combined phytoremediation for heavy metal contaminated mine tailings by a field-scale orthogonal experiment. *Ecotoxicol. Environ. Saf.* **2019**, *168*, 1–8. [[CrossRef](#)]
179. Yu, G.; Jiang, P.; Fu, X.; Liu, J.; Sunahara, G.I.; Chen, Z.; Xiao, H.; Lin, F.; Wang, X. Phytoextraction of cadmium-contaminated soil by *Celosia argentea* Linn.: A long-term field study. *Environ. Pollut.* **2020**, *266*, 115408. [[CrossRef](#)]
180. Yuan, X.; Xiong, T.; Yao, S.; Liu, C.; Yin, Y.; Li, H.; Li, N. A real field phytoremediation of multi-metals contaminated soils by selected hybrid sweet sorghum with high biomass and high accumulation ability. *Chemosphere* **2019**, *237*, 124536. [[CrossRef](#)]
181. Shen, S.; Chen, J.; Chang, J.; Xia, B. Using bioenergy crop cassava (*Manihot esculenta*) for reclamation of heavily metal-contaminated land. *Int. J. Phytoremediat.* **2020**, *22*, 1313–1320. [[CrossRef](#)]
182. Shu, W.S.; Xia, H.P.; Zhang, Z.Q.; Lan, C.Y.; Wong, M.H. Use of vetiver and three other grasses for revegetation of Pb/Zn mine tailings: Field experiment. *Int. J. Phytoremediat.* **2002**, *4*, 47–57. [[CrossRef](#)]
183. Liu, Y.; Zhong, X.; Huot, H.; Liu, W.; Liu, C.; Guo, M.; Li, Y.; Fei, Y.; Chao, Y.; Wang, S.; et al. Reclamation with organic amendments and plants remodels the diversity and structure of bacterial community in ion-adsorption rare earth element mine tailings. *J. Soils Sediments* **2020**, *20*, 3669–3680. [[CrossRef](#)]
184. Mi, J.; Hou, H.; Raval, S.; Yang, Y.; Zhang, S.; Hua, Y.; Wang, C.; Chen, F. Effect of crop cultivation on the soil carbon stock in mine dumps of the Loess Plateau, China. *Sci. Total Environ.* **2020**, *741*, 139809. [[CrossRef](#)]
185. Chen, F.; Yang, Y.; Mi, J.; Liu, R.; Hou, H.; Zhang, S. Effects of vegetation pattern and spontaneous succession on remediation of potential toxic metal-polluted soil in mine dumps. *Sustainability* **2019**, *11*, 397. [[CrossRef](#)]
186. Singh, A.K. Toxic aqueous discharge of iron and sulphur from spoiled coal mined lands and its control by phytostabilization process. *Curr. Sci.* **2018**, *115*, 529–534. [[CrossRef](#)]
187. Kamble, S.S.; Dod, R.D. Management of coal mine overburden and fly ash using bamboo cultivation. *Int. J. Innov. Technol. Explor. Eng.* **2019**, *8*, 4223–4226.
188. Jeżowski, S.; Mos, M.; Buckby, S.; Cerażny-Waliszewska, J.; Owczarzak, W.; Mocek, A.; Kaczmarek, Z.; McCalmont, J.P. Establishment, growth, and yield potential of the perennial grass *Miscanthus* × *Giganteus* on degraded coal mine soils. *Front. Plant Sci.* **2017**, *8*, 726. [[CrossRef](#)]
189. Bandyopadhyay, S.; Rana, V.; Maiti, S.K. Chronological Variation of Metals in Reclaimed Coal Mine Soil and Tissues of *Eucalyptus* Hybrid Tree After 25 Years of Reclamation, Jharia Coal Field (India). *Bull. Environ. Contam. Toxicol.* **2018**, *101*, 604–610. [[CrossRef](#)]
190. Pavlova, D.; Echevarria, G.; Mullaj, A.; Reeves, R.D.; Morel, J.-L.; Sulçe, S. Nickel hyperaccumulation by the species of *Alyssum* and *Thlaspi* (*Brassicaceae*) from the ultramafic soils of the Balkans. *Bot. Serbica* **2010**, *34*, 3–10.
191. Küpper, H.; Lombi, E.; Zhao, F.J.; McGrath, S.P. Cellular compartmentation of cadmium and zinc in relation to other elements in the hyperaccumulator *Arabidopsis halleri*. *Planta* **2000**, *212*, 75–84. [[CrossRef](#)]

192. Kubota, H.; Takenaka, C. *Arabis gemmifera* is a Hyperaccumulator of Cd and Zn. *Int. J. Phytoremediat.* **2003**, *5*, 197–201. [[CrossRef](#)]
193. Mesnoua, M.; Mateos-Naranjo, E.; Barcia-Piedras, J.M.; Pérez-Romero, J.A.; Lotmani, B.; Redondo-Gómez, S. Physiological and biochemical mechanisms preventing Cd-toxicity in the hyperaccumulator *Atriplex halimus* L. *Plant Physiol. Biochem.* **2016**, *106*, 30–38. [[CrossRef](#)]
194. Ugustyniak, M.; Arnawska, M.; Eimold, W. Uptake of cadmium, lead, nickel and zinc from soil and water solutions by the nickel hyperaccumulator. *Acta Biol. Cracoviensia* **2004**, *46*, 75–85.
195. Sun, Y.; Zhou, Q.; Wang, L.; Liu, W. Cadmium tolerance and accumulation characteristics of *Bidens pilosa* L. as a potential Cd-hyperaccumulator. *J. Hazard. Mater.* **2009**, *161*, 808–814. [[CrossRef](#)]
196. Wierzbicka, M.; Pielichowska, M.; Abratowska, A.; Wiłkomirski, B.; Wysocka, I.; Panufnik-Mędrzycka, D.; Bulska, E.; Panufnik-Mędrzycka, D.; Bulska, E. Thallium Hyperaccumulation in Polish Populations of *Biscutella laevigata* (Brassicaceae). *Acta Biol. Cracoviensia Ser. Bot.* **2016**, *58*, 7–19. [[CrossRef](#)]
197. Xiong, Z.T. Lead uptake and effects on seed germination and plant growth in a Pb hyperaccumulator *Brassica pekinensis* Rupr. *Bull. Environ. Contam. Toxicol.* **1998**, *60*, 285–291. [[CrossRef](#)]
198. Liu, W.; Zhang, C.; Hu, P.; Luo, Y.; Wu, L.; Sale, P.; Tang, C. Influence of nitrogen form on the phytoextraction of cadmium by a newly discovered hyperaccumulator *Carpobrotus rossii*. *Environ. Sci. Pollut. Res.* **2016**, *23*, 1246–1253. [[CrossRef](#)]
199. Zhang, X.H.; Liu, J.; Huang, H.T.; Chen, J.; Zhu, Y.N.; Wang, D.Q. Chromium accumulation by the hyperaccumulator plant *Leersia hexandra* Swartz. *Chemosphere* **2007**, *67*, 1138–1143. [[CrossRef](#)] [[PubMed](#)]
200. Liu, Z.; He, X.; Chen, W.; Yuan, F.; Yan, K.; Tao, D. Accumulation and tolerance characteristics of cadmium in a potential hyperaccumulator—*Lonicera japonica* Thunb. *J. Hazard. Mater.* **2009**, *169*, 170–175. [[CrossRef](#)] [[PubMed](#)]
201. Xue, S.G.; Chen, Y.X.; Reeves, R.D.; Baker, A.J.; Lin, Q.; Fernando, D.R. Manganese uptake and accumulation by the hyperaccumulator plant *Phytolacca acinosa* Roxb. (Phytolaccaceae). *Environ. Pollut.* **2004**, *131*, 393–399. [[CrossRef](#)] [[PubMed](#)]
202. Liu, K.; Yu, F.; Chen, M.; Zhou, Z.; Chen, C.; Li, M.S.; Zhu, J. A newly found manganese hyperaccumulator—*Polygonum lapathifolium* Linn. *Int. J. Phytoremediat.* **2016**, *18*, 348–353. [[CrossRef](#)]
203. Chen, T.; Wei, C.; Huang, Z.; Huang, Q.; Lu, Q.; Fan, Z. Arsenic hyperaccumulator *Pteris vittata* L. and its arsenic accumulation. *Chin. Sci. Bull.* **2002**, *47*, 902–905. [[CrossRef](#)]
204. Li, M.S.; Yang, S.X. Heavy metal contamination in soils and phytoaccumulation in a manganese mine Wasteland, South China. *Air Soil Water Res.* **2008**, *1*, 31–41. [[CrossRef](#)]
205. Yang, S.X.; Deng, H.; Li, M.S. Manganese uptake and accumulation in a woody hyperaccumulator, *Schima superba*. *Plant Soil Environ.* **2008**, *54*, 441–446. [[CrossRef](#)]
206. Yang, X.E.; Long, X.X.; Ye, H.B.; He, Z.L.; Calvert, D.V.; Stoffella, P.J. Cadmium tolerance and hyperaccumulation in a new Zn-hyperaccumulating plant species (*Sedum alfredii* Hance). *Plant Soil* **2004**, *259*, 181–189. [[CrossRef](#)]
207. Yang, X.; Long, X.; Ni, W.; Fu, C. *Sedum alfredii* H: A new Zn hyperaccumulating plant first found in China. *Chin. Sci. Bull.* **2002**, *47*, 1634–1637. [[CrossRef](#)]
208. Sahi, S.V.; Bryant, N.L.; Sharma, N.C.; Singh, S.R. Characterization of a lead hyperaccumulator shrub, *Sesbania drummondii*. *Environ. Sci. Technol.* **2002**, *36*, 4676–4680. [[CrossRef](#)]
209. Gao, Y.; Zhou, P.; Mao, L.; Shi, W.J.; Zhi, Y.E. Phytoextraction of cadmium and physiological changes in *Solanum nigrum* as a novel cadmium hyperaccumulator. *Russ. J. Plant Physiol.* **2010**, *57*, 501–508. [[CrossRef](#)]
210. Baker, A.J.M.; Reeves, R.D.; Hajar, A.S.M. Heavy metal accumulation and tolerance in British populations of the metallophyte *Thlaspi caerulescens* J. & C. Presl (Brassicaceae). *New Phytol.* **1994**, *127*, 61–68.
211. Sakakibara, M.; Watanabe, A.; Inoue, M.; Sano, S.; Kaise, T. Phytoextraction and phytovolatilization of arsenic from As-contaminated soils by *Pteris vittata*. In Proceedings of the Annual International Conference on Soils, Sediments, Water and Energy, Amherst, MA, USA, 16–19 October 2010; Volume 12.
212. Verma, S.; Verma, P.K.; Pande, V.; Tripathi, R.D.; Chakrabarty, D. Transgenic *Arabidopsis thaliana* expressing fungal arsenic methyltransferase gene (WaarsM) showed enhanced arsenic tolerance via volatilization. *Environ. Exp. Bot.* **2016**, *132*, 113–120. [[CrossRef](#)]
213. Guarino, F.; Conte, B.; Improta, G.; Sciarrillo, R.; Castiglione, S.; Cicatelli, A.; Guarino, C. Genetic characterization, micropropagation, and potential use for arsenic phytoremediation of *Dittrichia viscosa* (L.) Greuter. *Ecotoxicol. Environ. Saf.* **2018**, *148*, 675–683. [[CrossRef](#)]
214. Jia, Y.; Huang, H.; Sun, G.X.; Zhao, F.J.; Zhu, Y.G. Pathways and relative contributions to arsenic volatilization from rice plants and paddy soil. *Environ. Sci. Technol.* **2012**, *46*, 8090–8096. [[CrossRef](#)]
215. Hussein, H.S.; Ruiz, O.N.; Terry, N.; Daniell, H. Phytoremediation of mercury and organomercurials in chloroplast transgenic plants: Enhanced root uptake, translocation to shoots, and volatilization. *Environ. Sci. Technol.* **2007**, *41*, 8439–8446. [[CrossRef](#)]
216. Cabrita, M.T.; Duarte, B.; Cesário, R.; Mendes, R.; Hintelmann, H.; Eckey, K.; Dimock, B.; Caçador, I.; Canário, J. Mercury mobility and effects in the salt-marsh plant *Halimione portulacoides*: Uptake, transport, and toxicity and tolerance mechanisms. *Sci. Total Environ.* **2019**, *650*, 111–120. [[CrossRef](#)]
217. Battke, F.; Ernst, D.; Fleischmann, F.; Halbach, S. Phytoreduction and volatilization of mercury by ascorbate in *Arabidopsis thaliana*, European beech and Norway spruce. *Appl. Geochemistry* **2008**, *23*, 494–502. [[CrossRef](#)]
218. De Souza, M.P.; Chu, D.; Zhao, M.; Zayed, A.M.; Ruzin, S.E.; Schichnes, D.; Terry, N. Rhizosphere bacteria enhance selenium accumulation and volatilization by indian mustard. *Plant Physiol.* **1999**, *119*, 565–573. [[CrossRef](#)]

219. Terry, N.; Carlson, C.; Raab, T.K.; Zayed, A.M. Rates of Selenium Volatilization among Crop Species. *J. Environ. Qual.* **1992**, *21*, 341–344. [[CrossRef](#)]
220. Pilon-Smits, E.A.H.; De Souza, M.P.; Lytle, C.M.; Shang, C.; Lugo, T.; Terry, N. Selenium volatilization and assimilation by hybrid poplar (*Populus tremula x alba*). *J. Exp. Bot.* **1998**, *49*, 1889–1892. [[CrossRef](#)]
221. Parker, D.R.; Feist, L.J.; Varvel, T.W.; Thomason, D.N.; Zhang, Y. Selenium phytoremediation potential of *Stanleya pinnata*. *Plant Soil* **2003**, *249*, 157–165. [[CrossRef](#)]
222. Zieve, R.; Peterson, P.J. Volatilization of selenium from plants and soils. *Sci. Total Environ.* **1984**, *32*, 197–202. [[CrossRef](#)]
223. Shackira, A.M.; Puthur, J.T. Enhanced phytostabilization of cadmium by a halophyte—*Acanthus ilicifolius* L. *Int. J. Phytoremediat.* **2017**, *19*, 319–326. [[CrossRef](#)]
224. Nandillon, R.; Lebrun, M.; Miard, F.; Gaillard, M.; Sabatier, S.; Morabito, D.; Bourgerie, S. Contrasted tolerance of *Agrostis capillaris* metallicolous and non-metallicolous ecotypes in the context of a mining technosol amended by biochar, compost and iron sulfate. *Environ. Geochem. Health* **2019**, *43*, 1457–1475. [[CrossRef](#)] [[PubMed](#)]
225. Cristaldi, A.; Oliveri Conti, G.; Cosentino, S.L.; Mauromicale, G.; Copat, C.; Grasso, A.; Zuccarello, P.; Fiore, M.; Restuccia, C.; Ferrante, M. Phytoremediation potential of *Arundo donax* (Giant Reed) in contaminated soil by heavy metals. *Environ. Res.* **2020**, *185*, 109427. [[CrossRef](#)]
226. Visconti, D.; Álvarez-Robles, M.J.; Fiorentino, N.; Fagnano, M.; Clemente, R. Use of *Brassica juncea* and *Dactylis glomerata* for the phytostabilization of mine soils amended with compost or biochar. *Chemosphere* **2020**, *260*, 127661. [[CrossRef](#)]
227. Nadgórska-Socha, A.; Ptasieński, B.; Kita, A. Heavy metal bioaccumulation and antioxidative responses in *Cardaminopsis arenosa* and *Plantago lanceolata* leaves from metalliferous and non-metalliferous sites: A field study. *Ecotoxicology* **2013**, *22*, 1422–1434. [[CrossRef](#)]
228. Hussain, S.; Akram, M.; Abbas, G.; Murtaza, B.; Shahid, M.; Shah, N.S.; Bibi, I.; Niazi, N.K. Arsenic tolerance and phytoremediation potential of *Conocarpus erectus* L. and *Populus deltoides* L. *Int. J. Phytoremediat.* **2017**, *19*, 985–991. [[CrossRef](#)]
229. Raza, A.; Khan, A.H.A.; Nawaz, I.; Qu, Z.; Yousaf, S.; Ali, M.A.; Sayal, A.U.; Iqbal, M. Evaluation of Arsenic-Induced Stress in *Dahlia pinnata* Cav.: Morphological and Physiological Response. *Soil Sediment Contam.* **2019**, *28*, 716–728. [[CrossRef](#)]
230. Meier, S.; Alvear, M.; Borie, F.; Aguilera, P.; Ginocchio, R.; Cornejo, P. Influence of copper on root exudate patterns in some metallophytes and agricultural plants. *Ecotoxicol. Environ. Saf.* **2012**, *75*, 8–15. [[CrossRef](#)] [[PubMed](#)]
231. Wu, Q.; Wang, S.; Thangavel, P.; Li, Q.; Zheng, H.; Bai, J.; Qiu, R. Phytostabilization potential of *Jatropha curcas* L. in polymetallic acid mine tailings. *Int. J. Phytoremediat.* **2011**, *13*, 788–804. [[CrossRef](#)] [[PubMed](#)]
232. Santibáñez, C.; Verdugo, C.; Ginocchio, R. Phytostabilization of copper mine tailings with biosolids: Implications for metal uptake and productivity of *Lolium perenne*. *Sci. Total Environ.* **2008**, *395*, 1–10. [[CrossRef](#)] [[PubMed](#)]
233. García, S.; Zornoza, P.; Hernández, L.E.; Esteban, E.; Carpena, R.O. Response of *Lupinus albus* to Pb-EDTA indicates relatively high tolerance. *Toxicol. Environ. Chem.* **2017**, *99*, 1378–1388. [[CrossRef](#)]
234. Zhang, J.; Yang, S.; Huang, Y.; Zhou, S. The Tolerance and Accumulation of *Miscanthus Sacchariflorus* (maxim.) Benth., an Energy Plant Species, to Cadmium. *Int. J. Phytoremediat.* **2015**, *17*, 538–545. [[CrossRef](#)]
235. Pavel, P.B.; Puschenreiter, M.; Wenzel, W.W.; Diacu, E.; Barbu, C.H. Aided phytostabilization using *Miscanthus sinensis* × *giganteus* on heavy metal-contaminated soils. *Sci. Total Environ.* **2014**, *479–480*, 125–131. [[CrossRef](#)]
236. Lee, S.H.; Ji, W.H.; Lee, W.S.; Koo, N.; Koh, I.H.; Kim, M.S.; Park, J.S. Influence of amendments and aided phytostabilization on metal availability and mobility in Pb/Zn mine tailings. *J. Environ. Manag.* **2014**, *139*, 15–21. [[CrossRef](#)]
237. Kiran, B.R.; Prasad, M.N.V. Assisted phytostabilization of Pb-spiked soils amended with charcoal and banana compost and vegetated with *Ricinus communis* L. (Castor bean). *Environ. Geochem. Health* **2021**, *43*, 1507–1521. [[CrossRef](#)]
238. Varun, M.; Souza, R.D.; Pratas, J.; Paul, M.S.; De Ciências, D.; Coimbra, U. De Evaluation of phytostabilization, a green technology to remove heavy metals from industrial sludge using *Typha latifolia* L. *Biotechnol. Bioinf. Bioeng.* **2011**, *1*, 137–145.